

Applications of ultra-cold atoms

Several fundamental experiments can be done with such ultra-cold atomic clouds. One can produce an atomic fountain of such a cloud by projecting it upwards with a small velocity of a few cm/s. This can be done by making the two counter-propagating laser beams in the z direction slightly different in intensity. One can also produce a moving standing wave in the z direction by making the two counter-propagating waves along the z direction to have slightly different frequencies. The projected cloud will reach a certain height and fall back in just the same way as a ball projected upwards under the action of gravity.

Using such an atomic fountain of caesium in a Ramsey type interference experiment with microwave beam, it is possible to measure the clock frequency of Cs more precisely than with existing techniques. The precision of the time standard can be improved⁹ by three orders of magnitude to one part in 10^{16} . Using an atomic fountain of sodium and performing a Ramsey type experiment using optical beams, it has been shown¹¹ that the acceleration due to gravity could be measured to a precision of three parts in 10^8 . Rotational frequencies as small as 5×10^{-13} radians/second can be measured¹².

It is possible to perform experiments to test the foundations of quantum mechanics. One can verify to what accuracy the magnitude of the charge on the electron and proton are equal. Phase

changes introduced by fields on coherent atomic beams can also be measured.

Using laser cooling as the first stage and evaporative cooling as the second stage, Bose-Einstein condensation was first observed¹² in July 1995 in a cloud of two thousand ^{87}Rb atoms cooled below 170 nano Kelvin. Since then the number of atoms in the condensate has been increased to more than a million atoms and the properties of a weakly interacting Bose Condensate of alkali atoms have been measured and compared with theory. In 1997 the coherence of the Bose condensate was demonstrated¹³ by producing interference between two condensates. The condensates were created in a double potential well and allowed to diffuse into each other by removing the confining magnetic field and the blue detuned laser beam. It has also been possible to couple a fraction of the condensate out of the trap making the first atom laser¹⁴. Using lithography with such beams it will be possible to achieve information storage densities far in excess of what has been achieved so far. The possibilities for new applications seem endless.

The phenomenon of laser cooling and trapping has reached the present level of maturity only through the seminal work of Steven Chu, Cohen-Tannoudji and W. D. Phillips. It is therefore appropriate that the Nobel Committee has chosen these three scientists for the award of the Nobel Prize for 1997.

2. Lett, P. D., Watts, R. N., Westbrook, C. I. and Phillips, W. D., *Phys. Rev. Lett.*, 1988, **61**, 169-172.
3. Dalibard, J. and Cohen-Tannoudji, C., *J. Opt. Soc. Am.*, 1989, **B6**, 2023-2045.
4. Salomon, C., Dalibard, J., Phillips, W. D., Clairon, A. and Guellati, S., *Europhys. Lett.*, 1990, **12**, 683-688.
5. Migdall, A. L., Prodan, J. V., Phillips, W. D., Bergmann, T. H. and Metcalf, H. J., *Phys. Rev. Lett.*, 1985, **54**, 2596-2599.
6. Chu, S., Bjorkholm, J. E., Ashkin, A. and Cable, A., *Phys. Rev. Lett.*, 1986, **57**, 314-317.
7. Raab, E. L., Prentiss, M., Cable, A., Chus, S. and Pritchard, D. E., *Phys. Rev. Lett.*, 1987, **59**, 2631-2634.
8. Monroe, C., Swann, W. Robinson, H. and Wieman, C., *Phys. Rev. Lett.*, 1990, **65**, 1571-1574.
9. Clairon, A., Salomon, C., Guellati, S. and Phillips, W. D., *Europhys. Lett.*, 1991, **16**, 165-170.
10. Kasevich, M. and Chus, S., *Appl. Phys.*, 1992, **54**, 321-330.
11. Weiss, D. S., Young, B. N. and Chus, S., *Phys. Rev. Lett.*, 1993, **70**, 2706-2709.
12. Anderson, M. H., Ensher, J. R., Mathews, M. R., Wieman, C. E. and Cornell, E. A., *Science*, 1995, **269**, 198-201.
13. Andrews, M. R., Townsend, C. G., Miesner, H. J., Durfee, D. S., Kurn, D. M. and Ketterle, W., *Science*, 1997, **275**, 637-641.
14. Mewes, M. O., Andrews, M. R., Kurn, D. M., Durfee, D. S., Townsend, C. G. and Ketterle, W., *Phys. Rev. Lett.*, 1997, **78**, 582-585.

1. Chus, S., Hollberg, L., Bjorkholm, J., Cable, A. and Ashkin, A., *Phys. Rev. Lett.*, 1985, **55**, 48-51.

R. Srinivasan is in the Raman Research Institute, C.V. Raman Avenue, Bangalore 560 080, India,

COMMENTARY

The population and conservation status of Asian elephants in the Periyar Tiger Reserve, Southern India

Uma Ramakrishnan, J. A. Santosh, Uma Ramakrishnan and R. Sukumar

Populations of many animal species worldwide are threatened with loss of genetic variation due to substantial reductions in population size'. The genetically effective population size may

also decrease due to another demographic factor - a skewing of the ratio of reproducing males to females². The more unequal the sex ratio, the lower the effective population size. While the

sex ratios of polygynous' mammals are naturally female-biased in most instances due to a variety of reasons, the selective hunting of males would further exacerbate this situation. Among Asian

elephants (*Elephas maximus*), only the males possess tusks (there are also tuskless males) and thus are selectively targeted by poachers for their ivory³, resulting in very skewed sex ratios.

While ivory poaching has been a problem in most southern Indian population—several reports suggested that Periyar Tiger Reserve in Kerala is one of the worst affected⁴. We therefore initiated studies on the status, ecology and behaviour of elephants in Periyar to address several conservation concerns. Among the aspects we have studied are estimates of elephant population numbers, population structure and demography, simulation modelling of tusk harvest by poachers, and interrelationships of skewed sex ratio and changes in social behaviour. In this commentary we would like to summarize the present status of Periyar's elephants, the impact that ivory poaching has made on the population, and its demographic, genetic and conservation implications.

Periyar Tiger Reserve (777 km²) in Kerala state is constituted around a lake, with a waterspread of 26 km², created by a dam constructed across the Periyar river in 1895. The vegetation of the reserve consists of tropical wet evergreen and semi-evergreen forests, interspersed with moist deciduous forests and grasslands. Periyar was amongst the first Project Tiger Reserves declared in 1973, and more recently designated as an Elephant Reserve (No. 10) under India's Project Elephant¹.

A rapid survey was carried out at Periyar during March–May 1994, when density estimates were made and data on population structure obtained. We continued observation on the population structure and demography of elephants from June 1994 to June 1995 and during February–April 1997. A vegetation map of Periyar Tiger Research (PTR)⁶ formed the basis of our survey and stratification of the reserve for the purpose of sampling. The density of elephants was estimated using the line transect, indirect count method described elsewhere^{7,8}. A total of 64 transects was cut in different habitat types, and data analysed separately for each habitat type and also for areas around the lake (i.e. 0.5 km from the lake shore) and away from the lake. The daily defaecation rate of elephants was taken to be 16.3 (CV of 10%) based on data from wild elephants at Mudumalai⁹, while the daily dung decay rate of 0.013/day (CV of 10%) from calculations at Parambikulam Sanctuary¹⁰ where the climate and vegetation are in part similar to Periyar. Dung and elephant densities estimates were calculated for each habitat type using the programs TRANSECT¹¹, and GAJAH which uses a Monte Carlo method described in detail elsewhere¹². Elephants were aged by field techniques for constructing the population age structure³. Population structures were also obtained during another phase of field work during February–April 1997.

The total number of elephants in the sanctuary was found to be 1166 (95% confidence limits of 641–2115) during our 1994 survey. This is comparable to the estimates of 800 elephants (within high use areas alone) during 1978 through indirect count¹³, 935–1100 during 1987–89 by Chandran⁴ and dung densities in Thekkndy Range during the 1993 census by Kerala Forest Research Institute¹⁴.

Using direct elephant sightings, the adult male to female ratio was found to be 1:101 in the present study. Of the 5 adult males recorded, only two were tuskers and the rest were makhnas. It is possible that we may have missed another tusker according to local reports. The time series data on population structure across the various studies during the period 1969–97 clearly brings out a decline in the proportions of males in the population (Table 1). Concurrently, the numbers of males in the population would have also declined in the absence of any significant increase in the total population size. The population size is unlikely to have increased much, at least during the latter half of the above period (1969–97), because of a clear decline in recruitment into the population (discussed below).

Some of the data from previous studies such as Kurup¹⁵ and Chandran⁴ cannot be directly compared with ours because the former do not clearly define the various categories (calf, juvenile, sub-adult and adult) either in terms of

Table 1. Comparison of age structures and sex ratios of elephant populations in southern India

	Female (% population)				Male (% population)			
	Calf	Juvenile	Sub-adult	Adult	Calf	Juvenile	Sub-adult	Adult
Periyar 1969	1.219.0*		51.5	1.219.0*		8.0
Periyar 1978–80**	?	?	?	?	?	?	?	3.1
Periyar 1981–82	7.2	6.6†	8.1	56.7	7.2	6.6†	6.7	0.8
Periyar 1987–898.8†		18.5	60.28.8†		3.3	0.5
Periyar 1994–95	2.1	10.2	25.1	51.9	1.8	4.9	3.5	0.5
Periyar 1997	4.9	8.4	11.6	60.4	4.9	5.6	3.7	0.5
Satyamangalam-Chamarajanagar 1981–83	3.6	10.6	18.5	35.3	3.6	14.3	6.8	7.4
Mudumalai-Bandipur 1994	3.5	10.8	14.5	37.2	3.5	15.7	10.8	4.0

Note that classification into various age categories at Periyar during 1969 (ref. 15), 1978–80, 1981–82 (ref. 13) and 1987–89 (ref. 4) are not directly comparable with the rest of the data (Ref. 3 and our observations). See text for details.

*Combined data for juveniles and sub-adults.

**Data on herd classification not available for Periyar during 1978–80.

†Juveniles have not been sexed during 1981–82. Therefore equal numbers of males and females have been assumed.

*Combined data for calves and juveniles.

age or actual sizes, while we have defined these as calf (<1 yr), juvenile (1–5 yr), sub-adult (5–15 yr) and adult (>15 yr) based on size criteria³. Even if the definitions of elephant categories are not directly comparable, the distinct decline in the juvenile, sub-adult and adult male categories can be seen over this time period.

The ratio of adult male to female seems to have progressively skewed from about 1:6 (1969) to 1:19 (1977–79) and then skewed substantially to 1:71 (1980–82) reaching a peak of 1:122 (1987–89). It is important to remember that skewing of the adult sex ratio will proceed at a geometric rate with the progressive death of one sex, and thus the death of one or a few adult males will skew the ratio substantially as their numbers reduce in the population.

Interestingly, both the sub-adult and males categories seem to have reached very low frequencies even by 1987–89 when Chandran⁴ studied this population, and have not changed much over the next five years. These data indicate that ivory poaching began during the 1970s itself, with the adult male segment being targeted and then moved down to the sub-adult category by the 1980s. Severe poaching is indicated during the period prior to 1987 when even juvenile males have not been spared. Computer simulation modelling of elephant demography indicates that 336–388 tuskers of various ages have been killed and 3256–3334 kg of tusks harvested by poachers over the period 1974–94 (ref. 16).

Another demographic effect of the high rates of poaching and extreme skew in adult sex ratios is a decline in the birth rate of the population. Our 1994–95 data indicated that the average fecundity was only 0.075/adult female/year as compared to between 0.20 and 0.25 for the more productive populations in southern India³. The most likely cause of the fecundity decline is an Allee effect¹⁷. With an extremely low number of adult males in the populations, a substantial proportion of females coming into estrus would not be detected and mated. Another possible reason for the low numbers of calves and juveniles seen at Periyar is that females may have stopped reproducing under conditions of extreme stress

(poaching), or infant mortality may have been very high, as may have happened in an African elephant population at Ruaha, Tanzania¹⁸. The decline in recruitment into the population is not because of inter-annual variations in births as seen elsewhere in the south³ but a phenomenon which has persisted for several years as seen from the low numbers of juveniles in the 1994–95 age structure. Our 1997 observations, however, showed an increase in births with a fecundity rate of 0.16/adult female/year, still lower than the potential rate. It is possible that 1997 represents a genuine upswing in reproductive rates or that our limited sampling during three dry months represents only a single clan at Periyar.

Another aspect of reproduction at Periyar that has been observed is the presence of several barren adult females. Such female elephants show no sign of mammary gland activity and, further, have retracted nipples indicating that they have not reproduced or may be incapable of reproducing. This has also been noted by another experienced elephant observer (D. K. Lahiri-Choudhury, pers. commun.). It is possible that an alteration of reproductive physiology in female elephants is related to the near absence of adult males in the population over an extended period of time. This aspect requires further study, possibly through monitoring metabolites of reproductive hormones in faeces¹⁹.

The present situation of the elephant population at Periyar clearly calls for further research and active management. The elephant population here can possibly withstand the drastic reduction in the genetically effective population size (N_e), because this reduction has occurred over a single elephant generation (about 20 years). In theory, a one generation reduction of N_e to 20+ still retains about 98% of the original (prior to the poaching spate) heterozygosity, assuming random sampling of individuals. Allelic diversity could, however, be expected to suffer greater decline, with rare alleles having a high probability of being lost during a population bottleneck²⁰. Obviously the poaching of male elephants has been selective, at least in the initial years, and thus the genetic erosion could have been more serious²¹.

Management options for Periyar would include the translocation of adult male elephants from other populations with surpluses, in order to increase the genetically effective population size to 50 or above. This could be achieved by increasing the number of adult males from the present 6 to 13 individuals. Apart from increasing N_e , this would also serve to stimulate the birth rate of the population. Several questions would arise about such a proposed translocation. First of all, there is the question of protection of translocated bulls from poachers. From this angle, it may be better to translocate tuskless bulls (makhnas) into Periyar, given that ivory poaching is likely to be a continuing problem²². If breeding makhnas are translocated, the frequency of alleles for tusklessness would increase. Available data on captive elephant breeding (RS, unpublished) indicate that females are also carriers of the genes for tusks. Among sons born to females mated by a makhna, a certain proportion would thus still be tuskers (heterozygous genotype). It may thus be prudent to translocate makhnas or a mixture of makhnas and tuskers until protection from poachers is ensured.

The next aspect to be considered is that of the genetic consequences of introducing males from a population (such as Assam, where surplus makhnas are most likely to be available) which has been isolated from Periyar for a long period of time. Related to this is the health (parasite and disease) status of translocated bulls. Could elephants in Periyar have evolved resistance to certain local parasites against which elephants from another population may not have resistance? Could the translocation of such bulls disrupt locally-adapted gene complexes at Periyar? This calls for basic studies on patterns of genetic variation and levels of divergence among the elephant populations at Periyar and elsewhere. Concurrently, there have to be studies on the parasite and health status of elephants in these populations. If bulls are translocated into Periyar, it would be advisable to do this in stages after careful monitoring of these bulls through radio-telemetry to study breeding and possible conflict with people. It is time that such an elephant management programme is initiated at Periyar. The current status of

this important elephant population warrants immediate management action.

1. Advise, J. C. and Hamrick, J. L., *Conservation Genetics: Case Histories from Nature*, Chapman and Hall, New York, 1996.
2. Harris, R. B. and Allendorf, F. W., *Conserv. Biol.*, 1989, **3**, 181–191.
3. Sukumar, R., *The Asian Elephant: Ecology and Management*, Cambridge University Press, New York, 1989.
4. Chandran, P. M., Proceedings of the Elephant Symposium, Kerala Forest Department, Trivandrum, 1990, pp. 51–56.
5. Anonymous, *Project Eleplzant (Gajalme)*, Ministry of Environment and Forest, Govt. of India, 1993, p. 46.
6. Chandrasekharan, C., *Vegetation Map of Kerala*, Forest Department, Trivandrum, 1973.
7. Barnes, R. F. W. and Jensen, K. L., *IUCN African Elephant and Rhino Specialist Group Tech. Bull.*, 1987.
8. Varman, K. S., Ramakrishnan, U. and Sukumar, R., in *A Week with Elephants* (eds Daniel, J. C. and Datye, H.), Bombay Natural History Society, Bombay and Oxford University Press, New Delhi, 1995, pp. 331–339.
9. Watve, M. G., Ph D. thesis, Indian Institute of Science, Bangalore, 1992.
10. Dawson, S., M Sc thesis, Department of Zoology, University of Oxford, 1990.
11. White, G. C., *Program TRANSECT – Line transect Data Analysing Program*, Colorado State University, Fort Collins, 1987.
12. Santosh, J. A. and Sukumar, R., in *A Week with Elephants* (eds Daniel, J. C. and Datye, H.), Bombay Natural History Society, Bombay and Oxford University Press, New Delhi, 1995, pp. 394–404.
13. Nair, P. V., Ramachandran, K. K., Vijayan, V. S., Easa, P. S. and Balakrishnan, P. V., Kerala Forest Research Institute Report-24, Peechi, Kerala, 1985, p. 159.
14. Anonymous, *Wildlife Census – Kerala 1993, A Report*, Kerala Forest Research Institute, Peechi, Trichur, 1993, p. 129.
15. Kurup, G. U., *Cheetal*, 1971, **13**, 5–18.
16. Sukumar, R., Ramakrishnan, U. and Santosh, J. A., Ms in review.
17. Dennis, B., *Natural Resource Modeling*, 1989, **3**, 481–538.
18. Barnes, R. F. W. and Kapela, E. B., *Afr. J. Ecol.*, 1991, **29**, 289–294.
19. Lasley, B. L. and Kirkpatrick, J. F., *J. Zoo Anim. Wildlife Med.*, 1991, **22**, 23–31.
20. Franklin, I. R., in *Conservation Biology: An Evolutionary–Ecological Perspective* (eds Soulé, M. E. and Wilcox, B. A.), Sinauer Associates, Sunderland, Massachusetts, 1980, pp. 135–149.
21. Watve, M. G. and Sukumar, R., *Curr. Sci.*, 1997, **72**, 885–888.
22. Menon, V., Sukumar, R. and Kumar, A *God in Distress: Threats of Poaching and the Ivory Trade to the Asian Elephant in India*, Asian Elephant Conservation Centre, Bangalore and Wildlife Protection Society of India, New Delhi, 1997, p. 91.

ACKNOWLEDGEMENTS. We thank the Forest Department of Kerala for facilities and research permissions. This study was funded by the Ministry of Environment and Forests, Government of India. The preliminary survey was funded by the Calgary Zoological Society and Wildlife Preservation Trust International through the IUCN/SSC Asian Elephant Specialist Group.

The authors are from the Asian Elephant Conservation Centre and Centre for Ecological Sciences, Indian Institute of Science, Bangalore 560 012, India.

SCIENTIFIC CORRESPONDENCE

Typical liptinitic bodies from Neyveli lignite deposits, India

Coal and lignite, the sedimentary organic deposits consist of various micro-constituents called 'macerals' (= 'minerals' in inorganic rocks) which originate during the 'coalification' process, that is transformation of vegetal matter into coal^{1,2}. Different maceral groups are vitrinite – originated from woody plant remains, liptinite – derived from hydrogen-rich plant parts (pollen-spores, cuticles, resins, suberin, etc.), and inertinite – incorporating the carbon-rich oxidized plant remains, show different behaviour under normal and fluorescence modes'. Relatively hydrogen-rich liptinitic macerals fluoresce under fluorescence mode with wide range of colours. Based on morphological and optical properties, it is possible to correlate various macerals to their original plant parts and to reconstruct the palaeoflora^{1,2}.

Besides traditional macerals, typical multicellular bodies emitting the light of yellow and brownish-yellow colours have been found during the investiga-




Figure 1. Location map of Neyveli Lignite field and investigated bore-holes NLE-27, 35 and 36 in Mine III. (Source: Neyveli Lignite Corporation Limited).