

ment lowered R to 0.146. Scattering factors for non-hydrogen atoms were from Cromer & Waber (1965). Refinement of all non-hydrogen atoms with anisotropic temperature factors yielded $R = 0.096$. The positions of the 20 H atoms could be obtained from a difference map but all the H atoms attached to C atoms were fixed with $C-H = 1.1 \text{ \AA}$. Bond angles of 109.5° or 120.0° were used for tetrahedral and trigonal atoms, respectively. For H bonded to N atoms, $N-H = 1.0 \text{ \AA}$ and a bond angle of 120.0° were used. The H atoms were assigned the temperature factor of the carrier atom. With the scattering factor for H given by Stewart, Davidson & Simpson (1965), the refinement of positional and anisotropic thermal parameters of non-hydrogen atoms and positional and isotropic thermal parameters of H atoms yielded an R of 0.069. With the weighting scheme of Cruickshank (1961) further refinement converged to a final R of 0.053 for 2643 reflections. The final difference map was featureless. The shifts in the parameters at the end of the last cycle were $<0.1\text{a}$. The atomic parameters are given in Tables 1 and 2.*

* Lists of structure factors and anisotropic thermal parameters have been deposited with the British Library Lending Division as Supplementary Publication No. SUP 34782 (16 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH 1 2HU, England.

Table 1. Positional coordinates for the non-hydrogen atoms ($\times 10^4$)

	x	y	z
C(1)	2847 (3)	4976 (3)	-78 (2)
C(2)	3677 (3)	5085 (3)	-1050 (2)
C(3)	4812 (4)	4235 (4)	-1234 (2)
C(4)	5094 (4)	3281 (4)	-466 (3)
C(5)	4286 (3)	3180 (0)	513 (2)
C(6)	3140 (3)	3995 (3)	706 (2)
C(7)	2164 (3)	3816 (3)	1709 (2)
O(1)	3062 (2)	3961 (2)	2847 (1)
C(8)	3202 (2)	5138 (2)	3236 (2)
O(2)	2605 (2)	6018 (2)	2719 (1)
N(1)	4042 (2)	5185 (2)	4289 (1)
C(9)	4623 (2)	6361 (2)	4786 (2)
C(10)	5792 (3)	6908 (3)	4020 (2)
C(11)	5367 (3)	6089 (2)	6039 (2)
C(12)	3327 (2)	7282 (2)	4928 (2)
O(3)	3543 (2)	8395 (2)	4785 (2)
N(2)	2021 (2)	6827 (2)	5251 (1)
C(13)	715 (2)	7610 (2)	5499 (2)
C(14)	1183 (3)	8583 (3)	6445 (2)
C(15)	-477 (3)	6762 (3)	5957 (2)
C(16)	-17 (2)	8231 (2)	4365 (2)
O(4)	-938 (2)	9087 (2)	4433 (2)
N(3)	293 (2)	7735 (2)	3347 (2)
C(17)	-273 (3)	8250 (3)	2209 (2)
C(18)	933 (4)	9008 (4)	1678 (3)
C(19)	-845 (3)	7184 (3)	1393 (2)
O(5)	-415 (3)	7012 (4)	445 (2)
O(6)	-1880 (3)	6519 (3)	1853 (2)
C(20)	-2540 (6)	5507 (5)	1147 (4)

E.s.d.'s are given in parentheses.

Table 2. Positional ($\times 10^3$) and isotropic thermal ($\times 10$) parameters of the hydrogen atoms

E.s.d.'s are given in parentheses.

Bonded to	x	y	z	$B (\text{\AA}^2)$	
H(1)	C(1)	207 (5)	556 (5)	6 (4)	57 (9)
H(2)	C(2)	331 (7)	565 (7)	-170 (5)	86 (15)
H(3)	C(3)	551 (4)	428 (4)	-184 (3)	39 (8)
H(4)	C(4)	600 (5)	275 (6)	-72 (4)	62 (13)
H(5)	C(5)	457 (6)	244 (6)	118 (5)	70 (12)
H(6)	C(7)	121 (4)	440 (5)	165 (3)	47 (9)
H(7)	C(7)	173 (5)	286 (6)	180 (4)	68 (10)
H(8)	N(1)	431 (4)	456 (4)	446 (3)	27 (7)
H(9)	C(10)	517 (5)	706 (6)	300 (4)	66 (11)
H(10)	C(10)	628 (4)	778 (5)	429 (3)	50 (9)
H(11)	C(10)	663 (4)	635 (4)	399 (3)	36 (7)
H(12)	C(11)	619 (4)	555 (5)	608 (3)	43 (9)
H(13)	C(11)	583 (5)	685 (6)	639 (4)	44 (10)
H(14)	C(11)	453 (3)	569 (4)	655 (3)	30 (6)
H(15)	N(2)	216 (6)	626 (6)	553 (4)	68 (12)
H(16)	C(14)	204 (4)	900 (4)	632 (3)	40 (8)
H(17)	C(14)	2 (5)	913 (5)	657 (4)	57 (10)
H(18)	C(14)	183 (8)	824 (9)	721 (6)	125 (18)
H(19)	C(15)	-22 (4)	641 (4)	672 (3)	38 (7)
H(20)	C(15)	-140 (5)	713 (6)	640 (4)	60 (10)
H(21)	C(15)	-73 (5)	627 (5)	550 (3)	39 (8)
H(22)	N(3)	97 (4)	717 (4)	337 (3)	35 (8)
H(23)	C(17)	-64 (9)	904 (10)	844 (6)	112 (24)
H(24)	C(18)	197 (5)	837 (6)	151 (4)	53 (10)
H(25)	C(18)	62 (5)	953 (5)	95 (4)	49 (9)
H(26)	C(18)	121 (6)	965 (7)	229 (5)	86 (14)
H(27)	C(20)	-175 (7)	481 (8)	94 (5)	92 (15)
H(28)	C(20)	-328 (6)	516 (7)	158 (5)	90 (13)
H(29)	C(20)	-289 (6)	592 (6)	36 (4)	77 (12)

Results and discussion

A perspective view of the molecule down a is shown in Fig. 1. Bond lengths and angles are listed in Table 3. The bond lengths and angles of the peptide units




Fig. 1. Molecular conformation of Z-Aib-Aib-Ala-OME viewed down a .

Table 3. Bond distances (Å) and angles (°) with *e.s.d.*'s in parentheses

C(1)–C(2)	1.395 (4)	C(12)–N(2)	1.339 (3)
C(2)–C(3)	1.392 (5)	C(12)–O(3)	1.233 (3)
C(3)–C(4)	1.361 (5)	N(2)–C(13)	1.481 (3)
C(4)–C(5)	1.389 (4)	C(13)–C(14)	1.535 (4)
C(5)–C(6)	1.377 (4)	C(13)–C(15)	1.528 (4)
C(6)–C(1)	1.396 (4)	C(13)–C(16)	1.543 (3)
C(6)–C(7)	1.513 (3)	C(16)–O(4)	1.240 (3)
C(7)–O(1)	1.462 (3)	C(16)–N(3)	1.333 (3)
O(1)–C(8)	1.350 (3)	N(3)–C(17)	1.456 (3)
C(8)–O(2)	1.213 (3)	C(17)–C(18)	1.518 (5)
C(8)–N(1)	1.350 (3)	C(17)–C(19)	1.536 (4)
N(1)–C(9)	1.465 (3)	C(19)–O(5)	1.196 (4)
C(9)–C(10)	1.536 (3)	C(19)–O(6)	1.313 (4)
C(9)–C(11)	1.543 (3)	O(6)–C(20)	1.448 (6)
C(9)–C(12)	1.540 (3)		
C(1)–C(2)–C(3)	120.6 (3)	O(3)–C(12)–N(2)	123.0 (2)
C(2)–C(3)–C(4)	119.5 (3)	C(12)–N(2)–C(13)	123.5 (2)
C(3)–C(4)–C(5)	120.1 (3)	O(5)–C(19)–O(6)	124.9 (3)
C(4)–C(5)–C(6)	121.6 (3)	C(17)–C(19)–O(6)	111.7 (3)
C(5)–C(6)–C(1)	118.5 (2)	N(2)–C(13)–C(14)	111.4 (2)
C(6)–C(1)–C(2)	119.7 (3)	N(2)–C(13)–C(15)	107.4 (2)
C(5)–C(6)–C(7)	121.5 (2)	N(2)–C(13)–C(16)	111.0 (2)
C(1)–C(6)–C(7)	119.9 (2)	C(15)–C(13)–C(14)	108.6 (2)
C(6)–C(7)–O(1)	110.9 (2)	C(15)–C(13)–C(16)	107.6 (2)
C(7)–O(1)–C(8)	114.7 (2)	C(14)–C(13)–C(16)	110.6 (2)
O(1)–C(8)–O(2)	124.0 (2)	C(13)–C(16)–N(3)	116.7 (2)
O(1)–C(8)–N(1)	110.7 (2)	C(13)–C(16)–O(4)	119.7 (2)
C(8)–N(1)–C(9)	121.2 (2)	O(4)–C(16)–N(3)	123.3 (2)
O(2)–C(8)–N(1)	125.3 (2)	C(16)–N(3)–C(17)	122.8 (2)
N(1)–C(9)–C(10)	110.2 (2)	N(3)–C(17)–C(19)	108.5 (2)
N(1)–C(9)–C(11)	107.1 (2)	N(3)–C(17)–C(18)	111.8 (3)
N(1)–C(9)–C(12)	111.5 (2)	C(18)–C(17)–C(19)	111.4 (3)
C(10)–C(9)–C(11)	110.6 (2)	C(17)–C(19)–O(5)	123.4 (3)
C(10)–C(9)–C(12)	111.1 (2)	C(19)–O(6)–C(20)	116.7 (3)
C(11)–C(9)–C(12)	106.2 (2)		
C(9)–C(12)–N(2)	117.5 (2)		
C(9)–C(12)–O(3)	119.4 (2)		

compare well with those found in other peptides (Marsh & Donohue, 1967; Ramachandran, Kolaskar, Ramakrishnan & Sasisekharan, 1974; Benedetti, 1977). The average C–H = 1.0 Å and N–H = 0.8 Å. The angles involving H with tetrahedral C atoms are around 109° on average. There are few intramolecular contacts less than the normal limits given by Ramachandran & Sasisekharan (1968). The contacts C(7)–O(2) of 2.659 (4) Å and C(20)–O(5) of 2.668 (6) Å are less than the proposed extreme limit of 2.7 Å.

The molecular conformation is stabilized by an internal hydrogen bond between the CO of the urethan moiety and the NH of the alanyl residue. This conformation with a 4→1 type of intramolecular hydrogen bond, which corresponds to a type III' β -bend (Venkatachalam, 1968), has also been seen earlier in oligopeptides containing Aib residues (Shamala, Nagaraj & Balaram, 1977, 1978; Prasad, Shamala, Nagaraj, Chandrasekaran & Balaram, 1979; Smith,

Duax, Czerwinski, Kendrick, Marshall & Mathews, 1977). The conformational angles of the backbone are listed in Table 4. All the peptide units are *trans* and only small deviations of about 5° from planarity are observed.

Theoretical calculations suggest that the additional methyl substituent at C α of the Aib residue greatly restricts the allowed values of φ and ψ to the right- and left-handed 3_{10} - and α -helical regions (Marshall & Bosshard, 1972; Burgess & Leach, 1973). The values in Table 5 show that both the Aib residues occur in the left-handed helical region of the conformational map. In peptides containing L amino acids like *Z*-Aib-Pro-Aib-Ala-OMe (Nagaraj *et al.*, 1979), Boc-Pro-Aib-Ala-Aib-OMe (Smith *et al.*, 1977) and *Z*-Aib-Pro-NHMe (Prasad *et al.*, 1979) the φ , ψ values for the Aib residues fall in the right-handed helical region. However, in tosyl-(Aib)₅-OMe (Shamala *et al.*, 1977) both helical senses are equally probable as observed in the crystal, which belongs to a centrosymmetric space group. It is clear that the presence of an Aib residue

Table 4. Conformational angles (°) for the peptide backbone according to IUPAC–IUB Commission on Biochemical Nomenclature (1970)

ω_1 [O(1)–C(8)–N(1)–C(9)]	167.2 (2)
φ_2 [C(8)–N(1)–C(9)–C(12)]	58.1 (2)
ψ_2 [N(1)–C(9)–C(12)–N(2)]	36.8 (2)
ω_2 [C(9)–C(12)–N(2)–C(13)]	175.8 (2)
φ_3 [C(12)–N(2)–C(13)–C(16)]	68.3 (2)
ψ_3 [N(2)–C(13)–C(16)–N(4)]	18.6 (3)
ω_3 [C(13)–C(16)–N(4)–C(17)]	–177.7 (2)
φ_4 [C(16)–N(4)–C(17)–C(18)]	–136.2 (2)


Fig. 2. Packing diagram viewed down *b* (--- intermolecular hydrogen bonds; - - - intramolecular hydrogen bonds).

Table 5. Details of the hydrogen bonds

Donor <i>D</i>	Acceptor <i>A</i>	<i>D</i> ... <i>A</i>	$\angle D-H \cdots A$	$\angle H-D \cdots A$
N(3)	O(2)*	2.904 (7) Å	156.9 (25)°	16.5 (20)°
N(1)	O(3) [†]	2.997 (7)	144.5 (24)	27.2 (21)
N(2)	O(4) ^{††}	3.147 (8)	138.7 (24)	32.9 (22)

Symmetry code: superscript: (i) $1 - x, 1 - (y + \frac{1}{2}), 1 - z$; (ii) $-x, 1 - (y + f), 1 - z$.

* Intramolecular hydrogen bond.

induces β -bend formation and the consequent generation of a 3₁-helical segment.

The crystal structure viewed along **b** is shown in Fig. 2. In addition to the good intramolecular NH...O hydrogen bond between the CO of the urethan moiety and NH of the alanyl residue, the other peptide NH and CO groups are involved in intermolecular hydrogen bonds. The details of the inter- and intramolecular hydrogen bonds are given in Table 5.

This research was supported by the University Grants Commission. NS thanks the CSIR for financial support. RN was the recipient of a fellowship from the Department of Atomic Energy.

References

- BENEDETTI, E. (1977). *Peptides: Proceedings of the Fifth American Peptide Symposium*, edited by M. GOODMAN & J. MEINHOFER, pp. 257–273. New York: John Wiley.
- BURGESS, A. W. & LEACH, S. J. (1973). *Biopolymers*, **12**, 2599–2605.
- CROMER, D. T. & WABER, J. T. (1965). *Acta Cryst.* **18**, 104–109.

- CRUICKSHANK, D. W. J. (1961). In *Computing Methods and the Phase Problem in X-ray Crystal Analysis*. Oxford: Pergamon Press.
- GERMAIN, G., MAIN, P. & WOOLFSON, M. M. (1971). *Acta Cryst.* **A27**, 368–376.
- IUPAC-IUB COMMISSION ON BIOCHEMICAL NOMENCLATURE (1970). *J. Mol. Biol.* **52**, 1–17.
- KARLE, J. & HAUPTMAN, H. (1956). *Acta Cryst.* **9**, 635–651.
- KARLE, J. & KARLE, I. L. (1966). *Acta Cryst.* **21**, 849–859.
- MARSH, R. E. & DONOHUE, J. (1967). *Adv. Protein Chem.* **22**, 235–255.
- MARSHALL, G. R. & BOSSHARD, H. R. (1972). *Circ. Res. Suppl.* **II**, 30–31, 143–150.
- MARTIN, D. R. & WILLIAMS, R. J. P. (1976). *Biochem. J.* **153**, 181–190.
- NAGARAJ, R., SHAMALA, N. & BALARAM, P. (1979). *J. Am. Chem. Soc.* **101**, 16–20.
- PANDEY, R. C., CARTER COOK, J. JR & RINEHART, K. L. JR (1977). *J. Am. Chem. Soc.* **99**, 8469–8483.
- PRASAD, B. V. V., SHAMALA, N., NAGARAJ, R., CHANDRASEKARAN, R. & BALARAM, P. (1979). *Biopolymers*, **18**, 1635–1646.
- RAMACHANDRAN, G. N., KOLASKAR, A. S., RAMAKRISHNAN, C. & SASISEKHARAN, V. (1974). *Biochem. Biophys. Acta*, **359**, 298–302.
- RAMACHANDRAN, G. N. & SASISEKHARAN, V. (1968). *Adv. Protein Chem.* **23**, 283–437.
- SHAMALA, N., NAGARAJ, R. & BALARAM, P. (1977). *Biochem. Biophys. Res. Commun.* **79**, 292–298.
- SHAMALA, N., NAGARAJ, R. & BALARAM, P. (1978). *Chem. Commun.* pp. 996–997.
- SMITH, G. D., DUAX, W. L., CZERWINSKI, E. W., KENDRICK, N. E., MARSHALL, G. R. & MATHEWS, F. S. (1977). *Peptides: Proceedings of the Fifth American Peptide Symposium*, edited by M. GOODMAN & J. MEINHOFER, pp. 277–279. New York: John Wiley.
- STEWART, R. F., DAVIDSON, E. R. & SIMPSON, W. T. (1965). *J. Chem. Phys.* **42**, 3175–3187.
- VENKATACHALAM, C. M. (1968). *Biopolymers*, **6**, 1425–1436.
- WILSON, A. J. C. (1942). *Nature (London)*, **150**, 151–152.

Acta Cryst. (1980). **B36**, 110–114

The Crystal Structure of (+)-2-Dipropylamino-5-hydroxytetralin Hydrochloride

BY JOHAN GIESECKE

Department of Medical Biophysics, Karolinska Institute, S-104 01 Stockholm, Sweden

(Received 2 January 1979; accepted 30 August 1979)

Abstract

The title compound, C₁₆H₂₆NO⁺.Cl⁻, which is a potent dopaminergic drug, is monoclinic with $a = 7.601$ (1), $b = 18.508$ (3), $c = 11.408$ (2) Å, $\beta = 94.28$ (1)°, space group $P2_1$ and $Z = 4$. The structure was solved by

direct methods and refined to $R = 0.052$ for 2081 counter-measured observed reflections. The structure consists of infinite chains along the b axis, the molecules being connected by hydrogen-bonded chloride ions. Adjacent chains are held together by van de Waals forces only. The non-aromatic ring assumes a

0567-7408/80/010110-05\$01.00

© 1980 International Union of Crystallography