Iodide as Nucleophilic Trigger in Aryne Three-Component Coupling for the Synthesis of 2-Iodobenzyl Alcohols

Subrata Bhattacharjee, Avishek Guin, Rahul N. Gaykar, and Akkattu T. Biju*

Department of Organic Chemistry, Indian Institute of Science, Bangalore-560012, India

atbiju@iisc.ac.in

Supporting Information

1. General Information	S02
2. General Procedure for the Optimization of Reaction Conditions	S02
3. General Procedure for the Halide Triggered Aryne MCC with Aldehydes	S05
4. Procedure for the Iodide triggered aryne MCC with <i>N</i> -Methylisatin	S07
5. Procedure for the Iodide Triggered Aryne MCC with CO ₂	S07
6. Mechanistic Experiments	S08
7. X-ray Data of 3ac and 8a	S10
8. Synthesis and Characterization of 2-Halo Benzyl Alcohol Derivatives	S13
9. Functionalization of 2-Iodobenzyl Alcohols	S35
10. ¹ H and ¹³ C NMR Spectra of Products	S40

1. General Information

Unless otherwise specified, all reactions were carried out under an atmosphere of nitrogen in flame-dried reaction vessels with Teflon screw caps. 25 °C Corresponds to the room temperature (rt) of the lab when the experiments were carried out. THF was freshly purified by distillation over Na-benzophenone and was transferred under nitrogen. 18-Crown-6 was recrystallized from dry CH₃CN and KF, KI, KBr and KCl were dried by heating at 110 °C for 12 h and left to cool under nitrogen and stored in nitrogen filled glove-box. The 2(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** and the other symmetric and unsymmetrical aryne precursors were synthesized following literature procedure.¹

Analytical thin layer chromatography was performed on TLC Silica gel 60 F₂₅₄. Visualization was accomplished with short wave UV light or KMnO₄ staining solutions followed by heating. Flash chromatography was performed on silica gel (230-400 mesh) by standard techniques eluting with Pet. Ether-EtOAc solvent system.

All compounds were fully characterized. ¹H and ¹³C NMR spectra were recorded on Bruker Ultrashield spectrometer in CDCl₃ as solvent. Chemical shifts (δ) are given in ppm. The residual solvent signals were used as references and the chemical shifts converted to the TMS scale (CDCl₃: δ_H = 7.26 ppm, δ_C = 77.16 ppm). Infrared (FT-IR) spectra were recorded on a Perkin Elmer Spectrum BX spectrophotometer, ν-max in cm⁻¹. HRMS (ESI) data were recorded on a Waters Xevo G2-XS Q-TOF instrument.

2. General Procedure for the Optimization of the Reaction Conditions

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KI, KF and 18-crown-6 in a glove-box. Then aldehyde **1a** (0.25 mmol) was added outside the

¹ (a) Sato, Y.; Tamura, T.; Kinbara, A.; Morib, M *Adv. Synth. Catal.* **2007**, 349, 647. (b) Peña, D.; Cobas, A.; Pérez, D.; Guitián, E. *Synthesis*, **2002**, 1454.

glove-box under nitrogen atmosphere followed by addition of THF. To the stirring solution, aryne precursor **2a** was added and reaction mixture was allowed to stir at 25 °C for 24 h. After 24 h, the reaction was quenched by adding 4N HCl (HCl in dioxane, 0.4 mL) and the solvent was evaporated. Yields were determined by the ¹H NMR analysis of the crude reaction products using CH₂Br₂ as the internal standard.

Variation of the amount of Aryne precursor (2a)

entry	2a (equiv)	F ⁻ Source (equiv)	Yield ^a of 3a
1	1.3	KF (2.6), 18-crown-6 (2.6)	52%
2	1.5	KF (3), 18-crown-6 (3)	61%
3	2	KF (4), 18-crown-6 (4)	64%
4	2.5	KF (5), 18-crown-6 (5)	67%
5 ^b	2	CsF	18%
6	2	TBAF	<5%

Reaction condition: **1a** (0.25 mmol), KI (2 equiv), KF (THF (2.0 mL), 25 °C for 24 h. ^a Yields based on the ¹H NMR analysis of the crude reaction products using CH₂Br₂ as the internal standard. ^b Reaction was performed in 2 mL CH₃CN

Variation of Iodide source

entry	Iodide Source (equiv)	Yield ^a of 3a
1	KI (1.2 equiv)	32%
2	KI (2 equiv)	64%
3	KI (3 equiv)	61%
4	NaI (2 equiv)	<5%
5	TBAI (2 equiv)	37%

Reaction condition: **1a** (0.25 mmol), **2a** (0.50 mmol), KF (4 equiv), 18-crown-6 (4 equiv), THF (2.0 mL), 25 °C for 24 h. ^a Yields based on the ¹H NMR analysis of the crude reaction products using CH₂Br₂ as the internal standard.

Variation of Temperature

entry	Temperature (°C)	Yield ^a of 3a
1	0 °C	58%
2	rt (25 °C)	64%
3	40 °C	54%
4	50 °C	49%

Reaction condition: **1a** (0.25 mmol), **2a** (0.50 mmol), KI (2 equiv), KF (4 equiv), 18-crown-6 (4 equiv), THF (2.0 mL), for 24 h. ^a Yields based on the ¹H NMR analysis of the crude reaction products using CH₂Br₂ as the internal standard.

Further Optimizations

entry	2a (equiv)	F ⁻ Source (equiv)	Solvent (mL)	KI (equiv)	Yield ^a of 3a
1	2	KF (4), 18-crown-6 (4)	THF (2 mL)	2 equiv	64%
2	2	KF (4), 18-crown-6 (4)	DME (2 mL)	2 equiv	53%
3	2	KF (4), 18-crown-6 (4)	1,4-Dioxane (2 mL)	2 equiv	40%
4	2	KF (4), 18-crown-6 (4)	THF (1 mL)	2 equiv	67%
5	2.5	KF (5), 18-crown-6 (5)	THF (1 mL)	2 equiv	72%
6	2.5	KF (5), 18-crown-6 (5)	THF (1 mL)	2.5 equiv	73% (70)

Reaction condition: **1a** (0.25 mmol), for 24 h. ^a Yields based on the ¹H NMR analysis of the crude reaction products using CH₂Br₂ as the internal standard. Isolated yield in parenthesis.

3. General Procedure for the Halide Triggered Aryne MCC with Aldehydes

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KX (1.25 mmol), KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in a glove box. Then aldehyde **1** (0.5 mmol) was added outside the glove box under nitrogen atmosphere followed by addition of THF (2 mL). To the stirring solution, aryne precursor **2** (2.5 equiv) was added and reaction mixture was allowed to stir at rt for 24 h. After 24 h the reaction was quenched by adding 4N HCl (HCl in dioxane, 0.8 mL), the solvent was evaporated and the crude residue pre-adsorbed on silica gel and purified by flash column chromatography (Pet. ether /EtOAc = 90/10) on silica gel to afford the corresponding 2-halo benzyl alcohol derivatives **3** in moderate to good yields.

Procedure for the 3.0 mmol Scale Reaction for the synthesis of 3a

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KI (1.245 g, 7.5 mmol), KF (0.870 g, 15.0 mmol) and 18-crown-6 (3.960 g, 15.0 mmol) in a glove-box. Then 4-nitrobenzaldehyde **1a** (0.453 g, 3.0 mmol) was added outside the glove-box under nitrogen atmosphere followed by addition of THF (12 mL). To the stirring solution, 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (2.238 g, 7.5 mmol) was added and reaction mixture was allowed to stir at rt for 36 h. After 36 h, the reaction was quenched by adding 4N HCl (HCl in dioxane, 4.8 mL) at 0 °C, the solvent was evaporated and the crude residue was purified by flash column chromatography (Pet. ether /EtOAc = 90/10) on silica gel to afford (2-iodophenyl)(4-nitrophenyl)methanol **3a** as a yellow solid (0.733 g, 69 % yield).

Procedure for the 3.0 mmol Scale Reaction for the synthesis of 3i

Following the above procedure, treatment of methyl 4-formylbenzoate **1i** (0.493 g, 3.0 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (2.238 g, 7.5 mmol) and KI (1.245 g, 7.5 mmol) in the presence of KF (0.870 g, 15.0 mmol) and 18-crown-6 (3.960 g, 15.0 mmol) in THF (12.0 mL) at 25 °C for 36 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 4.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded methyl-4-(hydroxy(2-iodophenyl)methyl)benzoate **3i** as a yellow oil (0.762 g, 69% yield).

Procedure for the 3.0 mmol Scale Reaction for the synthesis of 3y

Following the procedure, treatment of 4-nitrobenzaldehyde **1a** (0.453 g, 3.0 mmol), 6-(trimethylsilyl)benzo[d][1,3]dioxol-5-yl trifluoromethanesulfonate **2c** (1.027 g, 7.5 mmol) and KI (1.245 g, 7.5 mmol) in the presence of KF (0.870 g, 15.0 mmol) and 18-crown-6 (3.960 g, 15.0 mmol) in THF (12.0 mL) at 25 °C for 36 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 4.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (6-iodobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol **3y** as a yellow oil (0.850 g, 71% yield).

Procedure for the 3.0 mmol Scale Reaction for the synthesis of 3ae

Following the procedure, treatment of 4-nitrobenzaldehyde **1a** (0.453 g, 3.0 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (2.238 g, 7.5 mmol) and KBr (0.893 g, 7.5 mmol) in the presence of KF (0.870 g, 15.0 mmol) and 18-crown-6 (3.960 g, 15.0 mmol) in THF (12.0 mL) at 25 °C for 36 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 4.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-bromophenyl)(4-nitrophenyl)methanol **3ae** as a yellow solid (0.691 g, 75% yield).

4. Procedure for the Iodide triggered aryne MCC with *N*-Methylisatin

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KI (0.104 g, 0.625 mmol), KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in a glove-box. Then 1-methylindoline-2,3-dione **4a** (0.040, 0.25 mmol) was added outside the glove-box under nitrogen atmosphere followed by addition of THF (1 mL). To the stirring solution, aryne precursor **2a** (0.186 g, 152 μ L, 0.625 mmol) was added and reaction mixture was allowed to stir at 60 °C for 24 h. After 24 h, the reaction was quenched by adding 4N HCl (HCl in dioxane, 0.4 mL), the solvent was evaporated and the crude residue pre-adsorbed on silica gel and purified by flash column chromatography (Pet. ether /EtOAc = 70/30) afforded 3-hydroxy-3-(2-iodophenyl)-1-methylindolin-2-one **5a** in 49% yield.

5. Procedure for the Iodide triggered aryne MCC with CO₂

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KI (0.104 g, 0.625 mmol), KF (0.029 g, 0.5 mmol) and 18-crown-6 (0.132 g, 0.5 mmol) inside the glove-box. Then the screw-capped test tube was evacuated and backfilled with CO₂ (balloon) for four times followed by the addition of THF (1 mL). Then aryne precursor (0.075 g, 0.25 mmol) was added, the reaction mixture was allowed to stir for 24 h under the atmosphere of CO₂ (balloon). After 24 h the reaction mixture was quenched with cold water and washed with dichloromethane three times. Then 1N HCl was added to the water layer and extracted with ethyl acetate (four times). Then solvent was evaporated to afford 2-iodobenzoic acid in 29% yield.

6. Mechanistic Experiments

Deuterium Labeling Experiment

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KI (0.104 g, 0.625 mmol), KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in a glove-box. Then D_2O (0.005 g, 4.5 μL , 0.25 mmol) was added outside the glove-box under nitrogen atmosphere followed by addition of THF (1 mL). To the stirring solution, aryne precursor **2a** (0.186 g, 152 μL , 0.625 mmol) was added and reaction mixture was allowed to stir at rt for 24 h. After 24 h the solvent was evaporated and the crude residue pre-adsorbed on silica gel and purified by flash column chromatography (Pet. ether) on silica gel to afford the iodobenzene **5** in 88% yield with 86% D incorporation at the 2-position.

^1H NMR spectrum of Iodobenzene-D (86% D)

Deuterium Labeling Experiment in presence of aldehyde:

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KI (0.104 g, 0.625 mmol), KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in a glove-box. Then 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol) and D₂O (0.005 g, 4.5 μ L, 0.25 mmol) was added outside the glove-box under nitrogen atmosphere followed by addition of THF (1 mL). To the stirring solution, aryne precursor **2a** (0.186 g, 152 μ L, 0.625 mmol) was added and reaction mixture was allowed to stir at rt for 24 h. After 24 h the reaction was quenched by adding 4N HCl (HCl in dioxane, 0.4 mL), the solvent was evaporated and the crude residue pre-adsorbed on silica gel and purified by flash column chromatography (Pet. ether /EtOAc = 90/10) on silica gel to afford the product **3a** in (0.036 g) 40% yield along with the iodobenzene with 90% D incorporation at the 2-position in (0.024 g) 49% yield.

¹H NMR spectrum of Iodobenzene-D (90% D)

Deuterium Labeling Experiment with excess D₂O:

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the KI (0.104 g, 0.625 mmol), KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in a

glove-box. Then 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol) and D₂O (0.013 g, 11.5 μ L, 0.625 mmol) was added outside the glove-box under nitrogen atmosphere followed by addition of THF (1 mL). To the stirring solution, aryne precursor **2a** (0.186 g, 152 μ L, 0.625 mmol) was added and reaction mixture was allowed to stir at rt for 24 h. After 24 h the reaction was quenched by adding 4N HCl (HCl in dioxane, 0.4 mL), the solvent was evaporated and the crude residue pre-adsorbed on silica gel and purified by flash column chromatography (Pet. ether /EtOAc = 90/10) on silica gel to afford the product **3a** (0.008 g) 9% yield along with the iodobenzene with 95% D incorporation at the 2-position in (0.052 g) 41% yield.

¹H NMR spectrum of Iodobenzene-D (95% D)

7. X-ray Data of **3ac** and **8a**

Data of **3ac**

Single crystal of **3ac** (recrystallized from CH₂Cl₂ at 25 °C) was mounted and the diffraction data was collected at 293 K on a Bruker SMART APEX CCD diffractometer using SMART/SAINT software. Intensity data were collected using graphite-monochromatized Mo-K α radiation (71.073 pm). The structure was solved by direct methods using the SHELX-97² and refined by full-matrix least-squares on F². Empirical absorption corrections were applied with SADABS.³ All Non-hydrogen atoms were refined anisotropically and hydrogen atoms were included in geometric positions. Structure was drawn using Olex-2 and ORTEP-3. The crystallographic refinement parameters are given below:

² G. M. SHELXL-2013 Sheldrick, University of Göttingen: Göttingen, Germany, 2014.

³ G. M. Sheldrick, SADABS, University of Göttingen, Göttingen, Germany, 1999.

CCDC	CCDC 1905768
Identification code	3ac
Empirical formula	C ₁₄ H ₁₂ INO ₄
Formula weight	385.15
Temperature/K	293(2) K
Crystal system	Triclinic
Space group	P -1
a/Å	11.2336(5)
b/Å	11.3868(6)
c/Å	12.1898(5)
$\alpha/^\circ$	93.014(4)
$\beta/^\circ$	106.905(4)
$\gamma/^\circ$	108.610(4)
Volume/Å ³	1395.58(12)
Z	4
$\rho_{\text{calc}}/\text{cm}^3$	1.833
μ/mm^{-1}	2.307
F(000)	752
Crystal size/mm ³	0.263 × 0.251 × 0.238
Radiation	MoK α (λ = 0.71073)
Theta range for data collection/ $^\circ$	1.769 to 27.484
Index ranges	-14 ≤ h ≤ 14, -14 ≤ k ≤ 14, -15 ≤ l ≤ 15
Reflections collected	20842
Independent reflections	6405 [R _{int} = 0.0482]
Data/restraints/parameters	6405/0/377
Goodness-of-fit on F ²	1.033
Final R indexes [I ≥ 2σ (I)]	R ₁ = 0.0399, wR ₂ = 0.0861
Final R indexes [all data]	R ₁ = 0.0538, wR ₂ = 0.0965
Largest diff. peak/hole / e Å ⁻³	1.057/-1.330

Ortep Diagram of **3ac** (thermal ellipsoids are shown with 50% probability).

CCDC 1905768 (**3ac**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from Data Centre via www.ccdc.cam.ac.uk/data_request/cif

Data of 8a

Single crystal of **8a** (recrystallized from CH₂Cl₂ at 25 °C) was mounted and the diffraction data was collected at 293 K on a Bruker SMART APEX CCD diffractometer using SMART/SAINT software. Intensity data were collected using graphite-monochromatized Mo-K α radiation (71.073 pm). The structure was solved by direct methods using the SHELX-97² and refined by full-matrix least-squares on F². Empirical absorption corrections were applied with SADABS.³ All Non-hydrogen atoms were refined anisotropically and hydrogen atoms were included in geometric positions. Structure was drawn using Olex-2 and ORTEP-3. The crystallographic refinement parameters are given below:

CCDC	CCDC 1911771
Identification code	8a
Empirical formula	C ₂₁ H ₁₅ NO ₃
Formula weight	329.355
Temperature/K	293(2) K
Crystal system	Monoclinic
Space group	C 2/c
a/Å	19.8644(17)
b/Å	13.8976(11)
c/Å	12.4126(8)
α /°	90 (4)
β /°	102.483 (4)
γ /°	90 (4)
Volume/Å ³	3345.7(5)
Z	4
$\rho_{\text{calc}}/\text{cm}^3$	1.308
μ/mm^{-1}	0.088
F(000)	1376
Radiation	MoK α (λ = 0.71073)
Index ranges	-25 $\leq$ h $\leq$ 25, -18 $\leq$ k $\leq$ 17, -16 $\leq$ l $\leq$ 16
Reflections collected	15233
Independent reflections	3840 [R(int) = 0.0844]

Data/restraints/parameters	3840 / 0 / 234
Goodness-of-fit on F^2	1.029
Final R indexes [$I \geq 2\sigma(I)$]	$R_1 = 0.0781$, $wR_2 = 0.1605$
Final R indexes [all data]	$R_1 = 0.1650$, $wR_2 = 0.2154$
Largest diff. peak/hole / $e \text{ \AA}^{-3}$	0.150/ -0.206

Ortep Diagram of **8a** (thermal ellipsoids are shown with 50% probability).

CCDC 1911771 (**8a**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from Data Centre via www.ccdc.cam.ac.uk/data_request/cif

8. Synthesis and Characterization of 2-HaloBenzyl Alcohol Derivatives

(2-Iodophenyl)(4-nitrophenyl)methanol (**3a**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 152 μL , 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 $^{\circ}\text{C}$ for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-iodophenyl)(4-nitrophenyl)methanol **3a** as a yellow solid (0.062 g, 70 % yield).

R_f (Pet. ether /EtOAc = 85/15): 0.40, ^1H NMR (400 MHz, CDCl_3) δ 8.17 (d, $J = 8.7$ Hz, 2H), 7.85 (d, $J = 8.2$ Hz, 1H), 7.60 (d, $J = 8.8$ Hz, 2H), 7.39-7.34 (m, 2H), 7.04-7.00 (m, 1H), 6.18 (s,

1H), 2.75 (s, 1H). ¹³C NMR (100 MHz, CDCl₃) δ 149.44, 147.42, 144.68, 139.92, 130.26, 129.13, 128.66, 127.86, 123.78, 98.72, 78.28. HRMS (ESI) calculated [M-OH]⁺ for C₁₃H₉INO₂: 337.9672, found: 337.9678. FTIR (cm⁻¹) 3400, 3074, 1601, 1718, 1598, 1517, 1008, 862.

(2-Iodophenyl)(phenyl)methanol (**3b**)⁴

Following the general procedure, treatment of benzaldehyde **1b** (0.027 g, 0.026 μL, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 152.0 μL, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-Iodophenyl)(phenyl)methanol **3b** as a yellow oil (0.021 g, 27% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.45; ¹H NMR (400 MHz, CDCl₃) δ 7.84 (dd, *J*₁ = 7.9, *J*₂ = 1.1 Hz, 1H), 7.53 (dd, *J*₁ = 7.8, *J*₂ = 1.6 Hz, 1H), 7.43-7.40 (m, 2H), 7.38-7.27 (m, 4H), 6.99 (td, *J*₁ = 7.7, *J*₂ = 1.7 Hz, 1H), 6.07 (d, *J* = 3.4, 1H), 2.38 (d, *J* = 3.7, 1H). ¹³C NMR (100 MHz, CDCl₃) δ 145.49, 142.24, 139.70, 129.59, 128.73, 128.63, 128.51, 127.93, 127.36, 98.79, 79.15. HRMS (ESI) calculated [M-OH]⁺ for C₁₃H₁₀I: 292.9822, found: 298.9829. FTIR (cm⁻¹) 3319, 3059, 2365, 1586, 1181, 1006.

(4-Bromophenyl)(2-iodophenyl)methanol (**3c**)

Following the general procedure, treatment of 4-bromobenzaldehyde **1c** (0.093 g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (4-bromophenyl)(2-iodophenyl)methanol **3c** as a yellow oil (0.132 g, 67% yield).

⁴ Cvengroš, J., Stolz, D., Togni, A. *Synthesis* **2009**, 16, 2818.

R_f (Pet. ether /EtOAc = 85/15): 0.5; **¹H NMR (400 MHz, CDCl₃)** δ 7.83 (d, *J* = 8.0 Hz, 1H), 7.46-7.44 (m, 3H), 7.36(t, *J* = 7.2 Hz, 1H), 7.27 (d, *J* = 8.3 Hz, 2H), 7.00 (td, *J*₁ = 7.6 Hz, *J*₂ = 1.6 Hz 1H), 6.00 (s, 1H), 2.61 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 145.06, 141.19, 139.75, 131.68, 129.80, 129.02, 128.83, 128.39, 121.85, 98.68, 78.50. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉BrI: 370.8927, found: 370.8933. Isotopic Mass: [M-OH+2]⁺ 372.8914. **FTIR (cm⁻¹)** 3350, 2923, 1639, 1178, 1007, 853.

(4-Chlorophenyl)(2-iodophenyl)methanol (**3d**)

Following the general procedure, treatment of 4-chlorobenzaldehyde **1d** (0.035 g, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 152.0 μL, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (4-chlorophenyl)(2-iodophenyl)methanol **3d** as a yellow oil (0.042 g, 49% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.45; **¹H NMR (400 MHz, CDCl₃)** δ 7.84 (dd, *J*₁ = 7.9 Hz, *J*₂ = 1.1 Hz, 1H), 7.47 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.6 Hz, 1H), 7.39-7.29 (m, 5H), 7.00 (td, *J*₁ = 7.6, *J*₂ = 1.7 Hz, 1H), 6.03 (s, 1H), 2.47 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 145.13, 140.67, 139.77, 133.68, 129.81, 128.85, 128.75, 128.71, 128.39, 98.67, 78.49. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉ClI: 326.9432, found: 326.9439. Isotopic Mass: [M-OH+2]⁺ 328.9413. **FTIR (cm⁻¹)** 3337, 2923, 1641, 1489, 1009, 855.

(4-Fluorophenyl)(2-iodophenyl)methanol (**3e**)

Following the general procedure, treatment of 4-fluorobenzaldehyde **1e** (0.062g, 53.6 μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash

column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (4-fluorophenyl)(2-iodophenyl)methanol **3e** as a white solid (0.074 g, 47% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.58; **¹H NMR (400 MHz, CDCl₃)** δ 7.84 (dd, *J*₁ = 8.0 Hz, *J*₂ = 0.9 Hz, 1H), 7.50 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.5 Hz, 1H), 7.40-7.34 (m, 3H), 7.04-6.98 (m, 3H), 6.01 (s, 1H), 2.60 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 162.35 (d, *J* = 246.2 Hz), 145.28, 139.74, 137.94 (d, *J* = 3.2 Hz), 129.68, 129.15 (d, *J* = 8.2 Hz), 128.77, 128.27, 115.44 (d, *J* = 21.5 Hz), 98.61, 78.48. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉FI: 310.9727, found: 310.9732. **FTIR (cm⁻¹)** 3584, 3322, 3059, 1602, 1507, 1225, 1007, 843.

[1,1'-Biphenyl]-4-yl(2-iodophenyl)methanol (**3f**)

Following the general procedure, treatment of [1,1'-biphenyl]-4-carbaldehyde **1f** (0.091 g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded [1,1'-biphenyl]-4-yl(2-iodophenyl)methanol **3f** as a yellow oil (0.070 g, 36% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.49; **¹H NMR (400 MHz, CDCl₃)** δ 7.87 (d, *J* = 7.9 Hz, 1H), 7.61-7.58 (m, 5H), 7.50-7.34 (m, 6H), 7.04-7.00 (m, 1H), 6.11 (s, 1H), 2.61 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 145.46, 141.25, 140.82, 140.74, 139.70, 129.61, 128.89, 128.76, 128.48, 127.77, 127.46, 127.33, 127.20, 98.78, 78.92. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉INO₂: 369.0135, found: 369.0142. **FTIR (cm⁻¹)** 3357, 3057, 2363, 1486, 1005, 855.

(2-Iodophenyl)(4-(trifluoromethyl)phenyl)methanol (**3g**)

Following the general procedure, treatment of 4-(trifluoromethyl)benzaldehyde **1g** (0.087 g, 68 μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent

purification via silica gel flash column chromatography (Pet. ether /EtOAc = 80/20) of the crude reaction mixture afforded (2-iodophenyl)(4-(trifluoromethyl)phenyl)methanol **3g** as a yellow oil (0.115 g, 61% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.51; **¹H NMR (400 MHz, CDCl₃)** δ 7.86-7.84 (m, 1H), 7.59 (d, *J* = 8.4 Hz, 2H), 7.53 (d, *J* = 8.2 Hz, 2H), 7.43-7.34 (m, 2H), 7.03-6.99 (m, 1H), 6.12 (s, 1H), 2.80 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 146.07, 144.94, 139.80, 129.98, 129.93 (d, *J* = 32.5 Hz), 128.96, 128.56, 127.42, 125.50 (q, *J* = 3.95 Hz), 122.86, 98.78, 78.50. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₄H₉IN: 317.9775, found: 317.9781. **FTIR (cm⁻¹)** 3583, 3303, 2926, 1325, 1167, 1124, 1066, 1011.

4-(Hydroxy(2-iodophenyl)methyl)benzonitrile (**3h**)

Following the general procedure, treatment of 4-formylbenzonitrile **1h** (0.065g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 80/20) of the crude reaction mixture afforded 4-(hydroxy(2-iodophenyl)methyl)benzonitrile **3h** as a yellow oil (0.114 g, 68% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.39; **¹H NMR (400 MHz, CDCl₃)** δ 7.87 (d, *J* = 7.6 Hz, 1H), 7.60 (d, *J* = 8.7 Hz, 2H), 7.54 (d, *J* = 8.3 Hz, 2H), 7.44-7.35 (m, 2H), 7.04-6.99 (m, 1H), 6.13 (s, 1H), 2.65 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 147.35, 144.61, 139.77, 132.29, 130.08, 128.97, 128.53, 127.64, 118.79, 111.41 98.63, 78.30. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₄H₉IN: 317.9774, found: 317.9781. **FTIR (cm⁻¹)** 3584, 3323, 2364, 1569, 1454, 1241, 1008, 886.

Methyl-4-(hydroxy(2-iodophenyl)methyl)benzoate (**3i**)

Following the general procedure, treatment of methyl 4-formylbenzoate **1i** (0.082 g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5mmol) and

18-crown-6 (0.660 g, 2.5mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded methyl-4-(hydroxy(2-iodophenyl)methyl)benzoate **3i** as a yellow oil (0.126 g, 68% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.41; **¹H NMR (400 MHz, CDCl₃)** δ 7.98-7.96 (m, 2H), 8.83 (d, *J* = 7.9 Hz, 1H), 7.47 (d, *J* = 8.1 Hz, 2H), 7.42 (dd, *J_I* = 7.8 Hz, *J₂* = 1.5 Hz, 1H), 7.34 (t, *J* = 7.5 Hz, 1H), 7.00-6.96 (m, 1H), 6.10 (s, 1H), 3.88 (s, 3H), 2.86 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 167.06, 147.36, 145.12, 139.72, 129.85, 129.83, 129.45, 128.85, 128.63, 127.11, 98.80, 78.65, 52.26. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₅H₁₂INO₂: 350.9876, found: 350.9881. **FTIR (cm⁻¹)** 3477, 2363, 1718, 1434, 1282, 1110, 1010, 870.

4-(Hydroxy(2-iodophenyl)methyl)benzaldehyde (**3j**)

Following the general procedure, treatment of terephthalaldehyde **1j** (0.034g, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 151 μL, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 80/20) of the crude reaction mixture afforded 4-(hydroxy(2-iodophenyl)methyl)benzaldehyde **3j** as a colourless oil (0.050 g, 59% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.41; **¹H NMR (400 MHz, CDCl₃)** δ 9.97 (s, 1H), 7.85-7.83 (m, 3H), 7.59 (d, *J* = 8.4 Hz, 2H), 7.42 (dd, *J_I* = 7.7 Hz, *J₂* = 1.6 Hz, 1H), 7.36 (t, *J* = 7.3 Hz, 1H), 7.03-6.99 (m, 1H), 6.15 (s, 1H), 2.76 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 192.16, 149.03, 144.97, 139.82, 136.77, 130.05, 130.02, 128.99, 128.68, 127.69, 98.81, 78.67. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₄H₁₀IO: 320.9771, found: 320.9774. **FTIR (cm⁻¹)** 3747, 3584, 2964, 2364, 1740, 1595, 1425, 1219.

(3-Bromophenyl)(2-iodophenyl)methanol (**3k**)⁴

Following the general procedure, treatment of 3-bromobenzaldehyde **1k** (0.093 g, 0.059 μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethane

sulfonate **2a** (0.373 g, 303 μ L, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (3-bromophenyl)(2-iodophenyl)methanol **3k** as a yellow oil (0.130 g, 66% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.53; **¹H NMR (400 MHz, CDCl₃)** δ 7.84 (dd, J_1 = 7.9 Hz, J_2 = 0.8 Hz, 1H), 7.58 (s, 1H), 7.45-7.35 (m, 3H), 7.31 (d, J = 7.8 Hz, 1H), 7.19 (t, J = 7.8 Hz, 1H), 7.00 (td, J_1 = 7.7, J_2 = 1.7 Hz, 1H), 6.02 (d, J = 2.6 Hz, 1H), 2.63 (d, J = 3.4 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 144.91, 144.49, 139.75, 130.94, 130.21, 130.12, 129.87, 128.89, 128.51, 125.91, 122.74, 98.76, 78.43. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉BrI: 370.8927, found: 370.8933. Isotopic Mass: [M-OH+2]⁺ 372.89143. **FTIR (cm⁻¹)** 3579, 3341, 2916, 1566, 1428, 1007.

(2-Fluorophenyl)(2-iodophenyl)methanol (**3l**)

Following the general procedure, treatment of 2-fluorobenzaldehyde **1l** (0.062g, 52.85 μ L, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μ L, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-fluorophenyl)(2-iodophenyl)methanol **3l** as a yellow oil (0.059 g, 36% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.56; **¹H NMR (400 MHz, CDCl₃)** δ 7.85 (dd, J_1 = 7.9 Hz, J_2 = 1.2 Hz, 1H), 7.49 (dd, J_1 = 7.9 Hz, J_2 = 1.7 Hz, 1H), 7.40-7.36 (m, 1H), 7.33-7.27 (m, 1H), 7.25-7.21 (m, 1H), 7.13-7.00 (m, 3H), 6.29 (d, J = 3.9 Hz, 1H), 2.59 (d, J = 4.3 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 160.74 (d, J = 217.8 Hz), 140.00, 139.71, 129.82, 129.74, 129.18 (d, J = 13.2 Hz), 128.65 (d, J = 3.8 Hz), 128.55, 128.37, 124.33 (d, J = 3.3 Hz), 115.56 (d, J = 22.1 Hz), 98.73, 73.63 (d, J = 3.7 Hz). **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉FI: 310.9727, found: 310.9730. **FTIR (cm⁻¹)** 3584, 3345, 3058, 2363, 1488, 1456, 1227, 1008.

(2-Chlorophenyl)(2-iodophenyl)methanol (**3m**)⁴

Following the general procedure, treatment of 2-chlorobenzaldehyde **1m** (0.070 g, 56.30 μ L, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethane sulfonate **2a** (0.373 g, 303 μ L, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-chlorophenyl)(2-iodophenyl)methanol **3m** as a yellow oil (0.065 g, 38% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.58; **¹H NMR (400 MHz, CDCl₃)** δ 7.87 (dd, J_1 = 7.8 Hz, J_2 = 0.9 Hz, 1H), 7.41-7.25 (m, 6H), 7.04-7.00 (m, 1H), 6.30 (d, J = 3.7 Hz, 1H), 2.58 (d, J = 4.0 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 143.88, 139.79, 139.52, 133.87, 129.77, 129.25, 128.56, 128.48, 127.11, 99.72, 76.49. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉ClI: 326.9432, found: 326.9436. Isotopic Mass: [M-OH+2]⁺ 328.9406. **FTIR (cm⁻¹)** 3583, 3301, 3059, 1626, 1468, 1436, 1055, 1009.

(2-Iodophenyl)(2-nitrophenyl)methanol (**3n**)

Following the general procedure, treatment of 2-nitrobenzaldehyde **1n** (0.076 g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μ L, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-iodophenyl)(2-nitrophenyl)methanol **3n** as a yellow oil (0.098 g, 55% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.35; **¹H NMR (400 MHz, CDCl₃)** δ 7.99 (dd, J_1 = 8.1 Hz, J_2 = 1.2 Hz, 1H), 7.83-7.81 (m, 1H), 7.56 (td, J_1 = 7.7 Hz, J_2 = 1.2 Hz, 1H), 7.51-7.45 (m, 2H), 7.39 (t, J = 7.5 Hz, 1H), 7.32-7.30 (m, 1H), 7.03 (td, J_1 = 7.7, J_2 = 1.7 Hz, 1H), 6.43 (d, J = 3.7 Hz, 1H), 3.31 (d, J = 4.0 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 149.50, 143.18, 139.76, 137.17, 133.74, 129.86, 129.78, 128.91, 128.56, 128.15, 124.83, 99.26, 74.78. **HRMS (ESI)**

calculated $[M-OH]^+$ for $C_{13}H_9INO_2$: 337.9672, found: 337.9681. **FTIR** (cm^{-1}) 3402, 2924, 1643, 1524, 1345, 1012.

(3,4-Dichlorophenyl)(2-iodophenyl)methanol (**3o**)

Following the general procedure, treatment of 3,4-dichlorobenzaldehyde **1o** (0.0875g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μ L, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 88/12) of the crude reaction mixture afforded (3,4-dichlorophenyl)(2-iodophenyl)methanol **3o** as a yellow oil (0.123 g, 65% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.49; **¹H NMR** (400 MHz, $CDCl_3$) δ 7.83 (dd, $J_1 = 7.8$ Hz, $J_2 = 0.7$ Hz, 1H), 7.50 (d, $J = 1.9$ Hz, 1H), 7.40-7.34 (m, 3H), 7.20-7.17 (m, 1H), 7.03-6.98 (m, 1H), 5.98 (d, $J = 3.0$ Hz, 1H), 2.89 (d, $J = 3.5$ Hz, 1H). **¹³C NMR** (100 MHz, $CDCl_3$) δ 144.55, 142.36, 139.79, 132.66, 131.79, 130.47, 130.02, 129.11, 128.96, 128.36, 126.57, 98.66, 77.94. **HRMS (ESI)** calculated $[M-OH]^+$ for $C_{13}H_8Cl_2I$: 360.9042, found: 360.9048. Isotopic Mass: $[M-OH+2]^+$ 362.9019, $[M-OH+4]^+$ 364.8993. **FTIR** (cm^{-1}) 3583, 3298, 3058, 2364, 1466, 1390, 1032, 889.

(2-Chloro-6-fluorophenyl)(2-iodophenyl)methanol (**3p**)

Following the general procedure, treatment of 2-chloro-6-fluorobenzaldehyde **1p** (0.0792g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μ L, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5mmol) and 18-crown-6 (0.660 g, 2.5mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-chloro-6-fluorophenyl)(2-iodophenyl)methanol **3p** as a white solid (0.110 g, 61% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.61; **¹H NMR** (400 MHz, $CDCl_3$) δ 7.87 (dd, $J_1 = 8.0$ Hz, $J_2 = 1.1$ Hz, 1H), 7.52 (d, $J = 7.7$ Hz, 1H), 7.38-7.34 (m, 1H), 7.28-7.22 (m, 2H), 7.03-6.96 (m, 2H),

6.32 (d, $J = 6.4$ Hz, 1H), 2.80(d, $J = 4.2$ Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ 161.785 (d, $J = 251.8$ Hz), 143.13 (d, $J = 1.7$ Hz), 139.88, 136.39 (d, $J = 6.5$ Hz), 130.05 (d, $J = 10.0$ Hz), 129.55, 128.26 (d, $J = 3.9$ Hz), 128.07, 127.00 (d, $J = 13.8$ Hz), 125.91 (d, $J = 3.6$ Hz), 115.35 (d, $J = 23.4$ Hz), 97.98, 74.88. **HRMS (ESI)** calculated $[\text{M-OH}]^+$ for $\text{C}_{13}\text{H}_8\text{ClFI}$: 344.9338, found: 344.9343. Isotopic Mass: $[\text{M-OH}+2]^+$ 346.9315. **FTIR (cm^{-1})** 3747, 3584, 2925, 2364, 1458, 1170, 1014, 902.

(3,5-Bis(trifluoromethyl)phenyl)(2-iodophenyl)methanol (**3q**)

Following the general procedure, treatment of 3,5-bis(trifluoromethyl)benzaldehyde **1q** (0.121 g, 82 μL 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL , 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (3,5-bis(trifluoromethyl)phenyl)(2-iodophenyl)methanol **3q** as a yellow oil (0.133 g, 60% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.56; ^1H NMR (400 MHz, CDCl_3) δ 7.91(s, 2H), 7.88 (d, $J = 8.1$ Hz, 1H), 7.81 (s, 1H), 7.41-7.35 (m, 2H), 7.07-7.02 (m, 1H), 6.20 (s, 1H), 2.80 (bs, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ 143.83, 143.35, 139.06, 130.82 (q, $J = 33$ Hz), 129.46, 128.29, 127.55, 126.23, 122.41 (q, $J = 272$ Hz), 120.79 (p, $J = 3.8$ Hz), 97.75, 77.06. **HRMS (ESI)** calculated $[\text{M-OH}]^+$ for $\text{C}_{15}\text{H}_8\text{IF}_6\text{I}$: 428.9569, found: 428.9573. **FTIR (cm^{-1})** 3319, 1374, 1280, 1173, 1134, 900.

(2-Iodophenyl)(naphthalen-2-yl)methanol (**3r**)

Following the general procedure, treatment of 2-naphthaldehyde **1r** (0.078 g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL , 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column

chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-iodophenyl)(naphthalen-2-yl)methanol **3r** as a yellow oil (0.056 g, 31% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.43; **¹H NMR (400 MHz, CDCl₃)** δ 7.91 (s, 1H), 7.87-7.80 (m, 4H), 7.53 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.5 Hz, 1H), 7.51-7.48 (m, 2H), 7.47-7.46 (m, 1H), 7.39-7.35 (m, 1H), 7.01 (td, *J*₁ = 7.7 Hz, *J*₂ = 1.7 Hz, 1H), 6.23 (s, 1H), 2.56 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 145.38, 139.74, 139.62, 133.34, 133.04, 129.69, 128.76, 128.73, 128.41, 128.31, 127.80, 126.32, 126.23, 126.08, 125.29, 99.04, 79.26. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₇H₁₂I: 342.9978, found: 342.9984. **FTIR (cm⁻¹)** 3340, 3055, 1562, 1432, 1007, 859.

(2-Iodophenyl)(pyren-1-yl)methanol (**3s**)

Following the general procedure, treatment of pyrene-1-carbaldehyde **1s** (0.115 g, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-iodophenyl)(pyren-1-yl)methanol **3s** as a brown oil (0.06 g, 28% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.43; **¹H NMR (400 MHz, CDCl₃)** δ 8.19-7.92 (m, 9H), 7.88 (d, *J* = 8.9 Hz, 1H), 7.31-7.23 (m, 2H), 6.98 (td, *J*₁ = 7.6 Hz, *J*₂ = 1.8 Hz, 1H), 6.93 (s, 1H), 2.73 (d, *J* = 2.8 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 145.29, 139.84, 135.05, 131.39, 131.22, 130.74, 129.73, 129.14, 128.66, 128.23, 127.67, 127.57, 126.14, 125.54, 125.44, 125.15, 125.05, 124.92, 124.82, 124.51, 123.10, 99.68, 76.72. **HRMS (ESI)** calculated [M-OH]⁺ for C₂₃H₁₄I: 417.0135, found: 417.0144. **FTIR (cm⁻¹)** 3742, 3585, 3276, 3043, 2363, 1428, 1008, 842.

(2-Iodophenyl)(thiophen-2-yl)methanol (**3t**)

Following the general procedure, treatment of thiophene-2-carbaldehyde **1t** (0.056g, 47μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8

mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 80/20) of the crude reaction mixture afforded (2-iodophenyl)(thiophen-2-yl)methanol **3t** as a yellow oil (0.038 g, 24% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.37; **¹H NMR (400 MHz, CDCl₃)** δ 7.83 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.0 Hz, 1H), 7.68 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.6 Hz, 1H), 7.44-7.40 (m, 1H), 7.29 (dd, *J*₁ = 4.3 Hz, *J*₂ = 2.2 Hz, 1H), 7.04-7.00 (m, 1H), 6.96-6.94 (m, 2H), 6.26 (d, *J* = 3.6 Hz, 1H), 2.55 (d, *J* = 3.9 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 146.37, 145.04, 139.64, 129.88, 128.84, 127.79, 126.89, 125.97, 125.90, 98.05, 75.75. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₁H₈IS: 298.9386, found: 298.9388. **FTIR (cm⁻¹)** 3744, 3585, 2923, 2363, 1461, 1429, 1005, 853.

Benzofuran-2-yl(2-iodophenyl)methanol (**3u**)

Following the general procedure, treatment of benzofuran-2-carbaldehyde **1u** (0.073 g, 61 μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5mmol) and 18-crown-6 (0.660 g, 2.5mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded benzofuran-2-yl(2-iodophenyl)methanol **3u** as a brown oil (0.090 g, 51% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.45; **¹H NMR (400 MHz, CDCl₃)** δ 7.86 (dd, *J*₁ = 7.9 Hz, *J*₂ = 0.8 Hz, 1H), 7.66 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.5 Hz, 1H), 7.52-7.47 (m, 2H), 7.44-7.40 (m, 1H), 7.31-7.26 (m, 1H), 7.24-7.20 (m, 1H), 7.06 (td, *J*₁ = 7.7 Hz, *J*₂ = 1.6 Hz, 1H), 6.46 (s, 1H), 6.17 (s, 1H), 2.91 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 157.09, 155.23, 142.25, 139.60, 130.14, 128.76, 128.41, 128.06, 124.58, 122.98, 121.38, 111.51, 105.31, 98.45, 74.18. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₅H₁₀IO: 332.9771, found: 332.9777. **FTIR (cm⁻¹)** 3322, 3059, 2364, 1585, 1454, 1253, 1007.

Cyclohexyl(2-iodophenyl)methanol (**3v**)

Following the general procedure, treatment of cyclohexanecarbaldehyde **1v** (0.056 g, 0.061 μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g,

1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded cyclohexyl(2-iodophenyl)methanol **3v** as a white solid (0.066 g, 42% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.58; **¹H NMR (400 MHz, CDCl₃)** δ 7.80 (dd, *J*₁ = 7.9 Hz, *J*₂ = 1.2 Hz, 1H), 7.44-7.41 (m, 1H), 7.37-7.33 (m, 1H), 6.98-6.94 (m, 1H), 4.71-4.69 (m, 1H), 1.93 (d, *J* = 3.5 Hz, 1H), 1.85-1.65 (m, 5H), 1.48-1.45 (m, 1H), 1.30-1.16 (m, 5H). **¹³C NMR (100 MHz, CDCl₃)** δ 145.66, 139.43, 129.18, 128.38, 128.13, 98.95, 81.59, 44.16, 29.86, 27.38, 26.51, 26.16. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₁₆I: 299.0291, found: 299.0294. **FTIR (cm⁻¹)** 3408, 3056, 2926, 2850, 1613, 1456, 1007.

1-(2-Iodophenyl)-2-methylpropan-1-ol (**3w**)

Following the general procedure, treatment of isobutyraldehyde **1w** (0.036 g, 0.046 μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KI (0.208 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded 1-(2-iodophenyl)-2-methylpropan-1-ol **3w** as a colourless oil (0.039 g, 28% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.82; **¹H NMR (400 MHz, CDCl₃)** δ 7.81 (d, *J* = 7.9 Hz, 1H), 7.45-7.43 (m, 1H), 7.38-7.34 (m, 1H), 6.98-6.94 (m, 1H), 4.70 (d, *J* = 5.6 Hz, 1H), 2.11-1.99 (m, 1H), 1.94 (bs, 1H), 0.98 (d, *J* = 3.3 Hz, 3H), 0.98 (d, *J* = 3.1 Hz, 3H). **¹³C NMR (100 MHz, CDCl₃)** δ 145.89, 139.49, 129.18, 128.39, 127.99, 98.74, 81.97, 34.18, 19.76, 16.65. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₀H₁₁I: 258.9978, found: 258.9979. **FTIR (cm⁻¹)** 3420, 2962, 1584, 1465, 1209, 1003.

(2-Iodo-4,5-dimethylphenyl)(4-nitrophenyl)methanol (**3x**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 4,5-dimethyl-2-

(trimethylsilyl)phenyl trifluoromethanesulfonate **2b** (0.204 g, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-iodo-4,5-dimethylphenyl)(4-nitrophenyl)methanol **3x** as a yellow oil (0.062 g, 65% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.45; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.19-8.16 (m, 2H), 7.6-7.59 (m, 3H), 7.07 (s, 1H), 6.13 (s, 1H), 2.63 (bs, 1H), 2.20 (s, 3H), 2.17 (s, 3H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 149.93, 147.27, 142.05, 140.32, 139.46, 138.03, 129.56, 127.59, 123.71, 95.02, 78.01, 19.66, 19.13. **HRMS (ESI)** calculated $[\text{M-OH}]^+$ for $\text{C}_{15}\text{H}_{13}\text{INO}_2$: 365.9985, found: 365.9992. **FTIR** (cm^{-1}) 3306, 2920, 2363, 1519, 1345, 859.

(6-Iodobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol (**3y**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 6-(trimethylsilyl)benzo[d][1,3]dioxol-5-yl trifluoromethanesulfonate **2c** (0.214 g, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 21.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (6-iodobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol **3y** as a yellow oil (0.074 g, 74% yield).

(Pet. ether /EtOAc = 85/15): 0.35; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.18 (d, J = 8.6 Hz, 2H), 7.59 (d, J = 8.6 Hz, 2H), 7.23 (s, 1H), 6.82 (s, 1H), 6.13 (s, 1H), 5.98 (d, J = 1.1 Hz, 1H), 5.95 (d, J = 1.1 Hz, 1H), 2.68 (bs, 1H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 149.67, 149.23, 148.69, 147.38, 138.37, 127.49, 123.79, 118.67, 108.49, 102.12, 86.65, 78.17. **HRMS (ESI)** calculated $[\text{M-OH}]^+$ for $\text{C}_{14}\text{H}_9\text{INO}_4$: 381.9571, found: 381.9578. **FTIR** (cm^{-1}) 3585, 2900, 2363, 1473, 1345, 1233, 1036, 859.

(2-Iodo-3,6-dimethylphenyl)(4-nitrophenyl)methanol (**3z**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 3,6-dimethyl-2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2d** (0.204 g, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 85/15) of the crude reaction mixture afforded (2-iodo-3,6-dimethylphenyl)(4-nitrophenyl)methanol **3z** as a yellow oil (0.052 g, 54% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.45; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.17-8.14 (m, 2H), 7.50-7.43 (m, 2H), 7.14 (d, $J = 7.7$ Hz, 1H), 7.02 (d, $J = 7$ Hz, 1H), 6.76 (d, $J = 4.3$ Hz, 1H), 2.86 (bs, 1H), 2.51 (s, 3H), 2.11 (s, 3H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 149.99, 146.90, 142.25, 140.56, 136.58, 132.31, 129.93, 126.46, 123.56, 76.84, 30.67, 20.38. **HRMS (ESI)** calculated $[\text{M}-\text{OH}]^+$ for $\text{C}_{15}\text{H}_{13}\text{INO}_2$: 365.9985, found: 365.9985. **FTIR** (cm^{-1}) 3583, 2922, 2015, 1598, 1516, 1342, 877.

(3-Iodonaphthalen-2-yl)(4-nitrophenyl)methanol (**3aa**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.0378g, 0.25 mmol), 3-(trimethylsilyl)naphthalen-2-yl trifluoromethanesulfonate **2e** (0.218 g, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.072 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 85/15) of the crude reaction mixture afforded (3-iodonaphthalen-2-yl)(4-nitrophenyl)methanol **3aa** as a yellow solid (0.055 g, 55% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.46; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.41(s, 1H), 8.18 (d, $J = 8.8$ Hz, 2H), 7.85 (s, 1H), 7.78-7.76 (m, 1H), 7.73-7.71 (m, 1H), 7.61 (d, $J = 8.5$ Hz, 2H), 7.53-7.49 (m, 2H), 6.29 (s, 1H), 2.81 (bs, 1H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 149.30, 147.45, 140.66, 139.52, 134.66, 132.82, 128.40, 128.26, 127.75, 127.46, 127.29, 126.62, 123.80, 96.30, 78.14.

HRMS (ESI) calculated $[M-OH]^+$ for $C_{17}H_{11}INO_2$: 387.9834, found: 387.9829. **FTIR** (cm^{-1}) 3741, 3584, 2921, 2363, 1517, 1344, 1041, 858.

(2-Iodonaphthalen-1-yl)(4-nitrophenyl)methanol (**3ab**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.0378 g, 0.25 mmol), 1-(trimethylsilyl)naphthalen-2-yl trifluoromethanesulfonate **2f** (0.218 g, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.072 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 85/15) of the crude reaction mixture afforded (2-iodonaphthalen-1-yl)(4-nitrophenyl)methanol **3ab** as a yellow solid (0.055 g, 54% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.43; **¹H NMR (400 MHz, CDCl₃)** δ 8.14 (d, J = 8.8 Hz, 2H), 8.01-7.94 (m, 2H), 7.79 (d, J = 8.1 Hz, 1H), 7.55 (t, J = 9.3 Hz, 3H), 7.43 (t, J = 7.6 Hz, 1H), 7.31-7.27 (m, 1H), 6.85 (d, J = 3.2 Hz, 1H), 3.01 (d, J = 3.8 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 150.00, 147.05, 139.60, 136.22, 134.69, 131.36, 131.11, 128.88, 126.80, 126.66, 126.30, 125.93, 123.79, 100.53, 81.11. **HRMS (ESI)** calculated $[M-OH]^+$ for $C_{17}H_{11}INO_2$: 387.9829, found: 387.9835. **FTIR** (cm^{-1}) 3745, 3584, 2363, 1601, 1515, 1342, 1061, 852.

(2-Iodo-6-methoxyphenyl)(4-nitrophenyl)methanol (**3ac**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 3-methoxy-2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2g** (0.205 g, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 85/15) of the crude reaction mixture afforded (2-iodo-6-methoxyphenyl)(4-nitrophenyl)methanol **3ac** as a yellow solid (0.077 g, 79% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.12; **¹H NMR (400 MHz, CDCl₃)** δ 8.15-8.12 (m, 2H), 7.56 (dd, J_1 = 7.9 Hz, J_2 = 0.8 Hz, 1H), 7.49 (d, J = 8.2 Hz, 2H), 7.02 (t, J = 8.1 Hz, 1H), 7.90 (d, J = 8.2

Hz, 1H), 6.23 (d, J = 11.3 Hz, 1H), 4.12 (d, J = 11.4 Hz, 1H), 3.68 (s, 3H). **^{13}C NMR (100 MHz, CDCl_3)** δ 157.28, 151.18, 147.05, 132.95, 132.79, 130.91, 126.28, 123.44, 112.31, 100.78, 79.69, 55.91. **HRMS (ESI)** calculated $[\text{M}-\text{OH}]^+$ for $\text{C}_{14}\text{H}_{11}\text{INO}_3$: 367.9778, found: 367.9786. **FTIR (cm^{-1})** 3434, 2722, 1567, 1517, 1344, 1024.

(2-Iodo-4-methylphenyl)(4-nitrophenyl)methanol (3ad) and (2-Iodo-5-methylphenyl)(4-nitrophenyl)methanol (3ad')

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 4-methyl-2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2h** (0.195 g, 0.625

mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded inseparable (1:1) regioisomeric mixture of (2-iodo-4-methylphenyl)(4-nitrophenyl)methanol (**3ad**) and (2-iodo-5-methylphenyl)(4-nitrophenyl)methanol (**3ad'**) as a yellow oil (0.046 g, 49% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.35; **^1H NMR (400 MHz, CDCl_3)** δ 7.73-7.10 (m, 1H), 7.62-7.58 (m, 4H), 7.17-7.15 (m, 1H), 6.85 (dd, J_1 = 8.1 Hz, J_2 = 1.8 Hz, 1H), 6.16 (s, 1H), 2.64-2.61 (m, 1H), 2.28 (s, 3H). **^{13}C NMR (100 MHz, CDCl_3)** δ 20.64, 78.08, 94.68, 123.75, 127.73, 128.71, 129.99, 139.35, 140.28, 141.80, 147.37, 149.60. Representative peak for other isomer: **^1H NMR (400 MHz, CDCl_3)** δ 8.20-8.17 (m, 4H), 7.73-7.10 (m, 1H), 7.22-7.20 (m, 1H), 7.17-7.15 (m, 1H), 6.16 (s, 1H), 2.64-2.61 (m, 1H), 2.30 (s, 3H). **^{13}C NMR (100 MHz, CDCl_3)** δ 21.18, 78.18, 98.72, 123.78, 127.75, 129.40, 131.34, 139.62, 140.53, 144.39, 147.40, 149.75. **HRMS (ESI)** calculated $[\text{M}-\text{OH}]^+$ for $\text{C}_{14}\text{H}_{11}\text{INO}_2$: 351.9829, found: 351.9834. **FTIR (cm^{-1})** 3399, 2853, 1600, 1518, 1345, 1011.

(2-Bromophenyl)(4-nitrophenyl)methanol (3ae)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate

2a (0.186 g, 152 μ L, 0.625 mmol) and KBr (0.074 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-bromophenyl)(4-nitrophenyl)methanol **3ae** as a yellow solid (0.057 g, 74% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.39; **¹H NMR (400 MHz, CDCl₃)** δ 8.17(d, *J* = 8.8 Hz, 2H), 7.60-7.55 (m, 3H), 7.46 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.7 Hz, 1H), 7.34 (td, *J*₁ = 7.6 Hz, *J*₂ = 1.1 Hz, 1H), 7.19 (td, *J*₁ = 7.6 Hz, *J*₂ = 1.7 Hz, 1H), 6.30 (s, 1H), 2.77 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 149.41, 147.36, 141.65, 133.20, 129.95, 128.72, 128.26, 127.72, 123.78, 122.79, 73.93. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉BrNO₂: 289.9811, found: 289.9814. Isotopic Mass: [M-OH+2]⁺ 291.9799. **FTIR (cm⁻¹)** 3584, 3072, 2364, 1518, 1345, 1016, 864.

(2-Chlorophenyl)(4-nitrophenyl)methanol (**3af**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 152 μ L, 0.625 mmol) and KCl (0.047 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-chlorophenyl)(4-nitrophenyl)methanol **3af** as a yellow solid (0.042 g, 62% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.39 **¹H NMR (400 MHz, CDCl₃)** δ 8.18 (d, *J* = 8.8 Hz, 2H), 7.59 (d, *J* = 8.8 Hz, 2H), 7.49 (dd, *J*₁ = 7.6 Hz, *J*₂ = 1.6 Hz, 1H), 7.38 (dd, *J*₁ = 7.7 Hz, *J*₂ = 1.4 Hz, 1H), 7.33-7.24 (m, 2H), 6.34 (d, *J* = 2.7 Hz, 1H), 2.63 (d, *J* = 3.6 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 149.40, 147.28, 140.00, 132.43, 129.83, 129.52, 128.19, 127.55, 127.53, 123.69, 71.71. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₉ClNO₂: 246.0316, found: 246.0322. Isotopic Mass: [M-OH+2]⁺ 248.0296. **FTIR (cm⁻¹)** 3584, 3072, 2363, 1519, 1345, 1025, 865.

(2-Bromophenyl)(phenyl)methanol (**3ag**)⁵

Following the general procedure, treatment of benzaldehyde **1b** (0.053 g, 0.051 μ L, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μ L, 1.25 mmol) and KBr (0.149 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-bromophenyl)(phenyl)methanol **3ag** as a yellow oil (0.100 g, 76% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.56; **¹H NMR (400 MHz, CDCl₃)** δ 7.60-7.54 (m, 2H), 7.42-7.40 (m, 2H), 7.37-7.27 (m, 4H), 7.16 (td, $J_1 = 7.7$ Hz, $J_2 = 1.7$ Hz, 1H), 6.19 (d, $J = 2.8$ Hz, 1H), 2.57 (d, $J = 2.8$ Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 142.59, 142.24, 132.94, 129.21, 128.58, 128.57, 127.88, 127.83, 127.16, 122.89, 74.85. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₁₀Br: 244.9960, found: 244.9967. Isotopic Mass: [M-OH+2]⁺ 246.9947. **FTIR (cm⁻¹)** 3355, 3061, 2363, 1437, 1181, 1013.

(2-Chlorophenyl)(phenyl)methanol (**3ah**)⁵

Following the general procedure, treatment of benzaldehyde **1b** (0.027 g, 0.026 μ L, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 152.0 μ L, 0.625 mmol) and KCl (0.047 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (2-chlorophenyl)(phenyl)methanol **3ah** as a white solid (0.030 g, 54% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.45; **¹H NMR (400 MHz, CDCl₃)** δ 7.62-7.60 (m, 2H), 7.41-7.39 (m, 2H), 7.36-7.27 (m, 4H), 7.26-7.21 (m, 1H), 6.23 (d, $J = 3.7$ Hz, 1H), 2.40 (d, $J = 3.8$ Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 142.35, 141.11, 132.65, 129.69, 128.91, 128.63, 128.17, 127.93, 127.24, 127.06, 72.83. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₃H₁₀Cl: 201.0466,

⁵ Kuriyama, M.; Shimazawa, R.; Enomoto, T.; Shirai, R. *J. Org. Chem.* **2008**, 73, 6939.

found: 201.0473. Isotopic Mass: $[M-OH+2]^+$ 203.0444. **FTIR** (cm^{-1}) 3324, 3029, 2364, 1598, 1443, 1019.

Benzofuran-2-yl(2-bromophenyl)methanol (**3ai**)

Following the general procedure, treatment of benzofuran-2-carbaldehyde **1u** (0.037 g, 30 μL , 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 152.0 μL , 0.625 mmol) and KBr (0.074 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded benzofuran-2-yl(2-bromophenyl)methanol **3ai** as a yellow oil (0.055 g, 71% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.48; **¹H NMR** (400 MHz, CDCl_3) δ 7.70 (dd, $J_1 = 7.8$ Hz, $J_2 = 1.6$ Hz, 1H), 7.58 (dd, $J_1 = 8.0$ Hz, $J_2 = 1.2$ Hz, 1H), 7.52-7.46 (m, 2H), 7.41-7.37 (m, 1H), 7.30-7.26 (m, 1H), 7.25-7.19 (m, 2H), 6.47 (s, 1H), 6.33 (d, $J = 4.3$ Hz, 1H), 2.82 (d, $J = 4.4$ Hz, 1H). **¹³C NMR** (100 MHz, CDCl_3) δ 157.05, 155.22, 139.37, 132.96, 129.91, 128.73, 128.08, 127.95, 124.58, 122.99, 122.91, 121.37, 111.51, 104.97, 69.67. **HRMS (ESI)** calculated $[M-OH]^+$ for $\text{C}_{15}\text{H}_{10}\text{BrO}$: 284.9910, found: 284.9918. Isotopic Mass: $[M-OH+2]^+$ 286.9898. **FTIR** (cm^{-1}) 3323, 3062, 2363, 1568, 1451, 1253, 1016.

Benzofuran-2-yl(2-chlorophenyl)methanol (**3aj**)⁶

Following the general procedure, treatment of benzofuran-2-carbaldehyde **1u** (0.037 g, 30 μL , 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.186 g, 152.0 μL , 0.625 mmol) and KCl (0.047 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded benzofuran-2-yl(2-chlorophenyl)methanol **3aj** as a yellow oil (0.031 g, 48% yield).

⁶ Xuan, Q.; Kong, W.; Song, Q. *J. Org. Chem.* **2017**, *82*, 7602.

R_f (Pet. ether /EtOAc = 95/15): 0.40; **¹H NMR (400 MHz, CDCl₃)** δ 7.71-7.69 (m, 1H), 7.52-7.50 (m, 1H), 7.48-7.45 (m, 1H), 7.41-7.37 (m, 1H), 7.36-7.26 (m, 3H), 7.23-7.19 (m, 1H), 6.48 (s, 1H), 6.37 (d, *J* = 4.4 Hz, 1H), 2.74 (d, *J* = 4.4 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 157.08, 155.22, 137.80, 132.85, 129.71, 129.62, 128.46, 128.09, 127.35, 124.58, 123.01, 121.37, 111.51, 104.76, 67.41. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₅H₁₀ClO: 241.0415, found: 241.0420. Isotopic Mass: [M-OH+2]⁺ 243.0392. **FTIR (cm⁻¹)** 3322, 3064, 2363, 1581, 1450, 1251, 1024.

1-(2-Bromophenyl)-2-methylpropan-1-ol (**3ak**)⁷

Following the general procedure, treatment of isobutyraldehyde **1w** (0.036 g, 0.046 μL, 0.5 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.373 g, 303 μL, 1.25 mmol) and KBr (0.149 g, 1.25 mmol) in the presence of KF (0.145 g, 2.5 mmol) and 18-crown-6 (0.660 g, 2.5 mmol) in THF (2.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.8 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded 1-(2-bromophenyl)-2-methylpropan-1-ol **3ak** as a yellow oil (0.063 g, 55% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.81; **¹H NMR (400 MHz, CDCl₃)** δ 7.53-7.48 (m, 2H), 7.34-7.31 (m, 1H), 7.12 (td, *J*₁ = 7.7 Hz, *J*₂ = 1.6 Hz, 1H), 4.87 (d, *J*₁ = 5.4 Hz, *J*₂ = 3.2 Hz, 1H), 2.13-2.01 (m, 1H), 1.90 (d, *J* = 3.5 Hz, 1H), 0.97 (d, *J* = 6.9 Hz, 1H), 0.95 (d, *J* = 6.9 Hz, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 143.03, 132.79, 128.79, 128.39, 127.55, 122.80, 77.71, 34.11, 19.61, 16.92. **FTIR (cm⁻¹)** 3420, 2963, 2871, 1466, 1199, 1007.

(6-Bromobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol (**3al**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 6-(trimethylsilyl)benzo[d][1,3]dioxol-5-yl trifluoromethanesulfonate **2c** (0.214 g, 0.625 mmol) and KBr (0.074 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 21.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica

⁷ Faggyas, R. J.; E. Calder, D. D.; Wilson, C.; Sutherland, A. *J. Org. Chem.* **2017**, *82*, 11585.

gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (6-bromobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol **3al** as a yellow oil (0.070 g, 79% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.34 **¹H NMR (400 MHz, CDCl₃)** δ 8.19-8.16 (m, 2H), 7.58 (d, *J* = 8.9 Hz, 2H), 6.98 (s, 1H), 6.87 (s, 1H), 6.25 (s, 1H), 5.98 (d, *J* = 1.3 Hz, 1H), 5.95 (d, *J* = 1.3 Hz, 1H), 2.72 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 149.63, 148.44, 148.16, 147.34, 135.12, 127.39, 123.79, 113.27, 112.76, 108.28, 102.18, 73.67. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₄H₉BrNO₄: 333.9709, found: 333.9713. Isotopic Mass: [M-OH+2]⁺ 335.9693. **FTIR (cm⁻¹)** 3584, 2916, 2363, 1474, 1346, 1235, 1036, 861.

(6-Chlorobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol (**3am**)

Following the general procedure, treatment of 4-nitrobenzaldehyde **1a** (0.038 g, 0.25 mmol), 6-(trimethylsilyl)benzo[d][1,3]dioxol-5-yl trifluoromethanesulfonate **2c** (0.214 g, 0.625 mmol) and KCl (0.047 g, 0.625 mmol) in the presence of KF (0.073 g, 1.25 mmol) and 18-crown-6 (0.330 g, 21.25 mmol) in THF (1.0 mL) at 25 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (6-chlorobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol **3am** as a yellow solid (0.043g, 56% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.34 **¹H NMR (400 MHz, CDCl₃)** δ 8.20-8.17 (m, 2H), 7.58 (d, *J* = 8.9 Hz, 2H), 6.89 (s, 1H), 6.83 (s, 1H), 6.28 (s, 1H), 5.99 (d, *J* = 1.3 Hz, 1H), 5.96 (d, *J* = 1.3 Hz, 1H), 2.55 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 149.68, 148.19, 147.52, 147.37, 133.52, 127.33, 124.40, 123.82, 109.94, 107.72, 102.18, 71.46. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₄H₉ClNO₄: 290.0215, found: 290.0219. Isotopic Mass: [M-OH+2]⁺ 292.0193. **FTIR (cm⁻¹)** 3584, 2918, 2364, 1475, 1346, 1240, 1036, 863.

3-Hydroxy-3-(2-iodophenyl)-1-methylindolin-2-one (**5a**)

Following the general procedure, treatment of 1-methylindoline-2,3-dione **4a** (0.040g, 0.25 mmol), 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.187 g, 152 μL, 0.625 mmol) and KI (0.104 g, 0.625 mmol) in the presence

of KF (0.072 g, 1.25 mmol) and 18-crown-6 (0.330 g, 1.25 mmol) in THF (1.0 mL) at 60 °C for 24 h followed by quenching the reaction mixture using 4N HCl (HCl in dioxane, 0.4 mL), and subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded 3-hydroxy-3-(2-iodophenyl)-1-methylindolin-2-one **5a** as a yellow solid (0.045 g, 49% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.23; **¹H NMR (400 MHz, CDCl₃)** δ 8.12 (d, *J* = 7.9 Hz, 1H), 7.80 (d, *J* = 7.5 Hz, 1H), 7.50-7.46 (m, 1H), 7.40-7.36 (m, 1H), 7.05-6.95 (m, 3H), 6.90 (d, *J* = 7.42 Hz, 1H), 3.39 (bs, 1H), 3.30 (s, 3H). **¹³C NMR (100 MHz, CDCl₃)** δ 175.76, 145.84, 142.01, 141.31, 130.59, 130.11, 129.53, 128.72, 128.24, 124.85, 123.48, 108.89, 93.91, 79.57, 26.90. **HRMS (ESI)** calculated [M-OH]⁺ for C₁₅H₁₁INO: 347.9880, found: 347.9884. **FTIR (cm⁻¹)** 3301, 2925, 1706, 1649, 1500, 1251, 1169, 928.

2-Iodobenzoic acid (**6a**)⁸

Following the procedure, treatment of 2-(trimethylsilyl)phenyl trifluoromethanesulfonate **2a** (0.075 g, 61 μL, 0.25 mmol) and KI (0.104 g, 0.625 mmol) in the presence of KF (0.029 g, 0.5 mmol) and 18-crown-6 (0.132 g, 0.5 mmol) in THF (1.0 mL) under CO₂ atmosphere at 25 °C for 24 h followed by quenching the reaction mixture using cold water, and washing with dichloromethane followed by extraction with ethyl acetate afforded 2-iodobenzoic acid **6a** as white solid (0.018 g, 29% yield).

R_f (Pet. ether /EtOAc = 50/50): 0.31; **¹H NMR (400 MHz, dMSO-*d*₆)** δ 13.31 (bs, 1H), 7.98 (dd, *J*₁ = 7.8 Hz, *J*₂ = 1.0 Hz, 1H), 7.71 (dd, *J*₁ = 7.7 Hz, *J*₂ = 1.6 Hz, 1H), 7.50-7.46 (m, 1H), 7.26-7.21 (m, 1H). **¹³C NMR (100 MHz, dMSO-*d*₆)** δ 168.15, 140.50, 136.92, 132.45, 130.05, 128.17, 94.11. **FTIR (cm⁻¹)** 3752, 3585, 3424, 2697, 2365, 1650, 1219, 1021.

9. Functionalization of 2-IodoBenzyl Alcohols

Using Sonogashira Coupling

⁸ Mukhopadhyay, S.; Batra, S. *Chem. Eur. J.*, **2018**, *24*, 14622.

(4-nitrophenyl)(2-(phenylethynyl)phenyl)methanol (**7a**)

Following the literature procedure,⁹ to a mixture of (2-iodophenyl)(4-nitrophenyl)methanol **3a** (107 mg, 0.3 mmol) in DMF (2 mL) were added Pd(PPh₃)₂Cl₂ (4.2 mg, 0.02 equiv) and CuI (2.3 mg, 0.04 equiv) under nitrogen. After the reaction mixture was stirred for 5 min. at rt, *N,N*-diisopropylamine (121 mg, 168 μ L, 4 equiv) was added by a syringe. The reaction mixture was then heated at 70 °C and phenylacetylene (37 mg, 40 μ L, 1.2 equiv) was added. After stirring the reaction mixture for 2 h at 70 °C, the reaction mixture was washed with saturated aq. NH₄Cl and water, and then extracted with EtOAc, dried over anhydrous Na₂SO₄ and concentrated under vacuum. Subsequent purification via silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded (4-nitrophenyl)(2-(phenylethynyl)phenyl)methanol **7a** as a yellow solid (90 mg, 91% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.40; **¹H NMR (400 MHz, CDCl₃)** δ 8.17 (d, *J* = 8.8 Hz, 2H), 7.66 (d, *J* = 8.8 Hz, 2H), 7.57-7.55 (m, 1H), 7.49-7.46 (m, 3H), 7.39-7.35 (m, 4H), 7.33-7.29 (m, 1H), 6.48 (s, 1H), 2.79 (bs, 1H). **¹³C NMR (100 MHz, CDCl₃)** δ 150.38, 147.27, 144.42, 132.83, 131.57, 129.33, 129.00, 128.71, 128.26, 127.32, 126.65, 123.73, 122.64, 121.44, 95.15, 87.08, 73.27. **HRMS (ESI)** calculated [M-OH]⁺ for C₂₁H₁₄NO₂: 312.1019, found: 312.1028. **FTIR (cm⁻¹)** 3582, 1518, 1344, 1179, 1034, 825.

Procedure for the Au-catalyzed cyclization of **7a**

To a flame-dried screw-capped test tube equipped with a magnetic stir bar was added the (PPh₃)AuCl (2.5 mg, 0.005 mmol) and AgOTf (1.3 mg, 0.005 mmol) in a glove-box. Then (4-nitrophenyl)(2-(phenylethynyl)phenyl)methanol **7a** (33 mg, 0.1 mmol) was added outside the glove box under nitrogen atmosphere followed by addition of THF (1 mL). The resulting

⁹ Li, D. Y.; Shi, K. J.; Mao, X. F.; Zhao, Z. L.; Wu, X. Y.; Liu, P. N. *Tetrahedron* **2014**, *70*, 7022.

solution was stirred at rt for 12 h. After 12 h, the solvent was evaporated and the crude residue pre-adsorbed on silica gel and purified by flash column chromatography (Pet. ether /EtOAc = 95/5) afforded (Z)-1-benzylidene-3-(4-nitrophenyl)-1,3-dihydroisobenzofuran **8a** as major isomer (27 mg, 82% yield) along with (Z)-1-benzylidene-3-(4-nitrophenyl)-1,3-dihydroisobenzofuran **8b** (5 mg, 15% yield).

(Z)-1-benzylidene-3-(4-nitrophenyl)-1,3-dihydroisobenzofuran 8a

Yellow solid (27 mg, 82% yield), R_f (Pet. ether /EtOAc = 85/15): 0.50; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.23 (d, J = 8.6 Hz, 2H), 7.79 (d, J = 8.8 Hz, 2H), 7.64 (d, J = 7.7 Hz, 1H), 7.55 (d, J = 8.6 Hz, 2H), 7.44-7.32 (m, 4H), 7.21-7.16 (m, 2H), 6.66 (s, 1H), 6.08 (s, 1H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 154.93, 148.08, 147.21, 141.22, 135.92, 134.33, 129.35, 129.10, 128.58, 128.22, 127.19, 125.95, 124.24, 122.07, 120.31, 97.60, 86.13. **HRMS (ESI)** calculated $[\text{M}+\text{H}]^+$ for $\text{C}_{21}\text{H}_{16}\text{NO}_3$: 330.1130, found: 330.1125. **FTIR** (cm^{-1}) 3052, 1659, 1523, 1349, 1107, 1038, 852.

(Z)-1-benzylidene-3-(4-nitrophenyl)-1,3-dihydroisobenzofuran 8b

Yellow solid (5 mg, 15% yield), R_f (Pet. ether /EtOAc = 90/10): 0.60; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.20 (d, J = 8.6 Hz, 2H), 7.75-7.73 (m, 2H), 7.57 (d, J = 8.6 Hz, 2H), 7.42-7.32 (m, 4H), 7.24-7.19 (m, 2H), 6.87 (d, J = 7.7 Hz, 1H), 6.53 (s, 1H), 6.37 (s, 1H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 152.14, 147.95, 147.18, 133.94, 131.48, 129.28, 129.05, 129.00, 128.60, 128.58, 126.99, 125.26, 125.08, 124.41, 123.72, 101.41, 78.89. **HRMS (ESI)** calculated $[\text{M}+\text{H}]^+$ for $\text{C}_{21}\text{H}_{16}\text{NO}_3$: 330.1130, found: 330.1125. **FTIR** (cm^{-1}) 3072, 3061, 2365, 1521, 1346, 1219, 1062, 855.

Using Heck Coupling:

(*E*)-(4-Nitrophenyl)(2-styrylphenyl)methanol (**9a**)

Following the literature procedure,¹⁰ a mixture of (2-iodophenyl)(4-nitrophenyl)methanol **3a** (44 mg, 0.125 mmol), styrene (26 mg, 29 μ L, 0.25 mmol), palladium acetate (2.8 mg, 0.0125 mmol), tri-*o*-tolylphosphine (7.6 mg, 0.025 mmol) and triethylamine (51 mg, 70 μ L, 0.5 mmol), in acetonitrile (1.25 ml) was heated at 80 °C for 12 h. The reaction mixture was concentrated under reduced pressure and then purified by flash column chromatography to afford (*E*)-(4-nitrophenyl) (2-styrylphenyl)methanol **9a** as a yellow oil (35 mg, 84%).

R_f (Pet. ether /EtOAc = 85/15): 0.39; $^1\text{H NMR}$ (400 MHz, CDCl_3) 8.17 (d, J = 8.8 Hz, 2H), 7.64 (d, J = 7.6 Hz, 1H), 7.56 (d, J = 8.6 Hz, 2H), 7.44-7.27 (m, 9H), 6.95 (d, J = 16 Hz, 1H), 6.26 (s, 1H), 2.53 (s, 1H). $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 150.53, 147.21, 139.77, 137.10, 136.32, 132.22, 128.92, 128.22, 127.85, 127.38, 127.00, 126.72, 125.33, 123.74, 72.86. **HRMS (ESI)** calculated $[\text{M}-\text{OH}]^+$ for $\text{C}_{21}\text{H}_{16}\text{NO}_2$: 314.1176, found: 314.1180. **FTIR** (cm^{-1}) 3583, 3025, 2364, 1600, 1518, 1344, 1027, 962, 850.

Using Suzuki Coupling

[1,1'-biphenyl]-2-yl(4-nitrophenyl)methanol (**10a**)

Following the literature procedure,¹¹ (2-iodophenyl)(4-nitrophenyl)methanol **3a** (44 mg, 0.125 mmol), phenylboronic acid (23 mg, 1.5 equiv), $\text{Pd}(\text{PPh}_3)_4$ (4.3 mg, 0.03equiv), Na_2CO_3 (40 mg, 3 equiv) was added into a flame-dried screw-capped test tube equipped with a magnetic stir bar. The screw-capped test tube was evacuated under vacuum and flushed with dry nitrogen three times and then 2 mL THF and 0.5 mL water was added. The reaction mixture was refluxed for 24 h. After cooling to room temperature, the mixture was poured into water and

¹⁰ Hirano, K.; Biju, A. T.; Piel, I.; Glorius, F. *J. Am. Chem. Soc.* **2009**, *131*, 14190.

¹¹ Chen, L.; Wang, Y.-H.; He, B.; Nie, H.; Hu, R.; Huang, F.; Qin, A.; Zhou, X.-S.; Zhao, Z.; Tang, B. Z. *Angew. Chem., Int. Ed.* **2015**, *54*, 4231.

extracted with dichloromethane three times. The combined organic layers were dried over anhydrous Na_2SO_4 . After filtration and solvent evaporation, a silica gel flash column chromatography (Pet. ether /EtOAc = 90/10) of the crude reaction mixture afforded [1,1'-biphenyl]-2-yl(4-nitrophenyl)methanol **10a** as a yellow liquid (30 mg, 79% yield).

R_f (Pet. ether /EtOAc = 85/15): 0.40; **^1H NMR (400 MHz, CDCl_3)** δ 8.10-8.06 (m, 2H), 7.43-7.35 (m, 6H), 7.34-7.25 (m, 5H), 6.02 (s, 1H), 2.33 (bs, 1H). **^{13}C NMR (100 MHz, CDCl_3)** δ 151.20, 147.07, 141.61, 140.50, 140.16, 130.47, 129.38, 128.58, 128.41, 128.25, 127.70, 127.35, 123.53, 71.84. **HRMS (ESI)** calculated $[\text{M}-\text{OH}]^+$ for $\text{C}_{19}\text{H}_{14}\text{NO}_2$: 288.1019, found: 288.1028. **FTIR (cm^{-1})** 3418, 3061, 2363, 1601, 1519, 1345, 1011, 839.

10. ^1H and ^{13}C NMR Spectra of Products

(2-Iodophenyl)(4-nitrophenyl)methanol (**3a**)

(2-Iodophenyl)(phenyl)methanol (3b)¹

(4-Bromophenyl)(2-iodophenyl)methanol (3c)

(4-Chlorophenyl)(2-iodophenyl)methanol (3d)

(4-Fluorophenyl)(2-iodophenyl)methanol (3e)

[1,1'-Biphenyl]-4-yl(2-iodophenyl)methanol (3f)

(2-Iodophenyl)(4-(trifluoromethyl)phenyl)methanol (3g)

4-(Hydroxy(2-iodophenyl)methyl)benzonitrile (3h)

Methyl-4-(hydroxy(2-iodophenyl)methyl)benzoate (3i)

4-(Hydroxy(2-iodophenyl)methyl)benzaldehyde (3j)

(3-Bromophenyl)(2-iodophenyl)methanol (3k)

(2-Fluorophenyl)(2-iodophenyl)methanol (3I)

(2-Chlorophenyl)(2-iodophenyl)methanol (3m)

(2-Iodophenyl)(2-nitrophenyl)methanol (3n)

(3,4-Dichlorophenyl)(2-iodophenyl)methanol (3o)

(2-Chloro-6-fluorophenyl)(2-iodophenyl)methanol (3p)

(3,5-Bis(trifluoromethyl)phenyl)(2-iodophenyl)methanol (3q)

(2-Iodophenyl)(naphthalen-2-yl)methanol (3r)

(2-Iodophenyl)(pyren-1-yl)methanol (3s)

(2-Iodophenyl)(thiophen-2-yl)methanol (3t)

Benzofuran-2-yl(2-iodophenyl)methanol (3u)

Cyclohexyl(2-iodophenyl)methanol (3v)

1-(2-Iodophenyl)-2-methylpropan-1-ol (3w)

(2-Iodo-4,5-dimethylphenyl)(4-nitrophenyl)methanol (3x)

(6-Iodobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol (3y)

(2-Iodo-3,6-dimethylphenyl)(4-nitrophenyl)methanol (3z)

(3-Iodonaphthalen-2-yl)(4-nitrophenyl)methanol (3aa)

(2-Iodonaphthalen-1-yl)(4-nitrophenyl)methanol (3ab)

(2-Iodo-6-methoxyphenyl)(4-nitrophenyl)methanol (3ac)

(2-Iodo-4-methylphenyl)(4-nitrophenyl)methanol and (2-Iodo-5-methylphenyl)(4-nitrophenyl)methanol (3ad)

(2-Bromophenyl)(4-nitrophenyl)methanol (3ae)

The figure displays the ^1H and ^{13}C NMR spectra of compound **3af**, which is 2-chloro-2-(4-nitrophenyl)-1-phenylethanol. The chemical structure is shown as an inset in both spectra.

^1H NMR Spectrum (Top): The spectrum shows peaks in the aromatic region (7.2–7.6 ppm) and a broad peak for the hydroxyl group (1.5–2.0 ppm). Integration values are provided below the peaks.

Chemical Shift (ppm)	Integration
7.60, 7.58, 7.50, 7.49, 7.48	2.16
7.39, 7.37, 7.31	1.05
7.29, 7.29, 7.28, 7.28, 7.27, 7.26	0.97
7.26, 7.26	2.38
2.63, 2.62	1.06
1.5–2.0	1.00

^{13}C NMR Spectrum (Bottom): The spectrum shows peaks in the aromatic region (123–149 ppm) and a cluster of peaks for the solvent (71–77 ppm).

Chemical Shift (ppm)
149.40, 147.28, 140.00, 132.43, 129.83, 129.52, 128.19, 127.55, 127.53, 123.69
77.38, 77.06, 76.74, 71.71

(2-Bromophenyl)(phenyl)methanol (3ag)

(2-Chlorophenyl)(phenyl)methanol (3ah)

Benzofuran-2-yl(2-bromophenyl)methanol (3ai)

Benzofuran-2-yl(2-chlorophenyl)methanol (3aj)

1-(2-Bromophenyl)-2-methylpropan-1-ol (3ak)

(6-Bromobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol (3al)

(6-Chlorobenzo[d][1,3]dioxol-5-yl)(4-nitrophenyl)methanol (3am)

3-Hydroxy-3-(2-iodophenyl)-1-methylindolin-2-one (5a)

2-Iodobenzoic acid (6a)

(4-Nitrophenyl)(2-(phenylethynyl)phenyl)methanol (7a)

(Z)-1-Benzylidene-3-(4-nitrophenyl)-1,3-dihydroisobenzofuran (8a)

(Z)-1-Benzylidene-3-(4-nitrophenyl)-1,3-dihydroisobenzofuran (8b)

(E)-(4-Nitrophenyl)(2-styrylphenyl)methanol (9a)

[1,1'-Biphenyl]-2-yl(4-nitrophenyl)methanol (10a)