PAPER

Does Hubble tension signal a breakdown in FLRW cosmology?

To cite this article: C Krishnan *et al* 2021 *Class. Quantum Grav.* **38** 184001

View the [article online](#) for updates and enhancements.

IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Does Hubble tension signal a breakdown in FLRW cosmology?

C Krishnan¹ , R Mohayaee², E Ó Colgáin^{3,4,*} ,
M M Sheikh-Jabbari⁵ and L Yin^{3,4}

¹ Center for High Energy Physics, Indian Institute of Science, Bangalore 560012, India

² Sorbonne Université, CNRS, Institut d'Astrophysique de Paris, 98bis Bld Arago, Paris 75014, France

³ Center for Quantum Spacetime, Sogang University, Seoul 121-742, Republic of Korea

⁴ Department of Physics, Sogang University, Seoul 121-742, Republic of Korea

⁵ School of Physics, Institute for Research in Fundamental Sciences (IPM), PO Box 19395-5531, Tehran, Iran

E-mail: chethan.krishnan@gmail.com, mohayaee@iap.fr, eoin@sogang.ac.kr,
shahin.s.jabbari@gmail.com and yinlu@sogang.ac.kr

Received 15 June 2021, revised 13 July 2021

Accepted for publication 4 August 2021

Published 23 August 2021

CrossMark

Abstract

The tension between early and late Universe probes of the Hubble constant has motivated various new FLRW cosmologies. Here, we reanalyse the Hubble tension with a recent age of the Universe constraint. This allows us to restrict attention to matter and a dark energy sector that we treat without assuming a specific model. Assuming analyticity of the Hubble parameter $H(z)$, and a generic low redshift modification to flat Λ CDM, we find that low redshift data ($z \lesssim 2.5$) and well-motivated priors only permit a dark energy sector close to the cosmological constant Λ . This restriction rules out late Universe modifications within FLRW. We show that early Universe physics that alters the sound horizon can yield an upper limit of $H_0 \sim 71 \pm 1 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Since various local determinations may be converging to $H_0 \sim 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$, a breakdown of the FLRW framework is a plausible resolution. We outline how future data, in particular strongly lensed quasar data, could also provide further confirmations of such a resolution.

Keywords: Hubble, tension, FLRW, cosmology

(Some figures may appear in colour only in the online journal)

* Author to whom any correspondence should be addressed.

‘How often have I said to you that when you have eliminated the impossible, whatever remains, however improbable, must be the truth?’—Sherlock Holmes

1. Introduction

A cursory glance at figure 1 of [1] reveals almost all local H_0 determinations are biased higher than Planck- Λ CDM [2]. This observation is agnostic about whether one calibrates type Ia supernovae with Cepheids [3–6] or TRGB [7–10]. If true, this has profound consequences. In particular, one can argue that dark energy (DE), a feature in multiple datasets—type Ia supernovae [11, 12], cosmic microwave background (CMB) [2] and baryon acoustic oscillations (BAO) [13]—cannot be described by a cosmological constant or by a minimally coupled scalar field theory [14–16]. Simply put, *uncoupled* quintessence models [18, 19] may already be ruled out by the Hubble tension [16]. This may leave only *coupled* quintessence models as viable options, e.g. [20–22]. The key point is that Hubble tension appears to be extremely challenging for late time DE.

Recall the age of Universe t_U within the Friedmann–Lemaître–Robertson–Walker (FLRW) paradigm:

$$t_U = \frac{977.8}{H_0} \int_0^\infty \frac{1}{(1+z)E(z)} dz \text{ Gyr}, \quad (1.1)$$

where H_0 is the Hubble constant and $E(z)$ denotes the normalised Hubble parameter. Since $E(z)$ increases as $E(z) \sim (1+z)^p$, $p \geq 3/2$ in the early Universe, the contribution of the early Universe physics to the integral is negligible. In other words, a competitive determination of t_U consistent with Planck, necessitates a late Universe modification to Planck- Λ CDM [17].

Since the high redshift contribution to t_U is essentially fixed by $p = 3/2$ and $\omega_m := \Omega_{m0}h^2$, which is in turn constrained by CMB, any alternative FLRW cosmology motivated by Hubble tension requires $H(z)$ higher than Planck at $z \approx 0$, but lower at cosmological redshifts. This can be interpreted as a ‘running of H_0 ’ with redshift [23]. More concretely, given two FLRW cosmologies, e.g. flat Λ CDM $H_{\Lambda\text{CDM}}(z)$ and a replacement $H_{\text{new}}(z)$, provided $H(z)$ is observationally determined, then $H_0^{\Lambda\text{CDM}}/H_0^{\text{new}} \sim E(z)_{\text{new}}/E(z)_{\Lambda\text{CDM}}$ must hold. Only when the two models are identical the left-hand side can remain a constant. Such a running must follow if Hubble tension can find a resolution within FLRW [23]. Tentative hints of such a feature already exist [24–28].

In this letter, given a recent competitive determination for the Universe’s age from globular clusters $t_U = 13.5 \pm 0.27$ Gyr [17, 29], we explore the possibility of the required late time modification, namely a higher H_0 , with lower $E(z)$ at redshifts $z \lesssim 2.5$. To be as model independent as possible we employ overlapping Taylor expansions about small and large z . The first quantifies our ignorance of the DE sector, while the second is fixed by CMB determinations of ω_m . We reduce the dependence of the latter on the DE sector by removing lower CMB multipoles $\ell \lesssim 30$ [30–33]. We also make use of a ‘SH0ES prior’ on the absolute magnitude of type Ia supernovae M_B [34] (see also [35]). It is worth noting that a prior on H_0 is stronger than a prior on M_B , as the former affects all the cosmological data. Our use of a SH0ES prior, in contrast to a TRGB prior, is supported by a number of other independent local determinations that favour $H_0 > 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. These include megamasers [36], surface brightness fluctuations [37] and Tully–Fisher relation [38], which are either completely independent of the Cepheid distance ladder, or show little variation in H_0 across Cepheids and TRGB.

This constitutes a minimal setup covering a matter sector and a late time deceleration–acceleration transition, without invoking any specific DE model. However, there is a loose end, namely the radius of the sound horizon r_d . When treated as a free parameter, we can allow

putative early Universe physics that alters r_d [39–43]. We will also fix it through a prior [33] and take the early Universe to be flat Λ CDM. While our work overlaps with earlier studies [44, 45], our approach is analytic, it is relatively model-agnostic and leads to an upper bound on an achievable Hubble constant, $H_0 \sim 71 \pm 1 \text{ km s}^{-1} \text{ Mpc}^{-1}$, which represents a competitive 1.4% error.

Overall, we find that any freedom given to the DE sector in the late Universe is removed by the data and priors: our best-fit DE sector is more or less Λ . From this observation, results follow. First, the data *must* adjust r_d in any attempt to accommodate a SH0ES prior. Unsurprisingly, we find no running in H_0 and this disfavors a resolution to Hubble tension in the DE sector $z \lesssim 2.5$, but leaves an early Universe modification on the table for $H_0 \sim 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ or lower. Higher values, e.g. $H_0 \sim 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [6], are consistent within 2σ , but the discrepancy in central values allows room for other explanations. An alternative scenario is the breakdown of FLRW itself. In section 6 we argue that H_0 values inferred from lensed quasars [24, 25] may provide the first glimpse of the validity of this scenario.

2. Setup

It is assumed that the Universe is dominated by pressure-less dust that is well-approximated by the equation of state (EOS), $w = 0$. Furthermore, we have evidence [11, 12] that the Universe goes through a deceleration–acceleration transition at late times. Therefore, at low redshifts, the minimal model contains two sectors, namely DE and matter, with a Hubble parameter:

$$H(z) = H_0 \sqrt{(1 - \Omega_{m0})f(z) + \Omega_{m0}(1 + z)^3}. \quad (2.1)$$

Observe that other putative sectors currently constitute unmotivated speculation. As remarked in [23], H_0 is an integration constant and this fixes the overall scale in any FLRW cosmology. Moreover, the Friedmann equation also determines the rhs of (2.1) subject to the matter EOS, $w = 0$. Finally, the unknown function $f(z)$ describes the DE sector, but consistency demands that $f(z = 0) = 1$. The flat Λ CDM model corresponds to $f(z) = 1$.

To study DE contribution it is convenient to write $H(z)$ as⁶,

$$H(z) = H_0 \sqrt{\Omega_{m0}} (1 + z)^{\frac{3}{2}} \sqrt{1 + X}, \quad (2.2)$$

$$X := \frac{\Omega_{\Lambda}(z)}{\Omega_m(z)} = \frac{(1 - \Omega_{m0})f(z)}{\Omega_{m0}(1 + z)^3}. \quad (2.3)$$

Now, if $X(z)$ is small and decreasing $X'(z) < 0$, we can treat it perturbatively at higher redshifts. Recall that by definition $\ddot{a}/a = \dot{H} + H^2$. As a result, when one moves from deceleration ($\ddot{a} < 0$) to acceleration ($\ddot{a} > 0$), necessarily $\dot{H} + H^2 = 0$ at the transition redshift z_t . We can get an idea for the size of $X(z_t)$ in the Chevallier–Polarski–Linder (CPL) parametrisation [51, 52], where concretely

$$f(z) = (1 + z)^{3(1+w_0+w_a)} e^{-\frac{3w_a z}{1+z}}. \quad (2.4)$$

To do this, one imports Planck Markov Chain Monte Carlo (MCMC) chains [2] and solves for $\ddot{a} = 0$ to identify z_t for each entry in the MCMC chain. One can see from figure 1, where

⁶ There are strong constraints on ‘early dark energy’ [46, 47] from CMB and BBN [48–50], but one can still play with the idea pre-recombination [39, 42, 43].

Figure 1. Distribution of the transition redshift z_t and $X(z_t)$ inferred from Planck MCMC chains of CMB + BAO + supernovae constraints on the CPL model.

we have used *getdist* [53], that in relaxing the DE model away from Λ , the $(z_t, X(z_t))$ values are still close to the Planck- Λ CDM values, $(z_t, X(z_t)) \approx (0.67, 0.5)$. The key point is that $X(z > z_t) < X(z_t) < 1$, so we can treat it perturbatively⁷.

3. Strategy

Following [16] we work perturbatively at low redshift by Taylor expanding $H(z)$ about $z = 0$. It is well documented that one cannot go beyond $z = 1$ [55] (see also appendix A of [56]), so here we expand up to an intermediate redshift $z_* \sim z_t$. Now, we can Taylor expand the unknown function $f(z)$,

$$f(z) = 1 + f_1 z + \frac{f_2}{2!} z^2 + \frac{f_3}{3!} z^3 + \dots, \quad (3.1)$$

where we have defined $f_n \equiv f^{(n)}(0)$. Adopting this expansion, $H(z)$ for $z < z_*$ becomes,

$$H(z) = H_0 \left(1 + h_1 z + h_2 z^2 + h_3 z^3 + \dots \right), \quad (3.2)$$

where we have further defined

$$\begin{aligned} h_1 &= \frac{3}{2} \Omega_{m0} + \frac{1}{2} (1 - \Omega_{m0}) f_1, \\ h_2 &= \frac{3}{2} \Omega_{m0} - \frac{1}{8} (3 \Omega_{m0} + (1 - \Omega_{m0}) f_1)^2 + \frac{1}{4} (1 - \Omega_{m0}) f_2, \end{aligned}$$

⁷ That z_t and $X(z_t)$ do not change much within the CPL model may not be so surprising, since CPL is biased towards Λ [54].

$$h_3 = -\frac{1}{4}(3\Omega_{m0} + (1 - \Omega_{m0})f_1) \left(3\Omega_{m0} - \frac{1}{4}(3\Omega_{m0} + (1 - \Omega_{m0})f_1)^2 \right. \\ \left. + \frac{1}{2}(1 - \Omega_{m0})f_2 \right) + \frac{1}{2} \left(\Omega_{m0} + \frac{1}{6}(1 - \Omega_{m0})f_3 \right). \quad (3.3)$$

We restrict the presentation to third order as it approximates flat Λ CDM well to z_t . Nevertheless, f_3 is degenerate with Ω_{m0} , so one can safely set $f_3 = 0$.⁸

To appreciate this fact, observe that the term in the square root in (2.2) can be expanded to third order in z as

$$(1 - \Omega_{m0})f(z) + \Omega_{m0}(1 + z)^3 = 1 + (1 - \Omega_{m0}) \left(f_1 z + \frac{f_2}{2!} z^2 + \frac{f_3}{3!} z^3 \right) \\ + \Omega_{m0}(3z + 3z^2 + z^3). \quad (3.4)$$

However, one can absorb f_3 into the definition of a new Ω'_{m0} , while also redefining f_1 and f_2 :

$$\Omega'_{m0} = \Omega_{m0} + \frac{f_3}{3!}(1 - \Omega_{m0}), \\ f'_2 = \frac{6(f_3 - f_2)}{f_3 - 6}, \quad f'_1 = \frac{3(f_3 - 2f_1)}{f_3 - 6}. \quad (3.5)$$

The end result is the original expression with $f_3 = 0$. This degeneracy between Ω_{m0} and f_3 allows one to set $f_3 = 0$ in (3.2).

We assume that (3.1) is a valid perturbative expression in the sense that higher order terms are smaller, i.e. $|f_n|z < n|f_{n-1}|$ in the redshift range of interest $z < z_*$. Recalling that a number of late-time resolutions to Hubble tension correspond at the background level to a w CDM model with $w \approx -1.2$ [58] (see also [59–61]), it is useful to note the expansion of (2.4) with representative values: $f(z)_{w=-1.2} = 1 - 0.6z + 0.48z^2 + O(z^3)$.

The idea now is to use (3.2) below $z = z_*$, but to use the following expression for $z \geq z_*$:

$$H(z) = H_0 \sqrt{\Omega_{m0}} (1 + z)^{\frac{3}{2}} \left(1 + \frac{1}{2}X - \frac{1}{8}X^2 + \dots \right),$$

where throughout we expand both $H(z)$ and $H(z)^{-1}$ to $O(X^{10})$. In line with expectations, the low redshift expansion (3.2) starts to deviate from the exact result before $z = 1$, whereas the high redshift expansion can be pushed close, but not as far as $z = 0$. The low redshift expansion performs worse, yet the error is within 1% for $\Omega_{m0} = 0.3$ through to $z_* \sim 0.8$. In contrast, the error in the high redshift expansion is negligible. Note that z_* constitutes a redundancy in our analysis, but as we show later (table 3), it does not change our results.

4. Data sets

We make use of the Pantheon supernovae dataset [62], a rich compilation of BAO data [63–73], as presented in table 1, and cosmic chronometer data [74]. To flesh out the analysis we employ

⁸ Our expansion covers interacting DE models which can be written as $w(z)$ CDM, e.g. [57]. However, curvature Ω_k is strictly speaking not covered by our definitions of distances, e.g. $d_L(z)$, $d_A(z)$. More precisely, curvature can be absorbed into f_i in $H(z)$, but it affects distances. However, this change will not be significant if the curvature is small, $\Omega_k \approx 0$.

Table 1. BAO data.

	z_{eff}	Constraint	Reference
6dF	0.106	$D_V/r_d = 2.98 \pm 0.13$	[63]
SDSS MGS	0.15	$D_V/r_d = 4.47 \pm 0.17$	[64]
		$D_A/r_d = 7.41 \pm 0.12$	
BOSS Galaxy	0.38	$D_H/r_d = 25.00 \pm 0.76$	[65]
		$D_A/r_d = 8.85 \pm 0.14$	
	0.51	$D_H/r_d = 22.38 \pm 0.58$	[65]
DES	0.81	$D_A/r_d = 10.75 \pm 0.43$	[66]
		$D_A/r_d = 10.51 \pm 0.19$	
eBOSS LRG	0.70	$D_H/r_d = 19.33 \pm 0.53$	[67, 68]
eBOSS ELG	0.85	$D_V/r_d = 18.33^{+0.57}_{-0.62}$	[69, 70]
		$D_A/r_d = 12.38 \pm 0.32$	
eBOSS Quasar	1.48	$D_H/r_d = 13.26 \pm 0.55$	[71, 72]
		$D_A/r_d = 11.29 \pm 0.57$	
Ly α -Ly α	2.33	$D_H/r_d = 8.93 \pm 0.28$	[73]
		$D_A/r_d = 11.20 \pm 0.51$	
Ly α -Quasar	2.33	$D_H/r_d = 9.08 \pm 0.34$	[73]

Table 2. Best fit values for the flat Λ CDM model with an t_U prior and a low multipole-subtracted prior on ω_m .

H_0 (km s $^{-1}$ Mpc $^{-1}$)	Ω_{m0}	r_d (Mpc)	M_B (mag)
$69.56^{+1.25}_{-1.23}$	$0.295^{+0.013}_{-0.012}$	$144.8^{+2.2}_{-2.0}$	$-19.37^{+0.03}_{-0.03}$

of a number of priors. First, we assume that $t_U = 13.50 \pm 0.27$ Gyrs [17]. Next, we employ the high redshift prior $\omega_m = 0.141 \pm 0.006$, where the lower multipoles have been removed to reduce dependence on the DE sector (see appendix A). Finally, we utilise a SH0ES prior on the absolute magnitude, $M_B = -19.244 \pm 0.042$ mag [34], which is consistent with [35], and an early Universe prior $r_d = 147.4 \pm 0.7$ [33]. Note that we adapt *a prior* on M_B and a H_0 prior would be stronger than an M_B prior.

5. Results

Consider first the flat Λ CDM model subject to the t_U prior and r_d, M_B treated as nuisance parameters. The best-fit values are shown in table 2. Evidently, H_0 is higher than Planck [2], but this is not so surprising since we have adopted a t_U prior. From our MCMC chains, where we used *emcee* [75], we find that $t_U = 13.62 \pm 0.16$ Gyr, while $\omega_m = 0.143 \pm 0.004$, which are both consistent with the priors. The higher H_0 appears to be driven by the t_U prior, while the nuisance parameters r_d and M_B come along for the ride. Nevertheless, r_d is consistent with the Planck value $r_d \sim 147$ Mpc at 1σ .

The flat Λ CDM fit simply serves to get oriented. Next, we repeat, but now let the DE sector vary through the f_i (3.1) and impose a SH0ES prior on M_B [34]. The best-fit values can be found in table 3, where we record results both with and without a prior on r_d [33]. Observe also that we have scanned over z_* , and as expected, it does not change our results. Throughout we find that H_0 , Ω_{m0} and t_U are all fully consistent with the imposed priors.

Table 3. Best fit values for our overlapping expansions with a t_U prior, low multipole-subtracted prior on ω_m and a SH0ES prior on M_B . In the lower entries for each z_* we impose a prior on r_d .

z_*	H_0 (km s ⁻¹ Mpc ⁻¹)	Ω_{m0}	f_1	f_2	r_d (Mpc)	M_B (mag)
0.5	$70.97^{+1.02}_{-1.00}$	$0.284^{+0.012}_{-0.010}$	$0.15^{+0.13}_{-0.17}$	$-0.08^{+0.25}_{-0.32}$	$141.88^{+1.8}_{-1.9}$	$-19.32^{+0.03}_{-0.03}$
	$69.34^{+0.83}_{-0.79}$	$0.292^{+0.013}_{-0.012}$	$0.08^{+0.14}_{-0.22}$	$-0.10^{+0.26}_{-0.32}$	$146.8^{+0.7}_{-0.6}$	$-19.37^{+0.02}_{-0.02}$
0.6	$71.18^{+1.04}_{-1.04}$	$0.279^{+0.013}_{-0.013}$	$0.17^{+0.13}_{-0.13}$	$-0.02^{+0.25}_{-0.29}$	$141.3^{+2.0}_{-1.9}$	$-19.31^{+0.03}_{-0.03}$
	$69.39^{+0.85}_{-0.83}$	$0.291^{+0.013}_{-0.013}$	$0.07^{+0.15}_{-0.23}$	$-0.06^{+0.25}_{-0.30}$	$146.7^{+0.7}_{-0.7}$	$-19.37^{+0.02}_{-0.02}$
0.7	$71.15^{+1.03}_{-1.03}$	$0.279^{+0.013}_{-0.013}$	$0.18^{+0.13}_{-0.12}$	$-0.03^{+0.22}_{-0.28}$	$141.2^{+2.0}_{-1.8}$	$-19.31^{+0.03}_{-0.03}$
	$69.38^{+0.84}_{-0.84}$	$0.290^{+0.013}_{-0.012}$	$0.07^{+0.15}_{-0.23}$	$-0.05^{+0.22}_{-0.27}$	$146.7^{+0.6}_{-0.6}$	$-19.37^{+0.02}_{-0.02}$

Figure 2. The ratio between the normalised Hubble parameters for our cosmological model (2.1) and flat Λ CDM with a prior on r_d and $z_* = 0.6$. The shaded blue and red envelopes are the 1σ confidence intervals, while the grey region is the overlap. That this ratio is basically a constant confirms there is no H_0 running in the sense discussed in [23].

It is worth noting that with the r_d prior, both f_1 and f_2 are consistent with their flat Λ CDM values within 1σ . There is clearly no freedom in the DE sector. The likely explanation for this is that our prior on ω_m leaves little freedom for non-zero f_i within the constraints imposed by BAO and supernovae on $H(z)$. Moreover, observe that the constraints on r_d and t_U lead to a value of M_B that is the same as the flat Λ CDM fit (table 2). Tellingly, the final M_B value is 2.7σ away from the original prior and this underscores the tension between M_B and r_d .

In figure 2 we offer visual confirmation that the reconstructed $E(z)_{\text{rec}}$ does not differ enough from flat Λ CDM $E(z)_{\Lambda\text{CDM}}$. There is no evidence for running in H_0 with redshift [23] as is clear from the 1σ confidence intervals for the ratio presented. For completeness, we compare to flat Λ CDM defined with respect to our MCMC chains (table 2) and the Planck MCMC chains [2]. The asymptotic, large z behaviour is governed by the best-fit values of Ω_{m0} .

Finally, when r_d is treated as a free parameter, f_2 is consistent with zero, but f_1 shows some marginal departure, which may be described by the simplest extension, the w CDM model. In

Figure 3. Best-fit values of H_0 with and without a prior on r_d . The purple and green bands represents 1σ from Planck and SH0ES [6], respectively.

essence, there may be some slight deviation from Λ in the DE sector, as suggested in [17]. However, the maximum H_0 value achievable within an FLRW cosmology coupled to Einstein gravity, regardless of the DE sector, is $H_0 \sim 71 \pm 1 \text{ km s}^{-1} \text{ Mpc}^{-1}$. We illustrate the difference the prior makes in figure 3, where the tension between SH0ES and r_d with a flat Λ CDM prior is evident. The figure is plotted for $z_* = 0.6$, but z_* does not affect the results (see table 3). This figure, although at low significance, clearly demonstrates that the early Universe solutions within FLRW come short of solving the Hubble tension.

6. Going beyond FLRW

H0LiCOW [24] has reported a descending trend of H_0 with lens redshift in the flat Λ CDM model and an extra time delay measurement was also obtained from DES [76]. This is interesting since all lenses should be subject to the same assumptions, so if there is a systematic, then all lenses are affected. Nevertheless, the descending trend in H_0 can be expected to be robust [25] and it has prompted various explanations [23, 77]. However, it is possible that it is not a trend with lens redshift, but due to the line of sight (LOS) anisotropy as we outline here.

Although the effect of local anisotropies on lensing is marginal, if these anisotropies extend to a far larger extent than predicted by LCDM model, as shown by bulk flow data, then they could have non-negligible impact on the lensing analyses. Indeed, a large over-density along the LOS will increase the estimated value of H_0 (by increasing the value of κ_{ext} , see e.g. [78] for details).

To begin, instead of focussing on the dependence of H_0 values on lens redshift, as is done in [24, 25], we study their angular distribution on the sky. In table 4 we present the coordinates of the H0LiCOW and DES lenses in the equatorial coordinate system where α and δ denote right ascension (RA) and declination (DEC), respectively. It is intriguing to compare the orientations of these few quasars with the CMB dipole [2], $\alpha = 168^\circ$, $\delta = -7^\circ$, approximately along which most of the superclusters in the local Universe are lined up. This anisotropy, if extended to larger scales as shown by non-convergence of the CMB dipole, is expected to effect the LOS contribution to the time-delay. The LOS contribution has been taken into account within certain models, but to our knowledge the effect of anisotropies or the aberration effect due to large velocity of the observer with respect to distant quasars has not been reported in these studies.

Table 4. Location of lenses on the celestial sphere from H0LiCOW and DES (last data point).

Lens	α	δ	H_0 (km s ⁻¹ Mpc ⁻¹)
B1608 + 656	242°	+66°	71.0 ^{+2.9} _{-3.3}
RXJ1131-1231	173°	-13°	78.2 ^{+3.4} _{-3.4}
HE 0435-1223	70°	-12°	71.7 ^{+4.8} _{-4.5}
SDSS 1206 + 4332	182°	+44°	68.9 ^{+5.4} _{-5.1}
WFI2033-4723	308°	-47°	71.6 ^{+3.8} _{-4.9}
PG 1115 + 080	170°	+8°	81.1 ^{+8.0} _{-7.1}
DES J0408-5354	62°	-54°	74.2 ^{+2.7} _{-3.0}

Figure 4. Sky position, in RA and DEC, of the lenses from the H0LiCOW sample and DES are shown on an Aitoff plot together with the direction of the CMB dipole. The inferred values of H_0 for each lens are shown next to the lens position. The data points are overlaid on a density plot of mass distribution in the local Universe (Virgo, Great Attractor, Shapley, Ophiuchus, Coma, Pavo-Indus, Hercule, Norma, Perseus-Pisces, Horologium and the local void are included). The large DES value is near the Horologium supercluster and the two large ones from H0LiCOW are near most other superclusters including Shapley. The supercluster on the top left is Perseus-Pisces. The dark blue region on the top right quadrant is the local void. GP is not shown.

Although we currently only have a tiny dataset of seven lensed quasars, we clearly see that the two lenses that are closest to the direction of the CMB dipole, namely RXJ1131-1231 and PG 1115 + 080, exhibit the highest values of H_0 . Furthermore, a large value of H_0 is inferred from DES data, which lies close to the direction of the Horologium supercluster. This may not be a coincidence or a simple fluctuation. A similar study has been carried out on the anisotropy of H_0 using supernovae, which due to poor data yielded a low-significant result [107]. Clearly these data are significantly impacted by the degeneracy in the mass distribution modelling of the lenses, however future similar data on time delay from LSST shall certainly provide us with further confirmations.

It is often assumed that the origin of the CMB dipole is the gravitational attraction of the local superclusters. However it seems that the observed superclusters up to the Shapley concentration cannot fully account for the CMB dipole, which consequently cannot be of purely kinematic origin (e.g. see [112] and references therein). Furthermore, since the sources in this dataset are all at high redshifts, then they might signal a large structural clustering well beyond what is permitted in an FLRW Universe. It is thus expected that the seemingly far-reaching

local anisotropy has some signature on the strongly lensed quasars and this indeed appears to be the case. The correlation between H_0 from the lensed quasars and distribution of most-prominent superclusters is suggestive even for such a tiny dataset and deserves further study as data improves.

We make one last comment before closing the section. We emphasise again that it is the relative differences in H_0 that interest us from the lensed quasars. In order to read into the correlations in figure 4, it is imperative that all lenses have been modelled uniformly. As is clear from recent results [79], the *absolute* value of H_0 inferred from strong lensing time delay depends on assumptions for mass sheet modelling, so for this reason we have not employed it as a prior in our analysis.

7. Discussion

In this letter we have explored a modification of the DE sector consistent with a recent determination of the age of Universe, t_U [17, 29]. As explained, a fixed t_U consistent with Planck results allows a higher H_0 , but lower $E(z)$ relative to Planck- Λ CDM, thus providing a realisation of a running H_0 [23]. Ultimately, the rigidity of the DE sector precludes this. As a result, data has a preference for an early Universe resolution, but only as high as $H_0 \sim 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Of course, this assumes a working model can be found, ideally something natural that penalises additional unobserved degrees of freedom, (see e.g. [80]), which is currently far from clear [81–88]. Note, our analysis in principle captures all early Universe resolutions, since r_d is a free parameter.

What is new here is that our analysis is minimal, yet covers (w_0, w_a) parameterisations of DE (see [89]), while the H_0 errors are meaningful. As is clear from table 3, we have a 1.4% error on $H_0 \sim 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ without assuming a prior on r_d . While this is only 1.2σ discrepant with SH0ES [6], once local determinations are combined, for example [90] or [91], this inches up to 1.4σ and 1.7σ , respectively. Putting aside caveats about combining data, admittedly, even as data improves, it is hard to imagine any better than a $\sim 2\sigma$ discrepancy with SH0ES. Of course, if claims of new early Universe physics are challenged by forthcoming results from ACT [92], SPT-3G [93], Simons Observatory [94] and CMB-S4 [95], we remain with the possibility of deviations from FLRW in the late Universe. Obviously, this deviation at late times should be considered with respect to a perturbed FLRW model. In contrast, when r_d is fixed, the resulting H_0 is discrepant at 2.5σ with SH0ES [6] and in excess of 3σ with local H_0 combinations [90, 91]. Late Universe modifications of the DE sector are firmly ruled out.

Throughout it should be kept in mind that FLRW is a working assumption. Given the CMB is more or less isotropic, FLRW is well motivated in the early Universe⁹. The 4.9σ -significant result of [98], which employs almost 1.5 million quasars primarily at $z > 1$ implies that the CMB restframe does not coincide with the restframe of distant quasars, thus directly undermining the FLRW Universe. Similar results were noted earlier, at lower significance for radio galaxies at lower redshifts and for smaller samples [99–106]. Intriguingly, as we have explained in section 6, strong lensing time delay may provide a new way to probe FLRW breakdown.

Moreover, deviations from FLRW are expected at smaller scales, where they may impact supernovae [107–113]. It is plausible that supernovae below $z \sim 0.15$ are affected by larger than expected bulk flows [108, 113–119]. Interestingly, [118, 119] study scaling relations in

⁹ Note that while there exist anomalies (see [96, 97] for reviews) we cut out low ℓ multipoles, and hence the corresponding anisotropies, in our analysis.

Galaxy clusters for redshifts $z \lesssim 0.3$ and find in line with figure 4 that H_0 varies across the sky within the flat Λ CDM model. However, the lensed quasars are much deeper in redshift, but ultimately may be pointing to the same physics. It is also worth noting that SH0ES remove supernovae below $z \sim 0.025$, precisely because of a documented higher H_0 [120]. However, recent studies show that the higher redshift supernovae are also affected by the local bulk flow as convergence to the CMB frame at such low redshifts cannot be assumed. The effect of the bulk flow on the Hubble parameter has already been the subject of a few studies (see e.g. [121]).

Overall, there are enough claims in the literature that $H_0 \sim 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is a compelling outcome of local H_0 measurements and our observation is that this central value may be offset from what any FLRW cosmology can accommodate. Different recent studies have already put the validity of FLRW in doubt (see e.g. [98] and references therein). Here we have argued that Hubble tension could indeed be yet another manifestation of the breakdown of the FLRW model. Alternatively, if local H_0 values converge lower, but still above $H_0 \sim 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, then we may be looking at new early Universe physics within FLRW.

Acknowledgments

We thank Silvia Galli, Raphael Gavazzi, Subir Sarkar, Leandros Perivolaropoulos, Paul Steinhardt & Kenneth Wong for correspondence and discussion. EOC is funded by the National Research Foundation of Korea (NRF-2020R1A2C1102899). MMSHJ would like to acknowledge SarAmadan Grant ISEF/M/400122. LY is supported by the CQeST of Sogang University (NRF-2020R1A6A1A03047877).

Data availability statement

No new data were created or analysed in this study.

Appendix A. Prior on ω_m

We can get a prior on $\omega_m \equiv \Omega_{m0}h^2$ directly from [30], where it is argued that one can remove the lower multipoles from the CMB to get a determination that is ‘as model independent as possible’ of the late-time cosmology. Excluding multipoles $\ell < 40$, while assuming that ω_b and ω_c are independent in the baryonic and cold dark matter (CDM) sectors, which will only lead to an overestimation, i.e. a conservative error, one arrives at the prior:

$$\omega_m = 0.145 \pm 0.007. \quad (\text{A.1})$$

This can be compared with the Planck value [2] $\omega_m = 0.143 \pm 0.0011$ [2]. Note that the error has increased by removing the lower multipoles.

Although the data analysed in [30] is older (WMAP5 + ACBAR), one can recast the results of [33] to get a more recent prior that is valid in the Planck era. From table 2 of [33], we have

$$\begin{aligned} \Omega_b^{\text{rec}} &= 0.1187 \pm 0.0034, \\ \Omega_c^{\text{rec}} &= 0.6378 \pm 0.071, \end{aligned} \quad (\text{A.2})$$

where it should be noted that the baryonic and CDM energy densities, Ω_b and Ω_c respectively, are evaluated at a recombination redshift of $z_{\text{rec}} = 1089.0 \pm 0.5$. Together with the radiation

sector, they sum to unity $\Omega_b + \Omega_c + \Omega_r = 1$, since the DE sector is negligible at higher redshifts. It is worth noting from table 2 of [33] that different models are considered and the energy densities for different models vary in a negligible way. This appears to back up the claim that the determinations are agnostic about the late-time cosmology. In contrast to the earlier paper [30], only the $\ell < 30$ multipoles are removed.

Once again adding the errors in quadrature, on the assumption that they are independent, one arrives at the following equality,

$$\frac{\omega_m(1+z_{\text{rec}})^3}{\omega_m(1+z_{\text{rec}})^3 + \omega_r(1+z_{\text{rec}})^4} = 0.7565 \pm 0.0078,$$

once one neglects the DE sector. Here we use $\omega_r = 2.47 \times 10^{-5}(1 + 0.2271N_{\text{eff}})$ with $N_{\text{eff}} = 3.046$. We further assume that errors in z_{rec} are independent and solve for ω_m for a large number of configurations $\sim 10\,000$ subject to the assumption that the errors are Gaussian. This ultimately leads to the conservative prior:

$$\omega_m = 0.141 \pm 0.006, \quad (\text{A.3})$$

which is marginally tighter than (A.1).

ORCID iDs

C Krishnan <https://orcid.org/0000-0002-6760-4883>

E Ó Colgáin <https://orcid.org/0000-0003-0813-5156>

References

- [1] Di Valentino E, Mena O, Pan S, Visinelli L, Yang W, Melchiorri A, Mota D F, Riess A G and Silk J 2021 arXiv:2103.01183 [astro-ph.CO]
- [2] Aghanim N *et al* (Planck) 2020 *Astron. Astrophys.* **641** A6
- [3] Freedman W L, Madore B F, Scowcroft V, Burns C, Monson A, Persson S E, Seibert M and Rigby J 2012 *Astrophys. J.* **758** 24
- [4] Riess A G *et al* 2016 *Astrophys. J.* **826** 56
- [5] Riess A G, Casertano S, Yuan W, Macri L M and Scolnic D 2019 *Astrophys. J.* **876** 85
- [6] Riess A G, Casertano S, Yuan W, Bowers J B, Macri L, Zinn J C and Scolnic D 2021 *Astrophys. J. Lett.* **908** L6
- [7] Jang I S and Lee M G 2017 *Astrophys. J.* **836** 74
- [8] Freedman W L, Madore B F, Hatt D, Hoyt T J, Jang I S, Beaton R L, Burns C R, Lee M G, Monson A J, Neeley J R *et al* 2019 arXiv:1907.05922 [astro-ph.CO]
- [9] Freedman W L, Madore B F, Hoyt T, Jang I S, Beaton R, Lee M G, Monson A, Neeley J and Rich J 2020 arXiv:2002.01550 [astro-ph.GA]
- [10] Soltis J, Casertano S and Riess A G 2021 *Astrophys. J. Lett.* **908** L5
- [11] Riess A G *et al* (Supernova Search Team) 1998 *Astron. J.* **116** 1009–38
- [12] Perlmutter S *et al* (Supernova Cosmology Project) 1999 *Astrophys. J.* **517** 565–86
- [13] Eisenstein D J *et al* (SDSS) 2005 *Astrophys. J.* **50** 560–74
- [14] Vagnozzi S, Dhawan S, Gerbino M, Freese K, Goobar A and Mena O 2018 *Phys. Rev. D* **98** 083501
- [15] Ó Colgáin E and Yavartanoo H 2019 *Phys. Lett. B* **797** 134907
- [16] Banerjee A, Cai H, Heisenberg L, Ó Colgáin E, Sheikh-Jabbari M M and Yang T 2021 *Phys. Rev. D* **103** L081305
- [17] Bernal J L, Verde L, Jimenez R, Kamionkowski M, Valcin D and Wandelt B D 2021 arXiv:2102.05066 [astro-ph.CO]
- [18] Wetterich C 1988 *Nucl. Phys. B* **302** 668–96

- [19] Ratra B and Peebles P J E 1988 *Phys. Rev. D* **37** 3406
- [20] Barros B J, Amendola L, Barreiro T and Nunes N J 2019 *J. Cosmol. Astropart. Phys.* **JCAP01(2019)007**
- [21] Gómez-Valent A, Pettorino V and Amendola L 2020 *Phys. Rev. D* **101** 123513
- [22] Peracaula J S, Gómez-Valent A, de Cruz Pérez J and Moreno-Pulido C 2020 *Class. Quantum Grav.* **37** 245003
- [23] Krishnan C, Ó Colgáin E, Sheikh-Jabbari M M and Yang T 2021 *Phys. Rev. D* **103** 103509
- [24] Wong K C et al 2020 *Mon. Not. R. Astron. Soc.* **498** 1420–39
- [25] Millon M et al 2020 *Astron. Astrophys.* **639** A101
- [26] Yang T, Banerjee A and Ó Colgáin E 2020 *Phys. Rev. D* **102** 123532
- [27] Krishnan C, Ó Colgáin E, Ruchika, Sen A A, Sheikh-Jabbari M M and Yang T 2020 *Phys. Rev. D* **102** 103525
- [28] Dainotti M G, De Simone B, Schiavone T, Montani G, Rinaldi E and Lambiase G 2021 arXiv:2103.02117 [astro-ph.CO]
- [29] Valcin D, Bernal J L, Jimenez R, Verde L and Wandelt B D 2020 *J. Cosmol. Astropart. Phys.* **JCAP12(2020)002**
- [30] Vonlanthen M, Räsänen S and Durrer R 2010 *J. Cosmol. Astropart. Phys.* **JCAP08(2010)023**
- [31] Audren B, Lesgourgues J, Benabed K and Prunet S 2013 *J. Cosmol. Astropart. Phys.* **JCAP02(2013)001**
- [32] Audren B 2014 *Mon. Not. R. Astron. Soc.* **444** 827–32
- [33] Verde L, Bellini E, Pigozzo C, Heavens A F and Jimenez R 2017 *J. Cosmol. Astropart. Phys.* **JCAP04(2017)023**
- [34] Efstathiou G 2021 arXiv:2103.08723 [astro-ph.CO]
- [35] Camarena D and Marra V 2021 arXiv:2101.08641 [astro-ph.CO]
- [36] Pesce D W et al 2020 *Astrophys. J. Lett.* **891** L1
- [37] Blakeslee J P, Jensen J B, Ma C-P, Milne P A and Greene J E 2021 *Astrophys. J.* **911** 65
- [38] Kourkchi E, Tully R B, Anand G S, Courtois H M, Dupuy A, Neill J D, Rizzi L and Seibert M 2020 *Astrophys. J.* **896** 3
- [39] Poulin V, Smith T L, Karwal T and Kamionkowski M 2019 *Phys. Rev. Lett.* **122** 221301
- [40] Kreisch C D, Cyr-Racine F Y and Dorè O 2019 arXiv:1902.00534 [astro-ph.CO]
- [41] Agrawal P, Cyr-Racine F Y, Pinner D and Randall L 2019 arXiv:1904.01016 [astro-ph.CO]
- [42] Niedermann F and Sloth M S 2019 arXiv:1910.10739 [astro-ph.CO]
- [43] Niedermann F and Sloth M S 2020 *Phys. Rev. D* **102** 063527
- [44] Bernal J L, Verde L and Riess A G 2016 *J. Cosmol. Astropart. Phys.* **JCAP10(2016)019**
- [45] Lemos P, Lee E, Efstathiou G and Gratton S 2019 *Mon. Not. R. Astron. Soc.* **483** 4803–10
- [46] Wetterich C 2004 *Phys. Lett. B* **594** 17–22
- [47] Doran M and Robbers G 2006 *J. Cosmol. Astropart. Phys.* **JCAP06(2006)026**
- [48] Bean R, Hansen S H and Melchiorri A 2001 *Phys. Rev. D* **64** 103508
- [49] Pettorino V, Amendola L and Wetterich C 2013 *Phys. Rev. D* **87** 083009
- [50] Ade P A R et al (Planck) 2016 *Astron. Astrophys.* **594** A14
- [51] Chevallier M and Polarski D 2001 *Int. J. Mod. Phys. D* **10** 213
- [52] Linder E V 2003 *Phys. Rev. Lett.* **90** 091301
- [53] Lewis A 2019 arXiv:1910.13970 [astro-ph.IM]
- [54] Ó Colgáin E, Sheikh-Jabbari M M and Yin L 2021 arXiv:2104.01930 [astro-ph.CO]
- [55] Cattoën C and Visser M 2007 *Class. Quantum Grav.* **24** 5985
- [56] Ó Colgáin E and Sheikh-Jabbari M M 2021 arXiv:2101.08565 [astro-ph.CO]
- [57] Gavela M B, Hernández D, Honorez L L, Mena O and Rigolin S 2009 *J. Cosmol. Astropart. Phys.* **JCAP07(2009)034**
Gavela M B, Hernández D, Honorez L L, Mena O and Rigolin S 2010 *J. Cosmol. Astropart. Phys.* **JCAP05(2010)e01**
- [58] Vagnozzi S 2020 *Phys. Rev. D* **102** 023518
- [59] Li X and Shafieloo A 2019 *Astrophys. J. Lett.* **883** L3
- [60] Felice A D, Geng C-Q, Pookkillath M C and Yin L 2020 *J. Cosmol. Astropart. Phys.* **JCAP08(2020)038**
- [61] Heisenberg L and Villarrubia-Rojo H 2021 *J. Cosmol. Astropart. Phys.* **JCAP03(2021)032**
- [62] Scolnic D et al 2018 *Astrophys. J.* **859** 101
- [63] Beutler F, Blake C, Colless M, Jones D H, Staveley-Smith L, Campbell L, Parker Q, Saunders W and Watson F 2011 *Mon. Not. R. Astron. Soc.* **416** 3017–32

- [64] Ross A J, Samushia L, Howlett C, Percival W J, Burden A and Manera M 2015 *Mon. Not. R. Astron. Soc.* **449** 835–47
- [65] Alam S et al (BOSS) 2017 *Mon. Not. R. Astron. Soc.* **470** 2617–52
- [66] Abbott T M C et al (DES) 2019 *Mon. Not. R. Astron. Soc.* **483** 4866–83
- [67] Bautista J E et al 2020 *Mon. Not. R. Astron. Soc.* **500** 736–62
- [68] Gil-Marín H et al 2020 *Mon. Not. R. Astron. Soc.* **498** 2492–531
- [69] Raichoor A et al 2020 *Mon. Not. R. Astron. Soc.* **500** 3254–74
- [70] de Mattia A et al 2020 arXiv:2007.09008 [astro-ph.CO]
- [71] Hou J et al 2020 *Mon. Not. R. Astron. Soc.* **500** 1201–21
- [72] Neveux R et al 2020 *Mon. Not. R. Astron. Soc.* **499** 210–29
- [73] du Mas des Bourboux H et al 2020 *Astrophys. J.* **901** 153
- [74] Moresco M et al 2016 *J. Cosmol. Astropart. Phys.* JCAP05(2016)014
- [75] Foreman-Mackey D, Hogg D W, Lang D and Goodman J 2013 *Publ. Astron. Soc. Pac.* **125** 306–12
- [76] Shajib A J et al (DES) 2020 *Mon. Not. R. Astron. Soc.* **494** 6072–102
- [77] Haslbauer M, Banik I and Kroupa P 2020 *Mon. Not. R. Astron. Soc.* **499** 2845–83
- [78] Suyu S H, Marshall P J, Auger M W, Hilbert S, Blandford R D, Koopmans L V E, Fassnacht C D and Treu T 2010 *Astrophys. J.* **711** 201
- [79] Birrer S et al 2020 *Astron. Astrophys.* **643** A165
- [80] Jedamzik K and Pogosian L 2020 *Phys. Rev. Lett.* **125** 181302
- [81] Hill J C, McDonough E, Toomey M W and Alexander S 2020 *Phys. Rev. D* **102** 043507
- [82] Ivanov M M, McDonough E, Hill J C, Simonović M, Toomey M W, Alexander S and Zaldarriaga M 2020 arXiv:2006.11235 [astro-ph.CO]
- [83] D’Amico G, Senatore L, Zhang P and Zheng H 2020 arXiv:2006.12420 [astro-ph.CO]
- [84] Niedermann F and Sloth M S 2020 arXiv:2009.00006 [astro-ph.CO]
- [85] Murgia R, Abellán G F and Poulin V 2020 arXiv:2009.10733 [astro-ph.CO]
- [86] Smith T L, Poulin V, Bernal J L, Boddy K K, Kamionkowski M and Murgia R 2020 arXiv:2009.10740 [astro-ph.CO]
- [87] Jedamzik K, Pogosian L and Zhao G B 2020 arXiv:2010.04158 [astro-ph.CO]
- [88] Lin W, Chen X and Mack K J 2021 arXiv:2102.05701 [astro-ph.CO]
- [89] Yang W, Di Valentino E, Pan S, Wu Y and Lu J 2021 *Mon. Not. R. Astron. Soc.* **501** 5845–58
- [90] Di Valentino E 2021 *Mon. Not. R. Astron. Soc.* **502** 2065–73
- [91] Verde L, Treu T and Riess A G 2019 *Nat. Astron.* **3** 891
- [92] Aiola S et al (ACT) 2020 *J. Cosmol. Astropart. Phys.* JCAP12(2020)047
- [93] Benson B Assa et al (SPT-3G) 2014 *Proc. SPIE Int. Soc. Opt. Eng.* **9153** 91531P
- [94] Ade P et al (Simons Observatory) 2019 *J. Cosmol. Astropart. Phys.* JCAP02(2019)056
- [95] Abazajian K et al 2019 arXiv:1907.04473 [astro-ph.IM]
- [96] Schwarz D J, Copi C J, Huterer D and Starkman G D 2016 *Class. Quantum Grav.* **33** 184001
- [97] Perivolaropoulos L and Skara F 2021 arXiv:2105.05208 [astro-ph.CO]
- [98] Secrest N J, von Hausegger S, Rameez M, Mohayaee R, Sarkar S and Colin J 2021 *Astrophys. J. Lett.* **908** L51
- [99] Blake C and Wall J 2002 *Nature* **416** 150–2
- [100] Singal A K 2011 *Astrophys. J. Lett.* **742** L23
- [101] Gibelyou C and Huterer D 2012 *Mon. Not. R. Astron. Soc.* **427** 1994–2021
- [102] Rubart M and Schwarz D J 2013 *Astron. Astrophys.* **555** A117
- [103] Tiwari P and Nusser A 2016 *J. Cosmol. Astropart. Phys.* JCAP03(2016)062
- [104] Colin J, Mohayaee R, Rameez M and Sarkar S 2017 *Mon. Not. R. Astron. Soc.* **471** 1045–55
- [105] Bengaly C A P, Maartens R and Santos M G 2018 *J. Cosmol. Astropart. Phys.* JCAP04(2018)031
- [106] Siewert T M, Schmidt-Rubart M and Schwarz D J 2020 arXiv:2010.08366 [astro-ph.CO]
- [107] Cooke R and Lynden-Bell D 2010 *Mon. Not. R. Astron. Soc.* **401** 1409–14
- [108] Colin J, Mohayaee R, Sarkar S and Shafieloo A 2011 *Mon. Not. R. Astron. Soc.* **414** 264–71
- [109] Dai D-C, Kinney W H and Stojkovic D 2011 *J. Cosmol. Astropart. Phys.* JCAP04(2011)015
- [110] Turnbull S J, Hudson M J, Feldman H A, Hicken M, Kirshner R P and Watkins R 2012 *Mon. Not. R. Astron. Soc.* **420** 447–54
- [111] Appleby S, Shafieloo A and Johnson A 2015 *Astrophys. J.* **801** 76
- [112] Colin J, Mohayaee R, Rameez M and Sarkar S 2019 *Astron. Astrophys.* **631** L13
- [113] Mohayaee R, Rameez M and Sarkar S 2020 arXiv:2003.10420 [astro-ph.CO]
- [114] Watkins R, Feldman H A and Hudson M J 2009 *Mon. Not. R. Astron. Soc.* **392** 743–56
- [115] Kashlinsky A, Atrio-Barandela F, Kocevski D and Ebeling H 2009 *Astrophys. J. Lett.* **686** L49–52

- [116] Feldman H A, Watkins R and Hudson M J 2010 *Mon. Not. R. Astron. Soc.* **407** 2328–38
- [117] Rameez M and Sarkar S 2019 arXiv:[1911.06456](#) [[astro-ph.CO](#)]
- [118] Migkas K, Schellenberger G, Reiprich T H, Pacaud F, Ramos-Ceja M E and Lovisari L 2020 *Astron. Astrophys.* **636** A15
- [119] Migkas K, Pacaud F, Schellenberger G, Erler J, Nguyen-Dang N T, Reiprich T H, Ramos-Ceja M E and Lovisari L 2021 arXiv:[2103.13904](#) [[astro-ph.CO](#)]
- [120] Jha S, Riess A G and Kirshner R P 2007 *Astrophys. J.* **659** 122–48
- [121] Heß S and Kitaura F-S 2016 *Mon. Not. R. Astron. Soc.* **456** 4247–55