



Hermitian–Yang–Mills Connections on Collapsing Elliptically Fibered $K3$ Surfaces

Ved Datar¹ · Adam Jacob²

Received: 30 October 2017 / Accepted: 28 June 2021 / Published online: 7 January 2022
© The Author(s) 2022

Abstract

Let $X \rightarrow \mathbb{P}^1$ be an elliptically fibered $K3$ surface, admitting a sequence ω_i of Ricci-flat metrics collapsing the fibers. Let V be a holomorphic $SU(n)$ bundle over X , stable with respect to ω_i . Given the corresponding sequence Ξ_i of Hermitian–Yang–Mills connections on V , we prove that, if E is a generic fiber, the restricted sequence $\Xi_i|_E$ converges to a flat connection A_0 . Furthermore, if the restriction $V|_E$ is of the form $\bigoplus_{j=1}^n \mathcal{O}_E(q_j - 0)$ for n distinct points $q_j \in E$, then these points uniquely determine A_0 .

Keywords Hermitian–Yang–Mills · Holomorphic degenerations · Elliptic fibrations · $K3$ surfaces

Mathematics Subject Classification 53B35 · 53C07 · 53C26 · 53C55

1 Introduction

In this paper, we study degenerations of Hermitian–Yang–Mills connections on a $K3$ surface. We are motivated by the work of Gross–Wilson [1], and later Gross–Tosatti–Zhang [2,3], who study Ricci-flat metrics on elliptically fibered Calabi–Yau’s as the volume of the fibers tends to zero (see also [4–7]). These types of degenerations relate to the conjectural picture of mirror symmetry put forth by Strominger–Yau–Zaslow [8], who postulate that mirror Calabi–Yau manifolds are given by dual torus fibrations over a real base with a singular affine structure. One major challenge when confronting this conjecture is the difficulty associated with constructing Lagrangian torus fibrations on

✉ Adam Jacob
ajacob@math.ucdavis.edu

Ved Datar
vvdatar@iisc.ac.in

¹ Department of Mathematics, Indian Institute of Science, Bengaluru, India

² Department of Mathematics, UC Davis, One Shields Ave, Davis, CA, USA

a given Calabi–Yau. However, if one instead considers degenerations of Calabi–Yau’s, the fibration structure often becomes apparent in the limit.

Vafa’s extension of the mirror symmetry conjecture to include holomorphic bundles raises the question of how Yang–Mills connections behave under these degenerations [9]. For a general abelian fibered Calabi–Yau, Fukaya makes the following conjecture: Given a sequence of Yang–Mills connections on a family of Calabi–Yau metrics with collapsing fibers, there exists a rectifiable set in the base of real codimension at least 2, such that on all fibers away from this set, the connections have bounded curvature, and the restriction to each torus fiber converges to a flat connection [10, Conjecture 5.5]. In the context of SYZ mirror symmetry, a fiberwise flat connection will define a Lagrangian submanifold \mathcal{L} in the mirror Calabi–Yau, and Fukaya further conjectures there exists a corresponding mirror sequence of Lagrangians converging to \mathcal{L} . In this paper we partially address the vector bundle portion of Fukaya’s conjecture, in the case of a fixed holomorphic $SU(n)$ bundle over a $K3$ surface.

Our setup is as follows. Let $\pi : X \rightarrow \mathbb{P}^1$ be an elliptic $K3$ surface. Let $\omega_{\mathbb{P}^1}$ be a Kähler form on \mathbb{P}^1 , and ω_X a Kähler form on X . Consider the family $\pi^*\omega_{\mathbb{P}^1} + t^2\omega_X$, which approaches the boundary of the Kähler cone as $t \rightarrow 0$, and let ω_t be the unique Ricci-flat metric in the class $[\pi^*\omega_{\mathbb{P}^1} + t^2\omega_X]$ given by Yau’s theorem [11]. Next, let $(V, \bar{\partial}_{\Xi})$ be a holomorphic $SU(n)$ bundle over X , with a fixed metric H_0 . Assume there exists a sequence $t_i \rightarrow 0$ such that the bundle V is stable with respect to ω_{t_i} . By the theorem of Donaldson, Uhlenback–Yau [12,13], there exists a corresponding sequence of connections Ξ_i solving the Hermitian–Yang–Mills (HYM) equations:

$$F_{\Xi_i} \wedge \omega_{t_i} = 0 \quad \text{and} \quad (F_{\Xi_i})^{0,2} = 0.$$

Furthermore, each Ξ_i is complex gauge equivalent to Ξ , so they define the same holomorphic structure (see (3.1)). We now state our main result:

Theorem 1.1 *With the set-up as above:*

- (1) *There exists a finite subset $Z \subset \mathbb{P}^1$, such that for any $x \in \mathbb{P}^1 \setminus Z$, if $E = \pi^{-1}(x)$, then the restriction $\Xi_i|_E$ converges smoothly, along a subsequence and modulo unitary gauge transformations, to a flat connection on the fiber.*
- (2) *Furthermore, if the restriction $V|_E$ is isomorphic to a direct sum of line bundles $\bigoplus_{j=1}^n \mathcal{O}_E(q_j - 0)$ for n distinct points $q_j \in E$, then the limiting flat connection is uniquely determined, and given by*

$$A_0 = \frac{\pi}{\text{Im}(\tau)} (\text{diag}\{\bar{q}_1, \dots, \bar{q}_n\}dz - \text{diag}\{q_1, \dots, q_n\}d\bar{z}), \tag{1.1}$$

where z is the holomorphic coordinate on the fiber, and τ determines the complex structure. In this case, we also have the following convergence:

$$\|\Xi_i|_E - A_0\|_{L^2_1(E, H_0, g_0, A_0)} \rightarrow 0.$$

Here g_0 is a flat reference metric on E , and the flat connection A_0 is used to compute derivatives.

Note that in the second point above, no gauge transformations are needed for convergence. The first point follows from a bubbling argument. Our sequence of connections has bounded Yang–Mills energy, thus there can only be a finite number of bubbles, and we show away from these bubbles the curvature of $\Xi_i|_E$ must approach zero in the C^0 norm. This step closely follows two cases from Dostoglou–Salamon, from their proof of the Atiyah–Floer conjecture [14,15]. The key difference here is that our Ricci flat metric ω_{t_i} is not a product metric, so we rely on certain convergence results for ω_{t_i} .

It then follows that $\Xi_i|_E$ must approach some limiting flat connection, and the main contribution of this paper is the explicit identification of the limit, under the assumption that the restriction of the holomorphic bundle $V|_E$ is isomorphic to a direct sum of line bundles $\bigoplus_{j=1}^n \mathcal{O}_E(q_j - 0)$. We use the observation that, because our sequence of connections Ξ_i all define a fixed holomorphic structure, there exists a sequence of Hermitian endomorphisms s_i satisfying $\Xi_i|_E = e^{s_i}(A_0)$ (where this action is defined by (3.1)). Although there is no hope of achieving C^0 control of s_i , we prove a gauge fixing result, and demonstrate that there exists a suitable normalization s'_i , defining the same connection, which in addition satisfies a uniform C^0 bound. This significant step is detailed in Theorem 5.1, which in particular hinges on a Poincaré inequality (5.1), where the explicit form of $V|_E$ is used. From here, convergence of $\Xi_i|_E$ to A_0 stated in Theorem 1.1 follows from standard theory.

Next, we turn to a specific geometric setup where Theorem 1.1 applies. Although this setup requires more assumptions, it has the benefit of producing explicit examples of bundles where the the limiting flat connection can be identified on a generic fiber. We now assume $\pi : X \rightarrow \mathbb{P}^1$ is a projective, elliptic K3 surface with a section σ and singular fibers of type I_1 or II . Assume the restriction of V to a generic fiber is semi-stable and regular (see Sect. 3 for relevant definitions). Then by the work of Friedman–Morgan–Witten in [16], there exists a divisor $D_V \in |n\sigma(\mathbb{P}^1) + kl|$, called the *spectral cover* associated to V , where l denotes the effective divisor class of the fibers, and $k \in \mathbb{Z}$ satisfies $0 \leq k \leq c_2(V)$. If D_V is reduced and irreducible, then V is stable with respect to $\pi^*[\omega_{\mathbb{P}^1}] + t^2[\omega_X]$ for any ample class $[\omega_X]$ on X , for $0 < t^2 \leq (\frac{n^3}{4}c_2(V))^{-1}$. Thus, for any sequence $t_i \rightarrow 0$ we can always find a corresponding sequence of HYM connections Ξ_i on V . More importantly, the intersection of the spectral cover D_V with a generic fiber E precisely picks out the points q_1, \dots, q_n from (1.1), and so the limiting flat connection is uniquely determined away from the ramification points of D_V .

Corollary 1.2 *Assume $\pi : X \rightarrow \mathbb{P}^1$ is a projective, elliptic K3 surface with a section, with singular fibers of type I_1 or II . Let V be a holomorphic $SU(n)$ bundle where the restriction to a generic fiber is semi-stable and regular, and assume the spectral cover D_V is reduced and irreducible. Then for any sequence $t_i \rightarrow 0$, there exists a sequence of HYM connections Ξ_i on V corresponding to ω_{t_i} . Furthermore, away from a finite number of fibers, there exists a HYM connection Ξ_0 uniquely determined by D_V , satisfying $\Xi_0|_E = A_0$, where A_0 is defined via (1.1). Specifically, on a generic fiber E the points q_1, \dots, q_n defining A_0 are given by*

$$D_V \cap E = q_1 + \dots + q_n.$$

On E we again have the convergence

$$\|\Xi_i|_E - \Xi_0|_E\|_{L^2_1(E, H_0, g_0, A_0)} \rightarrow 0.$$

The above setup is particularly attractive in that it allows for us to specify the limiting flat connection in a family that varies homomorphically in the base. Thus our result is a natural starting place to explore convergence in general, as opposed to only in the fiber direction.

Since adiabatic limits of Yang–Mills connections are fairly well studied, we now put our work in the context of previous results. Working on the product of two compact Riemann surfaces with trivial fibration $\pi : \Sigma_1 \times \Sigma_2 \rightarrow \Sigma_1$, J. Chen considers a family of metrics collapsing the fibers, and analyzes the convergence of a corresponding family of anti-self dual Yang–Mills connections [17, Theorem 4.10]. Assuming that Σ_2 has genus at least two, he proves that, modulo a sequence of gauge transformations and away from a bubbling set, the fiber component of the connection will converge continuously to a flat connection. Following this work, and using Fukaya’s gauge fixing theorem [18, Theorem 1.7], T. Nishinou improves upon Chen’s result, demonstrating smooth convergence away from a finite number of fibers and modulo gauge transformations, where the restriction of the limit to a fiber will be flat [19, Theorem 1.2]. This result requires the moduli space of flat connections over Σ_2 to be smooth and of expected dimension, with no reducible flat connections. The failure of such an assumption to hold over an elliptic curve is a major obstacle to extending the above results to elliptic fibrations.

In the case of $SU(2)$ bundles over the product of elliptic curves, in [20] Nishinou is able to partially extend his above results, after utilizing his gauge fixing theorem from [21, Theorem 3.11]. In some ways, our Theorem 5.1 can be thought of as a generalization to higher rank of this gauge fixing theorem, although we have already assumed existence of a fixed holomorphic structure. In fact, this assumption serves as a major simplification throughout our paper, compared to the general case of a sequence of anti-self dual Yang–Mills connections considered in [14, 17–20]. The most notable simplification is that, because our sequence of connections Ξ_i are all complex gauge equivalent to Ξ , we can bypass working with a sequence of holomorphic maps (which plays a role in [17, 19, 20]), as well as the more difficult type three bubbles of Dostoglou–Salamon [14, 15]. Additionally, this assumption allows us to prove the convergence in the second point of Theorem 1.1 directly, without relying on unitary gauge transformations.

Although our main result only applies to the restriction of Ξ_i to each fiber, one may hope to demonstrate convergence on any compact set away from a finite number of fibers. Nishinou achieves this in [19] and [20], as his assumptions allow a Poincaré type inequality in a neighborhood of a fiber, even in the elliptic curve case (this follows from the estimate in Lemma 6.43 from [18]). This estimate implies that once C^0 control of the complexified gauge transformation is demonstrated on one fiber, it holds for nearby fibers. Unfortunately we are unable to extend Lemma 6.43 from [18] to our setting, as our Poincaré inequality (Proposition 5.2) requires a normalization that only holds fiberwise. Another result in this direction is proven by Fu in [22], who considers a specific rank two bundle over the product of two elliptic curves

which is given by a two sheeted spectral cover. He defines a reference metric which satisfies desired asymptotic behavior near the ramification points of the cover, and then demonstrates that the a sequence of HYM metrics will converge smoothly to this reference metric. Because in our setting we consider a spectral cover as well, one may hope to extend Fu's result to the K3 surface case. Here, one major difficulty is the problem of constructing a reference metric near the singularities of the fibration. It is possible that the asymptotics of the metrics constructed in [23–25] may provide a clue, and we hope to investigate these types of constructions in future work.

Finally, we remark that after an earlier draft of this paper appeared, building on this work, the authors, along with Y. Zhang, were able to demonstrate convergence of Ξ_i on compact sets away from a finite number of fibers, under the assumptions of Corollary 1.2. We direct the reader to [26] for details.

Our paper is organized as follows. In Sect. 2 we describe in detail semi-flat Kähler metrics on a K3 surface, which serve as a local model for our degenerating Ricci-flat metrics away from the singular fibers. Next in Sect. 3 we introduce the necessary background on holomorphic vector bundles over elliptic fibrations, and state some preliminary results. Our bubbling argument is described Sect. 4. We then turn to identifying the limiting flat connection, and prove our gauge fixing result is Sect. 5. In Sect. 6 we complete the proof of our main theorem, and demonstrate convergence of our connections.

2 Semi-Flat Kähler Metrics

In this section, we review the construction of semi-flat Kähler metrics on a K3 surface. These metrics will not only describe the limiting behavior of the Ricci-flat metrics in dilated coordinates, and thus play a role in our bubbling argument, but they will also be useful for our understanding of the holomorphic structure of V . To begin, we introduce the notion of a special Kähler metric, which lives on the base of our elliptic fibration, and are a useful starting place to defining the semi-flat metric. We will closely follow the paper of Freed [27].

Let B be a Riemannian manifold of real dimension two. Assume TB admits a flat, torsion free connection ∇^B , which gives a covering of B by local affine coordinate charts. Furthermore, assume the coordinate transformations lie in $SL(2, \mathbb{R})$. Let (x^1, x^2) be coordinates in a local chart, and let $\phi_{ij} dx^i dx^j$ be a Hessian metric solving the real Monge-Ampere equation

$$\det(\phi_{ij}) = 1. \quad (2.1)$$

Here, we use the notation $\phi_{ij} := \frac{\partial^2 \phi}{\partial x^i \partial x^j}$ for a smooth function ϕ on B . We also denote $\frac{\partial \phi}{\partial x^i}$ by ϕ_i .

Consider the locally defined 2-form $\omega_B = dx^1 \wedge dx^2$. Because $SL(2, \mathbb{R}) = Sp(2, \mathbb{R})$, any matrix $A \in SL(2, \mathbb{R})$ preserves ω_B , and thus ω_B is well defined on all of B . It defines both a natural symplectic form and a volume form. Furthermore, $\nabla^B \omega_B = 0$, and so ∇^B is a symplectic connection. Taken together, ω_B and ϕ_{ij} define

an almost complex structure I on TB , which in coordinates can be expressed by

$$I\left(\frac{\partial}{\partial x^k}\right) = -\phi_{k2}\frac{\partial}{\partial x^1} + \phi_{k1}\frac{\partial}{\partial x^2},$$

for $k = 1, 2$. One can show explicitly that Nijenhuis tensor of I vanishes and thus it is integrable.

Definition 2.1 (B, ω_B, I) is special Kähler if it admits a real, flat, torsion-free, symplectic connection ∇^B satisfying

$$d_{\nabla^B} I = 0.$$

To see that B is special Kähler, note in affine coordinates the flat connection ∇^B is simply given by d , and so

$$\frac{\partial}{\partial x^k} I^p{}_q - \frac{\partial}{\partial x^q} I^p{}_k = 0,$$

which follows because ϕ_{ij} is a Hessian metric.

Given the complex structure I , we can give holomorphic coordinate functions on B .

Lemma 2.2 *The functions*

$$w = x^1 + \mathbf{i}\phi_2 \quad \text{and} \quad \xi = -x^2 + \mathbf{i}\phi_1.$$

are holomorphic with respect to the complex structure I .

Proof Taking the exterior derivative gives $dw = (1 + \mathbf{i}\phi_{12})dx^1 + \mathbf{i}\phi_{22}dx^2$ and $d\xi = \mathbf{i}\phi_{11}dx^1 - (1 - \mathbf{i}\phi_{12})dx^2$. By our explicit representation of I it is easy to check that $I(dw) = \mathbf{i}dw$ and $I(d\xi) = \mathbf{i}d\xi$. \square

For the remainder of the paper we choose w as our holomorphic coordinate on the base. We would like to better understand the holomorphic vector field $\frac{\partial}{\partial w}$. First, note that the coordinate transformation $T(x^1, x^2) \rightarrow (w, \bar{w})$ has pushforward matrix

$$T_* = \begin{pmatrix} 1 + \mathbf{i}\phi_{12} & \mathbf{i}\phi_{22} \\ 1 - \mathbf{i}\phi_{12} & -\mathbf{i}\phi_{22} \end{pmatrix} \quad \text{and} \quad T_*^{-1} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ -\mathbf{i}\frac{\phi_{11}}{1+\mathbf{i}\phi_{12}} & \mathbf{i}\frac{\phi_{11}}{1-\mathbf{i}\phi_{12}} \end{pmatrix},$$

where we have used (2.1). We now compute the partial derivative

$$\begin{aligned} \frac{\partial \xi}{\partial w} &= \frac{\partial \xi}{\partial x^1} \frac{\partial x^1}{\partial w} + \frac{\partial \xi}{\partial x^2} \frac{\partial x^2}{\partial w} \\ &= (\mathbf{i}\phi_{11}) \left(\frac{1}{2}\right) + (-1 + \mathbf{i}\phi_{12}) \left(\frac{-\mathbf{i}}{2} \frac{\phi_{11}}{1 + \mathbf{i}\phi_{12}}\right) \\ &= \frac{\mathbf{i}}{2} \phi_{11} \left(1 + \frac{1 - \mathbf{i}\phi_{12}}{1 + \mathbf{i}\phi_{12}}\right) = \mathbf{i} \frac{\phi_{11}}{1 + \mathbf{i}\phi_{12}} = \mathbf{i} \frac{1 - \mathbf{i}\phi_{12}}{\phi_{22}}, \end{aligned} \tag{2.2}$$

where the last equality follows from (2.1). For simplicity we will use the notation $\tau := \frac{\partial \xi}{\partial w}$, as τ will define the complex structure of our elliptic fibers. Then, by the explicit formula from Freed [27, Equation (1.12)], we see

$$\frac{\partial}{\partial w} = \frac{1}{2} \left(\frac{\partial}{\partial x^1} - \tau \frac{\partial}{\partial x^2} \right).$$

Next we construct a hyper-Kähler structure on TB . Quotienting out TB by a lattice Λ will give a local model for our elliptic fibration X away from the singular fibers. To begin, consider the following extension of our Hessian metric to TB :

$$g = \phi_{ij} \left(dx^i dx^j + dy^i dy^j \right).$$

With this metric we define three complex structures which make up a hyper-Kähler triple.

By a slight abuse of notation, let I denote the complex structure on TB induced from the complex structure I on the base. In particular I can be expressed as

$$I \left(\frac{\partial}{\partial x^k} \right) = -\phi_{k2} \frac{\partial}{\partial x^1} + \phi_{k1} \frac{\partial}{\partial x^2} \quad \text{and} \quad I \left(\frac{\partial}{\partial y^k} \right) = \phi_{k2} \frac{\partial}{\partial y^1} - \phi_{k1} \frac{\partial}{\partial y^2}$$

for $k = 1, 2$. The corresponding Kähler form is given by

$$\omega_I = dx^1 \wedge dx^2 - dy^1 \wedge dy^2. \tag{2.3}$$

In this complex structure the fibers are holomorphic subvarieties.

Next, we consider a complex structures J where the fibers are special Lagrangian, defined by

$$J \left(\frac{\partial}{\partial x^k} \right) = \frac{\partial}{\partial y^k} \quad \text{and} \quad J \left(\frac{\partial}{\partial y^k} \right) = \frac{\partial}{\partial x^k}.$$

Using the metric g the corresponding Kähler form is

$$\omega_J = \phi_{ij} dx^i \wedge dy^j.$$

Finally, one can define the complex structure $K = JI$, which together with g gives the Kähler form

$$\omega_K = dx^1 \wedge dy^2 - dx^2 \wedge dy^1.$$

It is easy to see that ω_I, ω_J and ω_K are closed, and by a lemma of Hitchin [28] it follows that I, J , and K are integrable complex structures. In the standard fashion we

can construct the following top dimensional holomorphic forms:

$$\begin{aligned}\Omega_I &= \omega_K + \mathbf{i}\omega_J \\ \Omega_J &= \omega_I + \mathbf{i}\omega_K \\ \Omega_K &= \omega_I + \mathbf{i}\omega_J,\end{aligned}$$

giving hyper-Kähler triple. Note the metric and complex structures defined above are invariant under translation in the y -coordinates. Thus, if Λ is the standard lattice $\langle 1, 1 \rangle$, the entire setup will descend to the elliptically fibered manifold TB/Λ .

We now construct complex coordinates on TB/Λ . We have a complex coordinate w on the base B , and in the fiber direction we define

$$z = \tau y^1 + y^2. \tag{2.4}$$

Here, τ represents the complex period of the elliptic curve, and is defined above in (2.2). By definition $\tau := \frac{\partial \xi}{\partial w}$, and since ξ is a holomorphic function, τ is holomorphic in w as well. This leads to the following:

Lemma 2.3 *The coordinates (w, z) are holomorphic coordinates on TB/Λ with respect to I .*

Proof Lemma 2.2 shows that w is holomorphic, and so it remains to be seen that $dz(V) = 0$ for all vector fields V of type $(0, 1)$. Taking the exterior derivative of z gives

$$dz = \tau dy^1 + y^1 \frac{\partial \tau}{\partial w} dw + dy^2, \tag{2.5}$$

where we used τ is holomorphic. At first glance, the term $y^1 \frac{\partial \tau}{\partial w} dw$ may seem out of place, however, it is important to remember that unless τ is constant, our local picture is not the cartesian product of the base with an elliptic curve, and so this term is expected.

Consider the vector field

$$\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial y^1} - \frac{\mathbf{i}\phi_{11}}{1 + \mathbf{i}\phi_{12}} \frac{\partial}{\partial y^2} \right).$$

From the explicit form of dz , we see that in order for $\frac{\partial}{\partial \bar{z}}$ to be anti-holomorphic, it needs to be killed by the form dw on the base.

Using the definition of I , and the fact that $\det(\phi_{ij}) = 1$, we compute

$$\begin{aligned}I \left(\frac{\partial}{\partial \bar{z}} \right) &= \frac{1}{2} \left(\phi_{12} - \frac{\mathbf{i}\phi_{11}\phi_{22}}{1 + \mathbf{i}\phi_{12}} \right) \frac{\partial}{\partial y^1} + \frac{1}{2} \left(-\phi_{11} + \frac{\mathbf{i}\phi_{11}\phi_{12}}{1 + \mathbf{i}\phi_{12}} \right) \frac{\partial}{\partial y^2} \\ &= \frac{1}{2} (-\mathbf{i}) \frac{\partial}{\partial y^1} + \frac{1}{2} \left(-\frac{\phi_{11}}{1 + \mathbf{i}\phi_{12}} \right) \frac{\partial}{\partial y^2} = -\mathbf{i} \frac{\partial}{\partial \bar{z}},\end{aligned}$$

which demonstrates this vector field is of type $(0, 1)$. By Lemma 2.2 it follows that $dw \left(\frac{\partial}{\partial \bar{z}} \right) = 0$. Additionally, we have

$$(\tau dy^1 + dy^2) \left(\frac{\partial}{\partial \bar{z}} \right) = \frac{1}{2}(\tau - \tau) = 0.$$

So $dz \left(\frac{\partial}{\partial \bar{z}} \right) = 0$. Since $\frac{\partial}{\partial w}$ and $\frac{\partial}{\partial \bar{z}}$ span all local $(0, 1)$ vector fields, we conclude that (w, z) are holomorphic coordinates. □

We conclude this section with a more detailed discussion of how the Ricci-flat Kähler metrics behave in the limit. We recall our setup from the introduction. Let $\pi : X \rightarrow \mathbb{P}^1$ be an elliptic K3 surface, and denote by Z_π the image of the singular fibers. Let $\omega_{\mathbb{P}^1}$ be the Fubini-Study metric on \mathbb{P}^1 , and ω_X a Kähler form on X . Let ω_t be the unique Ricci-flat metric in the class $[\pi^* \omega_{\mathbb{P}^1} + t^2 \omega_X]$. The convergence we need is local, so we fix a small, simply connected open set $U \subset \mathbb{P}^1$ away from Z_π , and define $X_U := \pi^{-1}(U)$. On U we can consider the Kähler form ω_B along with the Hessian metric ϕ_{ij} , which on this small open set is equivalent to $\omega_{\mathbb{P}^1}$. On X_U we have the fixed background metric ω_X , but we also have the Kähler form ω_I as defined above, called the *semi-flat metric*, and we denote it by $\omega_I =: \omega_{SF}$ for emphasis.

We will need the following uniform equivalence result. By [2, Lemma 4.1], there exists a constant C so that for t small enough

$$C^{-1} \left(\pi^* \omega_{\mathbb{P}^1} + t^2 \omega_X \right) \leq \omega_t \leq C \left(\pi^* \omega_{\mathbb{P}^1} + t^2 \omega_X \right). \tag{2.6}$$

We also need a result that demonstrates how ω_t degenerates. Consider the projection $p : U \times \mathbb{C} \rightarrow (U \times \mathbb{C})/\Lambda =: X_U$, and the coordinate transformation $L_t : U \times \mathbb{C} \rightarrow U \times \mathbb{C}$ defined by

$$L_t(x, y) = \left(x, \frac{y}{t} \right). \tag{2.7}$$

This coordinate transformation is a dilatation, designed so that the size of the fibers of π with respect to the metric $L_t^* p^* \omega_t$ are fixed. We will use the following result of Hein-Tosatti from [4, Proof of Theorem 1.1], which demonstrates how the semi-flat metric serves as a model for the limiting behavior of ω_t .

Proposition 2.4 (Hein-Tosatti [4]) *There exists a constant C so that for t small enough*

$$C^{-1} p^* \left(\pi^* \omega_B + \omega_{SF} \right) \leq L_t^* p^* \omega_t \leq C p^* \left(\pi^* \omega_B + \omega_{SF} \right).$$

This estimate also appears in [7], with the extra assumption that X is projective. For more details on convergence results, we direct the reader to [2,3].

3 Holomorphic Bundles over Elliptic Manifolds

In this section, we provide the necessary background on holomorphic vector bundles, including the relevant notions of stability needed to construct our sequence of HYM connections. We also introduce the construction of a spectral cover associated to V , following Friedman–Morgan–Witten [16] (see also [29–31]), and conclude the section with the construction of Ξ_0 used in Corollary 1.2.

To begin, suppose (X, ω) is a compact Kähler manifold of complex dimension m . Let $(V, \bar{\partial}_\Xi)$ be a holomorphic bundle over X . For any Hermitian metric H_0 on V , there exists a unique connection, called the Chern connection, compatible with both the metric and the holomorphic structure, which we denote by Ξ . The *degree* of V is defined by the following integral:

$$\text{deg}(V, \omega) = \mathbf{i} \int_X \text{Tr}(F_\Xi) \wedge \omega^{m-1}.$$

Given two metrics on V , the curvatures of the two corresponding Chern connections will differ by a $\partial\bar{\partial}$ -exact term, demonstrating that the degree is independent of a choice of metric. Furthermore, the degree does not depend on the representative of the Kähler class $[\omega]$. However, for $m > 1$, changing the class may change the degree.

Definition 3.1 (V, ω) is *stable* if, for all proper, torsion-free subsheaves $\mathcal{F} \subset E$,

$$\frac{\text{deg}(\mathcal{F}, \omega)}{\text{rk}(\mathcal{F})} < \frac{\text{deg}(V, \omega)}{\text{rk}(V)}.$$

(V, ω) is *semi-stable* if the above expression holds with a weak inequality.

Note that if \mathcal{F} is not locally free, its degree is defined by computing the degree of $\det(\mathcal{F})$, which is always a line bundle.

On any complex manifold, the space of one forms decomposes into the eigenspaces for $\pm \mathbf{i}$ with respect to the complex structure. This allows us to write the any connection Ξ as $\Xi = \Xi^{1,0} + \Xi^{0,1}$. Using this decomposition, one can define an action of the complexified gauge group on the space of connections. Specifically, if $\sigma \in GL(V)$, then

$$\sigma(\Xi) = \sigma^{*-1} \Xi^{1,0} \sigma^* + \sigma^{*-1} \partial \sigma^* + \sigma \Xi^{0,1} \sigma^{-1} - \bar{\partial} \sigma \sigma^{-1}, \tag{3.1}$$

Note that if σ is in fact unitary, the above action reduces to the standard action of the unitary gauge group. In this case, we use the standard notation $u^* \Xi$ for the unitary action.

We now turn to the HYM equations on a general Kähler manifold:

$$\mathbf{i} F_\Xi \wedge \omega^{m-1} = \frac{m \text{deg}(V, \omega)}{\text{rk}(V) \text{Vol}(X)} Id_V \omega^m \quad \text{and} \quad (F_\Xi)^{0,2} = 0. \tag{3.2}$$

For any metric g on X and connection Ξ on V , the Yang–Mills energy is defined by the following integral

$$\mathcal{YM}(\Xi, g) := \int_X |F_\Xi|_{H_0, g}^2 dV_g.$$

Critical points of this energy functional are called Yang–Mills connections, and one can check using the Kähler identities that HYM connections are a special class of Yang–Mills connections which are compatible with the complex structure on X .

Note that second equation in (3.2) stipulates that Ξ is compatible with the holomorphic structure on V . By definition this second equation is satisfied by the Chern connection Ξ , and one can check this compatibility is preserved along the action (3.1). This leads to the following question: Given a holomorphic vector bundle with fixed metric H_0 , does there exist a solution $\hat{\Xi}$ to (3.2) in the orbit of (3.1)? A definitive answer to this question was given by Donaldson, Uhlenbeck–Yau, in the following fundamental result.

Theorem 3.2 (Donaldson [12], Uhlenbeck–Yau [13]) *A holomorphic bundle V over (X, ω) admits a unique Hermitian–Yang–Mills connection in the complex gauge orbit of the Chern connection if and only if it is stable.*

In fact, one can prove that if $\hat{\Xi}$ is the unique Hermitian–Yang–Mills connection, it can be expressed as $\hat{\Xi} = e^s(\Xi)$, where s is a trace free Hermitian endomorphism of V .

Given this background, we return to our setup. Let $\pi : X \rightarrow \mathbb{P}^1$ be an elliptically fibered $K3$ surface, and let ω_t be the unique Ricci-flat Kähler metric in the class $[\pi^*\omega_{\mathbb{P}^1} + t^2\omega_X]$. Assume $(V, \bar{\partial}_\Xi)$ is a holomorphic $SU(n)$ bundle over X . This implies the curvature F_Ξ is trace free, and so $\deg(V, \omega_t) = 0$ for all t . Furthermore, assume that $(V, \bar{\partial}_\Xi)$ is stable with respect to ω_{t_i} for some sequence $t_i \rightarrow 0$. Let g_{t_i} be the Kähler metrics associated to ω_{t_i} . By the theorem of Donaldson–Uhlenbeck–Yau there exists a corresponding sequence of HYM connections Ξ_i on V . Note that in our particular setting, the HYM equations take the simpler form

$$F_\Xi \wedge \omega = 0 \quad \text{and} \quad (F_\Xi)^{0,2} = 0.$$

We will need the following Lemma, which states that the Yang–Mills energy of a HYM connection is a topological invariant. This result is standard, and can be found, for instance, in [32]. We include a proof for the reader’s convenience.

Lemma 3.3 *The Yang–Mills energy of Ξ_i with respect to the metric g_{t_i} , is fixed, i.e.*

$$\mathcal{YM}(\Xi_i, g_{t_i}) = \mathcal{YM}(\Xi, g_{t_0}).$$

Proof Let $\mathbf{i}\Lambda_\omega$ denote the adjoint of wedging with the Kähler form ω . Then, the equation $F_\Xi \wedge \omega = 0$ can be equivalently expressed as $\mathbf{i}\Lambda_\omega F_\Xi = 0$. Equality (4.4.5) in [33] shows that for any complex surface X one has

$$\int_X \text{Tr}(F_{\Xi_i} \wedge F_{\Xi_i}) = \|F_{\Xi_i}\|_{L^2(H_0, g_{t_i})}^2 - \|\mathbf{i}\Lambda_{\omega_{t_i}} F_{\Xi_i}\|_{L^2(H_0, g_{t_i})}^2.$$

Since F_{Ξ_i} is HYM with respect to g_{t_i} , the right most term vanishes, and so

$$\mathcal{YM}(\Xi_i, g_{t_i}) = \int_X \text{Tr}(F_{\Xi_i} \wedge F_{\Xi_i}).$$

The right-hand side above yields a topological invariant $[c_2(V) - \frac{1}{2}c_1^2(V)] \cup X$, and is thus independent of i , proving the lemma. \square

We now review the Friedman–Morgan–Witten construction of stable holomorphic bundles on elliptic fibrations with sections, and is needed for Corollary 1.2. We begin by looking at a single fiber. Let E be an elliptic curve, and $0 \in E$ the identity of the group law. Denote the trivial line bundle by \mathcal{O} , and given a point $q \in E$, let $\mathcal{O}_E(q - 0)$ be the line bundle associated to the divisor $q - 0$. We also define a sequence of rank r bundles (denoted \mathcal{I}_r) inductively, with $\mathcal{I}_1 = \mathcal{O}$ and \mathcal{I}_r the unique nontrivial extension of \mathcal{I}_{r-1} by \mathcal{O} . Recall the following theorem of Atiyah (Theorem 5 from [34]):

Theorem 3.4 (Atiyah [34]) *Any semi-stable, degree zero bundle V over E is isomorphic to a direct sum of bundles of the form $\mathcal{O}_E(q - 0) \otimes \mathcal{I}_r$, i.e.*

$$V \cong \bigoplus_{j=1}^{\ell} \mathcal{O}_E(q_j - 0) \otimes \mathcal{I}_{r_j}.$$

Definition 3.5 A semi-stable bundle is called regular if in the above direct sum $q_i \neq q_j$ for $i \neq j$.

Note that bundles of the form $\mathcal{O}_E(q_j - 0) \otimes \mathcal{I}_{r_j}$ do not admit flat connections unless $r_j = 1$. However, we can instead replace $\mathcal{O}_E(q_j - 0) \otimes \mathcal{I}_{r_j}$ with its Seshadri filtration $\mathcal{O}_E(q_j - 0)^{\oplus r_j}$, and define all bundles with the same Seshadri filtration to be \mathcal{S} -equivalent. We then see the \mathcal{S} -equivalence class of an $SU(n)$ bundle is determined by n points q_1, \dots, q_n (counted with multiplicities) satisfying $q_1 + \dots + q_n = 0$.

Thus, we can describe the moduli space of \mathcal{S} -equivalence classes of $SU(n)$ bundles as follows. Let $W := H^0(E, \mathcal{O}(n0))$ be the space of meromorphic functions ϕ that have a pole of at most order n at 0 , with no other poles. By Abel’s Theorem ϕ must have n zeros satisfying $q_1 + \dots + q_n = 0$. If ϕ has a pole of order less than n at 0 , we interpret this as some of the q_i are 0 . The zeros of ϕ are preserved under multiplication by an element of \mathbb{C}^* , and so the moduli space is $\mathbb{P}W \cong \mathbb{P}^{n-1}$.

Next consider a projective elliptic fibration $\pi : X \rightarrow \mathbb{P}^1$, with singular fibers of type I_1 or II (this extra assumption gives that X coincides with its Weierstrass model). Over each generic point x in the base there is an elliptic curve $E_x := \pi^{-1}(x)$ and a moduli space \mathbb{P}^{n-1}_x of $SU(n)$ bundles. Friedman–Morgan–Witten demonstrate that the projective spaces glue together to form a \mathbb{P}^{n-1} bundle over the base, which we denote by \mathcal{W} . A holomorphic $SU(n)$ bundle V over X which restricts to a semi-stable bundle on each fiber determines a section s of \mathcal{W} , which in turn defines a divisor $D_V \subset X$. Specifically, each point x in the base determines n points in E_x , thus D_V is an n -fold ramified cover of \mathbb{P}^1 . More precisely, in Sect. 4 of [16] it is demonstrated:

Theorem 3.6 (Friedman–Morgan–Witten [16]) *Let $\pi : X \rightarrow \mathbb{P}^1$ be an elliptic fibration with a section σ , with singular fibers of type I_1 or II . Let V be a holomorphic bundle of rank n over X . Assume that the restriction of V to a generic fiber of π is semi-stable and regular. Then, there exists a divisor*

$$D_V \in |n\sigma(\mathbb{P}^1) + kl|,$$

called the spectral cover associated to V , where l denotes the effective divisor class of the fibers of π , $k \in \mathbb{Z}$ satisfies $0 \leq k \leq c_2(V)$. For a generic $x \in \mathbb{P}^1 \setminus Z_\pi$, if $V|_{E_x} \cong \bigoplus_{j=1}^\ell \mathcal{O}_E(q_j - 0) \otimes \mathcal{I}_{r_j}$, then

$$D_V \cap E_x = \sum_{j=1}^\ell r_j q_j \in |n\sigma(x)|.$$

If V admits a spectral cover D_V which is reduced and irreducible, then D_V has a finite number of ramification points. Let Z_D denote the image of these ramification point under π . Then, for any $x \in \mathbb{P}^1 \setminus (Z_\pi \cup Z_D)$, we have $D_V \cap E_x = \sum_{j=1}^n q_j$ with all q_j distinct, and thus

$$V|_{E_x} \cong \mathcal{O}_E(q_1 - 0) \oplus \cdots \oplus \mathcal{O}_E(q_n - 0).$$

This verifies the holomorphic structure assumption on $V|_{E_x}$ in Theorem 1.1. Furthermore the points q_j vary holomorphically in x . The condition that D_V be reduced and irreducible also guarantees that the bundle V is stable with respect to ω_t for small t . This can be used to construct many examples of connections Ξ_i that satisfy the assumptions of our main theorem.

Theorem 3.7 (Theorem 7.4 in [16]) *If the spectral cover D_V constructed above is reduced and irreducible, then V is stable with respect to $\pi^*[\omega_{\mathbb{P}^1}] + t^2[\omega_X]$ for any ample class $[\omega_X]$ on X , for all $0 < t^2 \leq (\frac{n^3}{4}c_2(V))^{-1}$.*

We end this section with the construction of Ξ_0 from Corollary 1.2, which is a local *HYM* connection that determines the limit $A_0 = \Xi_0|_E$ on each fiber. Although the limiting connection A_0 is expressed in holomorphic coordinates in (1.1), here we find it easier to work with our coordinates (x^1, x^2, y^1, y^2) from the previous section. Both viewpoints are, of course, equivalent.

Consider $X_U := \pi^{-1}(U)$ for some simply connected $U \subset \mathbb{P}^1 \setminus (Z_\pi \cup Z_D)$. As a first step, we consider the case where V has rank one. Since it has degree zero, the bundle V is topologically trivial along each fiber and thus topologically trivial on X_U . We equip V with a trivial metric H_0 , and fix a unitary frame. For a fiber E_x we have assumed the restriction $V|_{E_x} \cong \mathcal{O}_E(q - 0)$, with q varying holomorphically in the base. We decompose q as follows

$$q(x^1, x^2) = \theta_1(x^1, x^2) - \tau\theta_2(x^1, x^2). \tag{3.3}$$

Recall E_x is determined by the quotient $T_x B/\Lambda$, and so the point q can be lifted to a point \tilde{q} in $T_x B \cong \mathbb{C}$. Now, if $\tau(x^1, x^2)$ gives the complex structure on E_x , we can define the holomorphic structure on V by

$$\bar{\partial}_q := \bar{\partial} - \frac{\pi \tilde{q}}{\text{Im}(\tau)} d\bar{z},$$

where z is the complex coordiante defined in (2.4). At the end of the section we will demonstrate that the connections we construct are independent of the lift from q to \tilde{q} , and therefore well defined.

Given the above holomorphic structure, and using that H_0 is the trivial metric, the Chern connection can be computed as

$$\Xi_0 = 2\pi \mathbf{i} \left(\theta_1 dy^1 + \theta_2 dy^2 \right).$$

Since each θ_j only depends on the base coordinates, $\Xi_0|_{E_x}$ is flat on each fiber. Using (3.3), one can check that the expression for $\Xi_0|_{E_x} =: A_0$ is equivalent to (1.1) in the statement of the main theorem. The holonomy around each period in E_x is given by $e^{2\pi \mathbf{i} \theta_1}$ and $e^{2\pi \mathbf{i} \theta_2}$, respectively.

Proposition 3.8 *The connection Ξ_0 is HYM with respect to all three complex structures $I, J,$ and $K,$ on X_U .*

Proof As a first step we show

$$\phi^{ij} \frac{\partial}{\partial x^i} \theta_j = 0 \tag{3.4}$$

and

$$\frac{\partial}{\partial x^i} \theta_j = \frac{\partial}{\partial x^j} \theta_i. \tag{3.5}$$

To see this, because both τ and q are holomorphic in the base, one can compute

$$\begin{aligned} 0 &= \frac{\partial}{\partial \bar{w}} q = \frac{1}{2} \left(\frac{\partial}{\partial x^1} - \bar{\tau} \frac{\partial}{\partial x^2} \right) (\theta_1 - \tau \theta_2) \\ &= \frac{1}{2} \left(\frac{\partial}{\partial x^1} \theta_1 - \bar{\tau} \frac{\partial}{\partial x^2} \theta_1 - \tau \frac{\partial}{\partial x^1} \theta_2 + |\tau|^2 \frac{\partial}{\partial x^2} \theta_2 \right). \end{aligned}$$

Now, using (2.2), the norm of τ is given by

$$|\tau|^2 = \frac{(1 - \mathbf{i}\phi_{12})(1 + \mathbf{i}\phi_{21})}{\phi_{22}^2} = \frac{1 + \phi_{12}^2}{\phi_{22}^2} = \frac{\phi_{11}}{\phi_{22}},$$

where for the last equality we used $\det(\phi_{ij}) = 1$. Thus

$$0 = \frac{1}{2\phi_{22}} \left(\phi_{22} \frac{\partial}{\partial x^1} \theta_1 + (\mathbf{i} - \phi_{12}) \frac{\partial}{\partial x^2} \theta_1 - (\mathbf{i} + \phi_{12}) \frac{\partial}{\partial x^1} \theta_2 + \phi_{11} \frac{\partial}{\partial x^2} \theta_2 \right).$$

Since both the real and imaginary parts vanish, (3.4) and (3.5) follow. In particular (3.5) allows us to simplify our notation and denote $\frac{\partial}{\partial x^i} \theta_j$ as θ_{ij} , where the indices commute.

The curvature of Ξ_0 is now given by

$$F_{\Xi_0} = 2\pi\mathbf{i} \theta_{ij} dx^i \wedge dy^j.$$

Right away it follows that $F_{\Xi_0} \wedge \omega_I = 0$ for all t . Furthermore, (3.4) implies $F_{\Xi_0} \wedge \omega_J = 0$ and (3.5) gives $F_{\Xi_0} \wedge \omega_K = 0$. Thus Ξ_0 is a holomorphic and HYM with respect to each complex structure. □

We now turn to the general case. Assume $V|_{E_x} \cong \bigoplus_{j=1}^n \mathcal{O}_E(q_j - 0)$, with each q_j is distinct. As before write $q_j = \theta_1^j - \tau\theta_2^j$, and construct diagonal matrices Θ_1 and Θ_2 with eigenvalues θ_1^j and θ_2^j , respectively. Consider the connection

$$\Xi_0 = 2\pi\mathbf{i} \left(\Theta_1 dy^1 + \Theta_2 dy^2 \right), \tag{3.6}$$

Its curvature is given by

$$F_{\Xi_0} = 2\pi\mathbf{i} \Theta_{ij} dx^i \wedge dy^j.$$

It is clear that $F_{\Xi_0}|_{E_x} = 0$ for every fiber in X_U . Furthermore, by Proposition 3.8 we have

$$F_{\Xi_0} \wedge \omega_I = F_{\Xi_0} \wedge \omega_J = F_{\Xi_0} \wedge \omega_K = 0.$$

Thus Ξ_0 is a local HYM connection with respect to each complex structure, although we only focus on I in this paper.

Finally, we demonstrate that the lift of each point q_i in E_x to \mathbb{C} is well defined. Recall that in the coordinates (y^1, y^2) , our lattice Λ is the standard lattice given by $(1, 1)$. Now, suppose we have another connection Ψ satisfying

$$\Xi_0 - \Psi = 2\pi\mathbf{i} \left(M_1 dy^1 + M_2 dy^2 \right),$$

where $M_1 = \text{diag}(\alpha_1, \dots, \alpha_n)$ and $M_2 = \text{diag}(\beta_1, \dots, \beta_n)$ are both diagonal matrices of integers, which means both Ξ_0 and Ψ define the same points on E_x . If u is the gauge transformation given by

$$u = \text{diag}(e^{2\pi\mathbf{i}(\alpha_1 y^1 + \beta_1 y^2)}, \dots, e^{2\pi\mathbf{i}(\alpha_n y^1 + \beta_n y^2)}),$$

we have $\Xi_0 - \Psi = -duu^{-1}$. Furthermore because all the α_i and β_i are integers, u descends to a smooth gauge transformation on the torus fibers, and thus the connections are gauge equivalent on X_U . Finally, since the points q_i add up to 0 in the group law on E_x , we can find a lift to \mathbb{C} where the points still add to 0, and Ξ_0 will be trace free.

4 Bubbling

We now present our bubbling argument, following the first two cases from [14]. Consider a sequence of Hermitian–Yang–Mills connections Ξ_i , corresponding to $\omega_{t_i} \in [\omega_{\mathbb{P}^1} + t_i^2 \omega_X]$ as $t_i \rightarrow 0$. Currently our argument depends on the sequence of connections we choose, although one can hope that with further analysis the set where bubbles occur can be uniquely identified by (V, ∂_{Ξ}) .

Choose a compact set $K \subset \mathbb{P}^1 \setminus Z_{\pi}$, where as before Z_{π} is the image of the singular fibers under π . In a neighborhood U of any point $x \in K$, we can choose affine coordinates (x^1, x^2) where x is at the origin, and coordinates (x^1, x^2, y^1, y^2) on $X_U := \pi^{-1}(U)$. The curvature of Ξ_i can be decomposed as

$$F_{\Xi_i} = F_{B_i} + F_{A_i} + \kappa_i,$$

where F_{B_i} and F_{A_i} denote the base and fiber directions of the curvature, and κ_i denotes the mixed terms. For each $x \in K$ we define the quantity

$$m_i(x) := \|F_{B_i}\|_{L^\infty(E_x, H_0, g_X)} + \frac{1}{t_i^2} \|F_{A_i}\|_{L^\infty(E_x, H_0, g_X)} + \|\kappa_i\|_{L^\infty(E_x, H_0, g_X)}^2,$$

where g_X is the metric associated to the fixed Kähler form ω_X .

Proposition 4.1 *There exists a finite number of points $\{p_1, \dots, p_\ell\} \subset K$ such that for any compact set $K' \subset K \setminus \{p_1, \dots, p_\ell\}$,*

$$\lim_{i \rightarrow \infty} t_i^2 \|m_i\|_{L^\infty(K')} = 0.$$

In particular, for any $x \in K \setminus \{p_1, \dots, p_\ell\}$,

$$\lim_{i \rightarrow \infty} \|F_{A_i}\|_{L^\infty(E_x)} = 0.$$

Proof Let x_i be a sequence of points in K for which $t_i^2 m_i(x_i)$ does not approach zero. We will show a finite amount of energy must bubble off along this sequence. Thus, by the total energy bound (Lemma 3.3), there can only be a finite number of points in K where bubbling occurs.

The proof closely follows the arguments of [14,17,19,21] and is divided into two cases. The first case occurs when $t_i^2 m_i(x_i)$ is unbounded, and the second when $t_i^2 m_i(x_i)$ stays bounded above, yet is also bounded away from zero. Unless mentioned otherwise, all bundle norms in this section are with respect to H_0 , so we suppress H_0 from our notation for simplicity.

Case 1 $t_i^2 m_i(x_i)$ is unbounded.

Given our sequence of points x_i , there exists corresponding points a_i in E_{x_i} where the supremum is obtained, and without loss of generality we can assume $(x_i, a_i) \rightarrow (x_0, a_0)$. Let $D_r(x_i)$ denote a disc of radius r in the metric $g_{\mathbb{P}^1}$ (corresponding to $\omega_{\mathbb{P}^1}$) in the base. We will show there exists a universal constant $\epsilon_0 > 0$, so that the inequality

$$\liminf_{i \rightarrow \infty} \int_{\pi^{-1}(D_r(x_i))} |F_{\Xi_i}|_{g_i}^2 dV_{g_i} > \epsilon_0 \tag{4.1}$$

holds for any small $r > 0$. Since the total energy is finite, a standard covering argument then shows that there can only be finitely many bubbles of this type.

Suppose (4.1) does not hold. Then there exists an r_0 so that for sufficiently large i ,

$$\int_{\pi^{-1}(D_{r_0}(x_i))} |F_{\Xi_i}|_{g_i}^2 dV_{g_i} \leq \epsilon_0.$$

It follows from (2.6) that there is a universal constant $c > 0$, so that the g_i -geodesic ball $B_i := B_{cr_0 t_i}(x_i, a_i)$ is contained in $\pi^{-1}(D_{r_0}(x_i))$. In particular we have the following bound

$$\int_{B_i} |F_{\Xi_i}|_{g_i}^2 dV_{g_i} \leq \epsilon_0.$$

We now rescale our coordinates and metrics. Consider the coordinate change

$$\lambda_i(x, y) = \left(t_i(x + x_i), y + a_i \right),$$

and let $\tilde{\omega}_i = t_i^{-2} \lambda_i^* \omega_i$. To compare this to the scaling L_{t_i} , defined in (2.7) in Sect. 2, note that pulling the metric back by L_{t_i} dilates the shrinking fibers, while $\tilde{\omega}_i$ is a combination of shrinking the base coordinates and then dilating the metric. Thus both scalings have the same effect, although with different coordinates.

By (2.6), $\tilde{\omega}_i$ is uniformly equivalent to the Euclidean metric in the scaled coordinates, which we denote by \tilde{g}_0 . Moreover the ball B_i pulls back to a \tilde{g}_i -geodesic ball \tilde{B}_i , which contains a Euclidean ball \tilde{B} . The ball \tilde{B} can be chosen to have uniform size independent of i .

Now, if $\tilde{\Xi}_i$ is the pull-back connection to these new coordinates, then the HYM equation is again satisfied

$$F_{\tilde{\Xi}_i} \wedge \tilde{\omega}_i = 0 \quad \text{and} \quad (F_{\tilde{\Xi}_i})^{0,2} = 0. \tag{4.2}$$

Change of variables, and the scale invariance of the Yang–Mills energy in dimension four, implies

$$\|F_{\tilde{\Xi}_i}\|_{L^2(\tilde{B}, \tilde{g}_0)}^2 \leq \|F_{\tilde{\Xi}_i}\|_{L^2(\tilde{B}_i, \tilde{g}_i)}^2 = \|F_{\Xi_i}\|_{L^2(B_i, g_i)}^2 \leq \epsilon_0.$$

Since \tilde{g}_i is uniformly equivalent to \tilde{g}_0 on \tilde{B} for large i , and $\tilde{\Xi}_i$ satisfies (4.2), we can apply the standard ϵ -regularity argument for Yang–Mills connections on a fixed ball \tilde{B} [35, Theorem 4.8]. Thus, for ϵ_0 small enough (depending only on the real dimension 4 of X), the above L^2 control implies

$$|F_{\tilde{\Xi}_i}|_{\tilde{g}_i}^2(0) \leq C,$$

for some constant C independent of i . Equivalence of metrics implies control of $F_{\tilde{\Xi}_i}$ with respect to \tilde{g}_0 , which in components gives the following bound

$$|F_{\tilde{B}_i}|_{\tilde{g}_0}(0) + |F_{\tilde{A}_i}|_{\tilde{g}_0}(0) + |\tilde{\kappa}_i|_{\tilde{g}_0}^2(0) \leq C.$$

Scaling back, we see

$$t_i^2 |F_{B_i}|_{g_X}(0) + |F_{A_i}|_{g_X}(0) + t_i^2 |\kappa_i|_{g_X}^2(0) \leq C.$$

Hence we achieve control of $t_i^2 m_i(0)$, which we have assumed diverges, a contradiction. Let W_1 denote the set of points in K at which bubbles of this type appear.

Case 2 $t_i^2 m_i(x_i)$ is bounded above and away from zero.

In this case an instanton on $\mathbb{C} \times E$ bubbles off. We follow the outline of [19,21], and use an energy quantization result of Wehrheim.

Suppose $x_i \rightarrow x_0 \in K \setminus W_1$, and let $D_{2\rho}(x_i)$ denote a disc of radius 2ρ in the metric $g_{\mathbb{P}^1}$ in the base. Suppose there exists constants δ and Λ so that

$$\delta < t_i^2 m_i(x_i) \leq \sup_{D_{2\rho}(x_i)} t_i^2 m_i < \Lambda.$$

The rightmost inequality holds since for large enough i we can assume $D_{2\rho}(x_i) \subset K \setminus W_1$. By making ρ smaller if necessary, we can furthermore assume that $\pi^{-1}(D_{2\rho}(x_i))$ is topologically a product between a ball in \mathbb{C} and an elliptic curve E , although the complex structure may vary.

We preform the same scaling as in Case 1, and define $\tilde{\omega}_i = t_i^{-2} \lambda_i^* \omega_i$. Again this involves rescaling the metric and applying a dilation. The disk $D_{2\rho}(x_i)$ pulls back to $\tilde{D}_{2\rho/t_i}(0)$, the geodesic disk with respect to the Euclidean metric \tilde{g}_0 in the scaled coordinates. Starting from Proposition 2.4, the arguments used in the proof of [36, Theorem 1.1] (cf. pages 2936-2937) give that $\tilde{\omega}_i$ converge sub-sequentially and smoothly to a limiting flat product metric ω_∞ on $\mathbb{C} \times E$.

Our sequence of scaled connections $\tilde{\Xi}_i$ is defined on $\pi^{-1}(\tilde{D}_{2\rho/t_i}(0))$, and for any point $p \in \tilde{D}_{2\rho/t_i}(0)$, we have

$$\begin{aligned} |F_{\tilde{B}_i}|_{\tilde{g}_0}(p) + |F_{\tilde{A}_i}|_{\tilde{g}_0}(p) + |\tilde{\kappa}_i|_{\tilde{g}_0}^2(p) &= t_i^2 |F_{B_i}|_{g_0}(p) + |F_{A_i}|_{g_0}(p) + t_i^2 |\kappa_i|_{g_0}^2(p) \\ &< \sup_{D_{2\rho}(x_i)} t_i^2 m_i < \Lambda. \end{aligned}$$

This implies $|F_{\tilde{\Xi}_i}|_{\tilde{g}_i}$ is uniformly bounded. By strong Uhlenbeck compactness [37, Corollary 1.4 and Theorem 1.5], this bound implies there exists a subsequence of

connections which converges smoothly, modulo unitary gauge transformations, to a limiting connection $\tilde{\Xi}_\infty$ on the trivial $SU(n)$ bundle over $\mathbb{C} \times E$. The connection $\tilde{\Xi}_\infty$ will be ASD with respect to the limiting product metric ω_∞ . Furthermore, by assumption, we have

$$|F_{\tilde{B}_i}|_{\tilde{g}_0}(0) + |F_{\tilde{A}_i}|_{\tilde{g}_0}(0) + |\tilde{\kappa}_i|_{\tilde{g}_0}^2(0) > \delta,$$

and it follows that the limiting connection is not flat. An energy quantization result of Wehrheim [38, Remark 1.2] implies there exists a universal constant $\epsilon_0 > 0$ so that

$$\mathcal{YM}(\tilde{\Xi}_\infty, \tilde{g}_0) \geq 2\epsilon_0.$$

This implies that there exists an $R > 0$, so that for i sufficiently large,

$$\epsilon_0 < \int_{\tilde{D}_{R/2}(0) \times E} |F_{\tilde{\Xi}_i}|_{\tilde{g}_i}^2 dV_{\tilde{g}_i} = \int_{\pi^{-1}(D_{R_i}(x_i))} |F_{\Xi_i}|_{g_i}^2 dV_i.$$

Thus there can be only finitely many bubbles of this type, and denote the set of all such bubbles by W_2 . This concludes the proof of Proposition 4.1. □

We conclude this section by noting that Proposition 4.1, in conjunction with our convergence argument in Sect. 6, implies that on a generic fiber, the connections A_i will converge to a limiting flat connection A_0 . The following section is devoted to identifying this limiting flat connection explicitly.

5 Gauge Fixing over an Elliptic Curve

In this section we work on a fixed fiber of π , denoted E for simplicity, satisfying $V|_E \cong \bigoplus_{j=1}^n \mathcal{O}_E(q_j - 0)$ with each q_j distinct and $q_1 + \dots + q_n = 0$. By the results of Sect. 3, if V defines a spectral cover D_V which is reduced and irreducible, this happens generically.

Equip E with the fixed Kähler form $\omega_0 = dy^1 \wedge dy^2$ and let g_0 denote the corresponding metric. Recall that E carries the complex coordinate $z = \tau y^1 + y^2$. Denote the restriction $V|_E$ by V_0 . Since V_0 is of the form $\bigoplus_{j=1}^n \mathcal{O}_E(q_j - 0)$, it can be naturally identified with $E \times \mathbb{C}^n$, equipped with the complex structure

$$\bar{\partial}_{A_0} := \bar{\partial} - \frac{\pi Q}{\text{Im}(\tau)} d\bar{z},$$

where Q is a diagonal matrix with entries \tilde{q}_i (recall \tilde{q}_i are the lifts of the points q_i to \mathbb{C}). Let H_0 be the trivial metric on \mathbb{C}^n , and let A_0 be the Chern connection associated to $\bar{\partial}_{A_0}$ and H_0 . Using (3.3), in addition to $dz|_E = \tau dy^1 + dy^2$, one can explicitly check that that $A_0 = \Xi_0|_E$, where Ξ_0 is given by (3.6).

Now, given our sequence of connections Ξ_i on V , we have the sequence of restricted connections $A_i := \Xi_i|_E$ on V_0 . Again, because our sequence of connections arises

from the Donaldson–Uhlenbeck–Yau Theorem, we know that A_i lies in the complexified gauge orbit of A_0 , and thus A_i and A_0 define isomorphic complex structures. As a result, after transforming A_0 by a unitary gauge transformation if necessary, we can write $A_i = e^{s_i}(A_0)$ for a trace free Hermitian endomorphism s_i . The main result of this section is:

Theorem 5.1 *Let $e^s(A_0)$ be a connection on V_0 given by the action of a trace free Hermitian endomorphism s . There exists constants $\epsilon_0 > 0$, and $C_0 > 0$, depending only on g_0, A_0 , and H_0 , so that the following holds. If the curvature of $e^s(A_0)$ satisfies*

$$\|F_{e^s(A_0)}\|_{C^0(g_0, H_0)}^2 \leq \epsilon_0,$$

then there exists another trace free Hermitian endomorphism s' which satisfied both

$$e^s(A_0) = e^{s'}(A_0) \quad \text{and} \quad \|s'\|_{C^0(g_0, H_0)} \leq C_0.$$

Because V_0 is poly-stable, and not stable, the above theorem is the best C^0 control that one can expect. For example, since A_0 is flat, if e^s is a diagonal matrix of constants c_1, \dots, c_n , then $e^s(A_0)$ will still be flat. However, one eigenvalue c_i can be arbitrarily large while still preserving the condition $\det(e^s) = 1$ (recall that $e^s \in SL(V_0)$). Thus, one can never expect C^0 control for s . The main idea of the above theorem is that, by a suitable choice of normalization, one can construct a related complex gauge transformation that yields the same connection, yet with the desired C^0 control.

We first demonstrate several preliminary results. For the remainder of the section, unless specified, all norms are taken with respect to the metrics g_0 and H_0 , and we remove this from our notation for simplicity.

Proposition 5.2 *Let s be a trace free Hermitian endomorphism such that the diagonal entries of s have zero average when integrated over E . Then there exists a constant C_p , independent of s , so that*

$$\|s\|_{L^2(E)} \leq C_p \|\bar{\partial}_{A_0} s\|_{L^2(E)}. \tag{5.1}$$

Proof Note that diagonal entries of s can always be defined as the entries that preserve each subbundle $\mathcal{O}_E(q_i - 0) \subset V_0$. Now, assume that the inequality does not hold. Then there exists a sequence of endomorphisms s_k satisfying the assumptions of the proposition, along with the inequality

$$\int_E |\bar{\partial}_{A_0} s_k|^2 \leq \frac{1}{k} \int_E |s_k|^2.$$

Let $\tilde{s}_k := s_k / \|s_k\|_{L^2(E)}$. Then

$$\int_E |\bar{\partial}_{A_0} \tilde{s}_k|^2 \leq \frac{1}{k}.$$

Because s is Hermitian with respect to H_0 , we have $|\bar{\partial}_{A_0}\tilde{s}_k| = |\partial_{A_0}\tilde{s}_k|$. This allows us to conclude that the sequence \tilde{s}_k converges weakly in L^2_1 (and strongly in L^2) to an endomorphism s_∞ satisfying

$$\|s_\infty\|_{L^2(E)} = 1 \quad \text{and} \quad \|\bar{\partial}_{A_0}s_\infty\|_{L^2(E)} = 0.$$

Now, because Q is diagonal, if s_∞ has entries (a_{ij}) , then the diagonal entries of $\bar{\partial}_{A_0}s_\infty$ are of the form $\bar{\partial} a_{ii}$. Since $\|\bar{\partial}_{A_0}s_\infty\|_{L^2} = 0$, we see the diagonal entries of s_∞ are constant. Furthermore, by assumption the diagonal entries of \tilde{s}_k have zero average, and by strong convergence in L^2 we conclude the diagonal entries of s_∞ must also have zero average. Thus these entries vanish entirely. Now, because the points q_i are distinct, the automorphism group of V_0 is precisely n dimensional [16, Lemma 1.13]. Thus if the diagonal entries of s_∞ vanish, s_∞ must vanish entirely. In other words, if s_∞ had any non-vanishing off diagonal entries, they would define a holomorphic map between line bundles $\mathcal{O}_E(q_i - 0)$ and $\mathcal{O}_E(q_j - 0)$ for distinct points q_i and q_j , which is impossible. So $s_\infty = 0$ yet $\|s_\infty\|_{L^2(E)} = 1$, a contradiction. \square

Next consider the following function spaces, equipped with the L^2 norm.

Definition 5.3 Let $\text{Herm}_0(V_0)$ be the space of trace free Hermitian endomorphisms of V_0 , and furthermore let $\text{Herm}^{\perp 0}(V_0)$ denote the subspace consisting of those endomorphisms whose diagonal entries have zero average on E .

Consider $\Upsilon(\cdot) \in \text{End}(\mathfrak{gl}(V_0))$, defined by

$$\Upsilon(s) = \frac{e^{\text{ad}_s} - 1}{\text{ad}_s}.$$

From the definition of the complexified gauge action (3.1), we have

$$\partial_{e^s(A_0)} = \partial_{A_0} + e^{-s}(\partial_{A_0}e^s) \quad \text{and} \quad \bar{\partial}_{e^s(A_0)} = \bar{\partial}_{A_0} - (\bar{\partial}_{A_0}e^s)e^{-s}. \tag{5.2}$$

This allows one to compute

$$e^s(A_0) = A_0 + \Upsilon(-s)\partial_{A_0}s - \Upsilon(s)\bar{\partial}_{A_0}s \tag{5.3}$$

(for instance, see [39, Appendix A]).

Define the map $\mathcal{N}(s) := e^s(A_0)$, which maps $\text{Herm}_0(V_0)$ into the affine space of connections centered at A_0 , equipped with the L^2 norm. Let \mathcal{A} denote the image of the map \mathcal{N} . Using (5.3), we see the derivative of \mathcal{N} at 0 is given by

$$L(s) := \mathcal{N}'(0)(s) = \partial_{A_0}s - \bar{\partial}_{A_0}s.$$

The tangent space to $\text{Herm}_0(V_0)$ is again $\text{Herm}_0(V_0)$. Note that for any $s \in \text{Herm}_0(V_0)$, if \hat{s} is a diagonal matrix of constants given by the averaging the diagonals of s over E , then $L(s) = L(s - \hat{s})$, and so both $\text{Herm}_0(V_0)$ and $\text{Herm}^{\perp 0}(V_0)$ have the same image under L . Proposition 5.2 shows that L has trivial Kernel on $\text{Herm}^{\perp 0}(V_0)$. Thus, not only

can we conclude that the restriction of L to $\text{Herm}_0^\perp(V_0)$ is invertible, but (5.1) shows in addition that L has bounded inverse. The contraction mapping principle implies there exists a small neighborhood $\mathcal{U} \subset \mathcal{A}$ of the connection A_0 , and a set $\mathcal{V} \subset \text{Herm}_0^\perp(V_0)$ in the tangent space to $\text{Herm}_0(V_0)$ at 0, so that $\mathcal{V} \rightarrow \mathcal{U}$ is a diffeomorphism onto its image.

Summing up, we have proved the following:

Lemma 5.4 *There exist constants δ_0 and Λ_0 , which depend only on A_0, H_0 and g_0 , so that the following holds. If $A \in \mathcal{A}$ satisfies $\|A - A_0\|_{L^2(E)} < \delta_0$, there exists $s \in \text{Herm}_0^\perp$, such that*

$$A = e^s(A_0),$$

and

$$\|s\|_{L^2(E)} \leq \Lambda_0 \|e^s(A_0) - A_0\|_{L^2(E)}.$$

We now turn to one final lemma. Consider the same constant $\delta_0 > 0$ from above, and let $C_0 > 0$ be a fixed constant, to be determined in the proof of Theorem 5.1.

Lemma 5.5 *Let s be a trace free Hermitian endomorphism, and A a flat connection on V_0 . Given constants $\delta_0 > 0$ and $C_0 > 0$, there exists a constant ϵ_0 , depending only on H_0, g_0, δ_0 , and C_0 , so that if $\|s\|_{C^0} \leq C_0$ and $\|F_{e^s(A)}\|_{C^0(E)} < \epsilon_0$, then*

$$\|e^s(A) - A\|_{L^2(E)} < \frac{\delta_0}{2}.$$

Proof To begin, we see how the curvature of A is related to the curvature of $e^s(A)$. Using (5.2) one can compute

$$e^{-s} F_{e^s(A)} e^s - F_A = \bar{\partial}_A(e^{-2s} \partial_A e^{2s}) = e^{-2s} \left(\bar{\partial}_A \partial_A e^{2s} - \bar{\partial}_A e^{2s} e^{-2s} \partial_A e^{2s} \right),$$

which implies

$$-\Delta_{g_0} \text{Tr}(e^{2s}) + |e^{-s} \partial_A e^{2s}|_{g_0}^2 = \text{Tr}(e^{2s} i \Lambda_{\omega_0} F_{e^s(A)}). \tag{5.4}$$

Integrating over E yields

$$\int_E |e^{-s} \partial_A e^{2s}|^2 < e^{C_0} \epsilon_0.$$

The C_0 bound for s , along with $\det(e^s) = 1$, demonstrates the eigenvalues of e^s are bounded above and below. Thus the left hand side above controls the L^2 norm of the difference $e^s(A) - A$, and so

$$\|e^s(A) - A\|_{L^2(E)} < C \epsilon_0.$$

Here C only depends on C_0, g_0 , and H_0 . Choose ϵ_0 small so $C \epsilon_0 < \delta_0/2$. □

As we turn to the proof of Theorem 5.1, recall that the constants δ_0 and Λ_0 depend only on A_0 , H_0 and g_0 . Thus, by the above lemma, if we can show C_0 depends only on these quantities, ϵ_0 will depend only on these quantities.

Proof of Theorem 5.1 Our first task is to specify C_0 . Fix an endomorphism $s \in \text{Herm}_0(V_0)$ satisfying

$$\|F_{e^s(A_0)}\|_{C^0}^2 \leq \epsilon_0.$$

Using this curvature bound, along with the inequality

$$-\Delta_{g_0}|s|^2 \leq |s| |F_{e^s(A_0)}|$$

(see for instance Proposition A.6 in [39]), we can apply Moser iteration to conclude

$$\max\{|s|^2, 1\} \leq C_1 \|s\|_{L^2} (1 + \epsilon_0). \tag{5.5}$$

Here C_1 only depends on g_0 and H_0 . We now set $C_0 := 2C_1 \Lambda_0$. This shows C_0 , and subsequently ϵ_0 , depends only on the initial setup.

The main idea of the proof is as follows. We construct a path of Hermitian endomorphisms, so that the curvature of the induced connections along this path is bounded by ϵ_0 . We show the endpoint of our path satisfies the conclusion of the theorem, and then apply a method of continuity argument to conclude our desired result for s . Naively, one may first try to connect e^s to Id_{V_0} by the path e^{ts} for $t \in [0, 1]$. However, for arbitrary initial s it is not clear that curvature stays bounded by ϵ_0 along this path, which is an important for the argument. Instead, we follow the Yang–Mills flow.

Following Donaldson [12, Sect. 1.1], we consider a path of complex gauge transformations $g(t)$, satisfying

$$\dot{g}(t)g(t)^{-1} = -i \star F_{g(t)(A_0)} \quad g(0) = e^s.$$

On a Riemann surface, the above flow is referred to as the Kempf–Ness flow, and given a solution, the corresponding connections $A(t) := g(t)(A_0)$ solve the Yang–Mills heat flow:

$$\dot{A}(t) = -d_{A(t)}^* F_{A(t)} \quad A(0) = e^s(A_0).$$

By a Theorem of Rade [40, Theorem 2], there exists a limiting Yang–Mills connection A_∞ for which

$$\|A(t) - A_\infty\|_{L^2_1} \leq c t^{-\beta}.$$

Since V_0 is polystable, the limit connection A_∞ is also flat, and in the unitary gauge orbit of A_0 , and so there exists a unitary gauge transformation u_∞ for which $A_\infty = u_\infty^* A_0$.

Consider the trivial flow $u_\infty(t) = u_\infty$, which again satisfies the Kempf–Ness flow equation

$$\dot{u}_\infty(t)u_\infty(t)^{-1} = -i\star F_{u_\infty(t)(A_0)} = -i\star F_{A_\infty} = 0.$$

Define $\eta(t) \in \text{Herm}_0(V_0)$, and a path of unitary gauge transformations $u(t)$, by the equation

$$u_\infty u(t)e^{\eta(t)} = g(t). \tag{5.6}$$

Thus, $u(t)e^{\eta(t)}$ relates our two solutions of the Kempf–Ness flow. By Proposition 4.13 in [41], both $u(t)$ and $\eta(t)$ are bounded in L^2_2 . In fact, this L^2_2 bound is proven following the general argument of [42]. The authors demonstrate that if M is a complete, connected, simply connected Riemannian manifold of nonpositive sectional curvature, which admits a function $\Phi : M \rightarrow \mathbb{R}$ which is convex along geodesics, then given two negative gradient flow lines of Φ , the geodesic distance between these two flow lines stays bounded. In our case, the role of M is taken by the space of Hermitian metrics, and the function Φ is Donaldson’s functional (see [12,41,42], for a precise definition of Φ).

The bound on $\eta(t)$, along with convergence of $F_{A(t)}$ to zero in L^2 , allows us to conclude by Lemma 5.5 that there exists a T sufficiently large, so that

$$\|(u_\infty u(T)e^{\eta(T)}u(T)^{-1}u_\infty^{-1})((u_\infty u(T))^*A_0) - (u_\infty u(T))^*A_0\|_{L^2} \leq \frac{\delta_0}{2}.$$

For simplicity we denote the fixed unitary gauge transformation $u_\infty u(T)$ by u , and define the path of Hermitian endomorphism $e^{\kappa(t)}$ by $ue^{\eta(t)}u^{-1}$. Then the above estimate can be written

$$\|e^{\kappa(T)}(u^*A_0) - u^*A_0\|_{L^2} \leq \frac{\delta_0}{2}. \tag{5.7}$$

It is along the path $e^{\kappa(t)}$ that we can now apply our method of continuity argument.

Let $\tilde{A}(t)$ be the path of connections given by $e^{\kappa(t)}(u^*A_0)$. Since $\tilde{A}(t) = (u_\infty u(t)u^{-1})^*A(t)$, where $A(t)$ solves the Yang–Mills flow and $u(t)$ is given by (5.6), we conclude that $\|F_{\tilde{A}(t)}\|_{C^0(E)} \leq \epsilon_0$ for all $t \in [0, T]$. This follows because the curvature is decreasing along the Yang–Mills flow, and the action of a unitary gauge transformation will not affect this norm. Also, the path $\tilde{A}(t)$ is smooth for $t \in [0, T]$.

To set up the method of continuity, consider the set $I \subseteq [0, T]$ consisting of times t for which there exists a trace free Hermitian endomorphism $\kappa'(t)$ which satisfies both

$$e^{\kappa'(t)}(u^*A_0) = e^{\kappa(t)}(u^*A_0) \tag{5.8}$$

and

$$\|\kappa'(t)\|_{C^0(g_0, H_0)} \leq C_0. \tag{5.9}$$

We prove $I = [0, T]$. First, we demonstrate $T \in I$ to conclude I is non-empty. By the estimate (5.7), we can apply Lemma 5.4 to $\tilde{A}(T)$, and conclude there exists a trace free Hermitian endomorphism $\kappa'(T)$ satisfying both (5.8) and an L^2 bound. By our Moser iteration bound this L^2 control can be improved to C_0 , and so $\kappa'(T)$ satisfies (5.9) as well. Thus $T \in I$. We now need that I is both open and closed with respect to the topology induced from the C^2 topology on $\text{Herm}(V_0)$. For the rest of the proof we use the notation $A := u^*A_0$.

Our next step is to show I is open. Let $t_0 \in I$, and consider the corresponding endomorphism e^{κ_0} . Construct a small neighborhood of e^{κ_0} with radius $\rho > 0$, where ρ is chosen so

$$\|e^\kappa - e^{\kappa_0}\|_{C^2(E)} < \rho$$

implies

$$\|e^\kappa(A) - e^{\kappa_0}(A)\|_{L^2(E)} < \delta_0/2.$$

Now, because $e^{\kappa_0} \in I$, there exists an endomorphism $e^{\kappa'_0}$ satisfying both $e^{\kappa_0}(A) = e^{\kappa'_0}(A)$ and $\|\kappa'_0\|_{C^0} \leq C_0$. Given our choice of ϵ_0 , by Lemma 5.5 we have

$$\|e^{\kappa_0}(A) - A\|_{L^2} = \|e^{\kappa'_0}(A) - A\|_{L^2} < \delta_0/2.$$

By the triangle inequality $\|e^\kappa(A_0) - A_0\|_{L^2} < \delta_0$, and thus Lemma 5.4 implies there exists an endomorphism $\kappa' \in \text{Herm}_0^1$ such that $e^\kappa(A) = e^{\kappa'}(A)$ and

$$\|\kappa'\|_{L^2} \leq \Lambda_0 \|e^{\kappa'}(A) - A\|_{L^2} < \Lambda_0$$

(we assumed $\delta_0 < 1$). By our Moser iteration bound (5.5), we conclude

$$\|\kappa'\|_{C^0} < 2\Lambda_0 C_1 = C_0,$$

which completes the proof of openness.

Finally we prove I is closed. Let t_i be a sequence of times in I converging to t , and let κ_i be the corresponding sequence of endomorphisms converging to κ in the C^2 topology. For each i , there exists κ'_i which are uniformly bounded in C^0 and satisfy $e^{\kappa_i}(A) = e^{\kappa'_i}(A)$. The complexified gauge action gives

$$e^{\kappa_i} \circ \bar{\partial}_A \circ e^{-\kappa_i} = e^{\kappa'_i} \circ \bar{\partial}_A \circ e^{-\kappa'_i},$$

from which we conclude $\bar{\partial}_A(e^{-\kappa_i} e^{\kappa'_i}) = 0$. Since $A = u^*A_0$, this implies $\bar{\partial}_{A_0}(u^{-1} e^{-\kappa_i} e^{\kappa'_i} u) = 0$. Since A_0 has only diagonal entries, we see the diagonal entries of $u^{-1} e^{-\kappa_i} e^{\kappa'_i} u$ must be constant. As before it then follows that the off diagonal terms must vanish, otherwise one would have a holomorphic map between line bundles

$\mathcal{O}_E(q_i - 0)$ and $\mathcal{O}_E(q_j - 0)$ for distinct points q_i and q_j . Taking the complex conjugate and using the fact that κ_i and κ'_i are Hermitian, we see that $h_i := u^{-1}e^{\kappa'_i}e^{-\kappa_i}u$ will be a diagonal matrix of constants.

Now, C^2 convergence of κ_i together with the C^0 control of κ'_i gives that the matrices h_i are uniformly bounded above and below. Since each h_i is a diagonal matrix of constants, after passing to a subsequence the h_i converge in C^0 to a limit h . This allows us to define an endomorphism $e^{\kappa'} := uhu^{-1}e^\kappa$, and by convergence of h_i and κ_i we have

$$e^{\kappa'_i} = uh_iu^{-1}e^{\kappa_i} \rightarrow e^{\kappa'}$$

in C^0 . Since $e^{\kappa'_i}$ is Hermitian, so is $e^{\kappa'}$, and thus κ' is Hermitian and satisfies (5.9). Furthermore, since for each i we have $\bar{\partial}_A(uh_iu^{-1}) = 0$, we in fact have $uh_iu^{-1}e^{\kappa_i} \rightarrow e^{\kappa'}$ in C^1 , and as a result we conclude $e^{\kappa'}(A) = e^\kappa(A)$. Thus (5.8) is also satisfied and $t \in I$.

Thus, I is open, closed, and nonempty, and as a result $e^{\kappa(0)} \in I$. In particular, there exists a $\kappa'(0)$ satisfying both $e^{\kappa(0)}(A) = e^{\kappa(0)'}(A)$ and $\|\kappa(0)'\|_{C^0} \leq C_0$. Now, define s' by

$$e^{s'} = u^{-1}e^{\kappa(0)'}u.$$

We see s' satisfies the desired C^0 bound. Furthermore,

$$e^{s'}(A_0) = u^{-1*}e^{\kappa(0)'}(u^*A_0) = u^{-1*}e^{\kappa(0)}(u^*A_0) = e^{\eta(0)}(A_0).$$

Yet we started the flow (5.6) at $g(0) = e^s$, so $\eta(0) = s$. This completes the proof of the theorem. □

6 Convergence

In this section, we complete the proof of Theorem 1.1. As before, let Z_π denote image of the singular fibers under π , and W_1 and W_2 the bubbling sets for our sequence of connections. We fix a point $x \in \mathbb{P}^1 \setminus (Z_\pi \cup W_1 \cup W_2)$, and denote the fiber over x by $E := \pi^{-1}(x)$. We use the notation $A_i := \Xi_i|_E$ and $A_0 := \Xi_0|_E$. As above equip E with the fixed flat metric $\omega_0 = dy^1 \wedge dy^2$. Unless otherwise specified, in this section all norms are taken with respect to the metrics g_0 and H_0 .

Recall our bubbling sequence at x is defined by

$$m_i(x) := \|F_{B_i}\|_{C^0(E, g_X)} + \frac{1}{t_i^2} \|F_{A_i}\|_{C^0(E, g_X)} + \|\kappa_i\|_{C^0(E, g_X)}^2.$$

Since $x \notin W_1 \cup W_2$, we have $t_i^2 m_i \rightarrow 0$ as $i \rightarrow \infty$, so $\|F_{A_i}\|_{C^0(E)} \rightarrow 0$. As mentioned at the end of Sect. 4 (and as we shall see below), this is enough to prove that A_i converges, along a subsequence and modulo gauge transformations, to a limiting flat connection. Our main result is identifying this limit.

Assume $V|_E = \mathcal{O}_E(q_1 - 0) \oplus \cdots \oplus \mathcal{O}_E(q_n - 0)$ for q_1, \dots, q_n distinct. Writing $A_i = e^{s_i}(A_0)$ for a sequence of Hermitian endomorphism s_i , for i large enough we can apply Theorem 5.1 to conclude there exists gauge transformations s'_i , which are uniformly bounded in C^0 , and satisfy $A_i = e^{s'_i}(A_0)$. Thus, as in the proof of Lemma 5.5,

$$\|A_i - A_0\|_{L^2(E)} \leq C \|e^{-s'_i} \partial_{A_0} e^{2s'_i}\|_{L^2(E)} \leq C \|F_{A_i}\|_{L^2(E)} \rightarrow 0. \tag{6.1}$$

Furthermore, since A_0 is flat, we can integrate by parts and change the order of derivatives to conclude:

$$\begin{aligned} \|\nabla^0(A_i - A_0)\|_{L^2(E)}^2 &= \int_{E_p} \text{Tr}(\nabla^0(e^{-2s_i} \nabla^0 e^{2s_i}) (\nabla^0(e^{-2s_i} \nabla^0 e^{2s_i}))^*) \\ &= \int_{E_p} \text{Tr}(\bar{\nabla}^0(e^{-2s_i} \nabla^0 e^{2s_i}) (\bar{\nabla}^0(e^{-2s_i} \nabla^0 e^{2s_i}))^*) \\ &= \|F_{A_i}\|_{L^2(E)}^2 \rightarrow 0. \end{aligned}$$

Thus we have demonstrated

$$\|A_i - A_0\|_{L^2_1(E)} \rightarrow 0,$$

which is the stated convergence in Theorem 1.1.

Next we prove smooth convergence, allowing for the action of unitary gauge transformations. Specifically, since x is away from the bubbling set, there exists a small disk $D_\rho(x)$ that does not intersect $W_1 \cup W_2$. We use the same coordinate transformations λ_i in the proof Proposition 4.1 sending x to the origin and scaling. Consider the rescaled metrics $\tilde{\omega}_i = t_i^{-2} \lambda_i^* \omega_i$. The set $\pi^{-1}(D_\rho(x))$ rescales to a set topologically equivalent to $\tilde{D}_{\frac{\rho}{t_i}}(0) \times E$ (although the complex structure will not be a product).

Our sequence of connections Ξ_i pulls back to $\tilde{\Xi}_i$, with fiber and base components

$$\tilde{A}_j^i = A_j^i \quad \text{and} \quad \tilde{B}_j^i = t_i B_j^i.$$

As before, the connections $\tilde{\Xi}_i$ are HYM with respect to $\tilde{\omega}_i$, and each $\tilde{\omega}_i$ is uniformly equivalent to the Euclidian metric for large i by Proposition 2.4. Also, since $D_\rho(x)$ is away from W_1 , the function $t_i^2 m_i(x)$ is uniformly bounded above on the disk. This implies the curvature $|F_{\tilde{\Xi}_i}|$ is uniformly bounded on $\pi^{-1}(\tilde{D}_{\frac{\rho}{t_i}}(0))$.

Applying strong Uhlenbeck compactness [37, Corollary 1.4 and Theorem 1.5] on the fixed compact set

$$\pi^{-1}(\tilde{D}_1(0)) \subset \pi^{-1}(\tilde{D}_{\frac{\rho}{t_i}}(0)),$$

there exists a sequence of gauge transformations u_i so that along a subsequence, $u_i^* \tilde{\Xi}_i$ converges smoothly to a limiting Yang–Mills connection $\tilde{\Xi}_\infty$. Restricting our attention

to the fiber E over the origin yields a sequence of connections $u_i^* \tilde{A}_i$ which converges smoothly to a limiting flat connection \tilde{A}_∞ . Note that our fiber coordinates are not scaled, and the restriction of $\tilde{\omega}_i$ to E is equivalent to the standard metric $\omega_0 = dy^1 \wedge dy^2$. Thus, on E , we see $u_i^* A_i$ converges smoothly to a flat connection A_∞ . The connection A_∞ may not equal A_0 , but it will lie in the unitary gauge orbit.

We conclude by remarking that estimates of the form (6.1) are common in these types of degeneration problems, for example see Proposition 3.1 in [18] or Theorem 1 in [43]. In our estimate, the curvature term is not raised to a power, and this holds because the specific form of our complex structure $V|_E = \bigoplus_{j=1}^n \mathcal{O}_E(q_j - 0)$ implies that the Yang–Mills energy functional is Morse–Bott at A_0 (see Definition 7.5 in [43]). Essentially, the argument in our proof of Proposition 5.2 gives that the kernel of the Hessian operator of the Yang–Mills energy functional can be identified with one forms valued in constant diagonal matrices, which also gives the tangent space to Yang–Mills connections at A_0 . We direct the reader to [43] for further details.

Acknowledgements We would like to thank T. Nishinou, for providing a revised draft of [20], and P. Feehan for introducing us to reference [43]. We also thank the referee for many valuable suggestions. V. Datar is partially supported by NSF RTG Grant DMS-1344991 and A. Jacob is partially supported by a grant from the Hellman Foundation, and a Simons collaboration grant.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Gross, M., Wilson, P.M.H.: Large complex structure limits of $K3$ surfaces. *J. Differ. Geom.* **55**(3), 475–546 (2000)
2. Gross, M., Tosatti, V., Zhang, Y.: Collapsing of abelian fibered Calabi–Yau manifolds. *Duke Math. J.* **162**(3), 517–551 (2013)
3. Gross, M., Tosatti, V., Zhang, Y.: Gromov–Hausdorff collapsing of Calabi–Yau manifolds. *Commun. Anal. Geom.* **24**(1), 93–113 (2016)
4. Hein, H.-J., Tosatti, V.: Remarks on the collapsing of torus fibered Calabi–Yau manifolds. *Bull. Lond. Math. Soc.* **47**(6), 1021–1027 (2015)
5. Tosatti, V.: Limits of Calabi–Yau metrics when the Kähler class degenerates. *J. Eur. Math. Soc. (JEMS)* **11**(4), 755–776 (2009)
6. Tosatti, V.: Adiabatic limits of Ricci-flat Kähler metrics. *J. Differ. Geom.* **84**(2), 427–453 (2010)
7. Tosatti, V., Zhang, Y.: Collapsing hyperkähler manifolds. *Ann. Sci. Éc. Norm. Supér.* **53**(3), 751–786 (2020)
8. Strominger, A., Yau, S.-T., Zaslow, E.: Mirror symmetry is T-duality. *Nucl. Phys. B* **479**(1–2), 243–259 (1996)
9. Vafa, C.: Extending mirror conjecture to Calabi–Yau with bundles. *Commun. Contemp. Math.* **1**, 65–70 (1999)
10. Fukaya, K.: Multivalued Morse theory, asymptotic analysis and mirror symmetry. In: *Proceedings of Symposium on Pure Mathematics*, vol. 73, pp. 205–278. American Mathematical Society, Providence, RI (2005)

11. Yau, S.-T.: On the Ricci curvature of compact Kähler manifolds and the complex Monge–Ampère equation. I. Commun. Pure Appl. Math. **31**(3), 339–411 (1978)
12. Donaldson, S.K.: Anti self-dual Yang–Mills connections over complex algebraic surfaces and stable vector bundles. Proc. Lond. Math. Soc. **50**, 1–26 (1985)
13. Uhlenbeck, K., Yau, S.-T.: On the existence of Hermitian–Yang–Mills connections in stable vector bundles. Commun. Pure Appl. Math. **39–S**, 257–293 (1986)
14. Dostoglou, S., Salamon, D.: Self-dual instantons and holomorphic curves. Ann. Math. **139**(3), 581–640 (1994)
15. Dostoglou, S., Salamon, D.: Corrigendum: "Self-dual instantons and holomorphic curves". Ann. Math. **165**(2), 665–673 (2007)
16. Friedman, R., Morgan, J., Witten, E.: Vector bundles over elliptic fibrations. J. Algebraic Geom. **8**(2), 279–401 (1999)
17. Chen, J.: Convergence of anti-self-dual connections on $SU(n)$ -bundles over product of two Riemann surfaces. Commun. Math. Phys. **196**(3), 571–590 (1998)
18. Fukaya, K.: Anti-self-dual equation on 4-manifolds with degenerate metric. Geom. Funct. Anal. **8**(3), 466–528 (1998)
19. Nishinou, T.: Convergence of adiabatic family of anti-self-dual connections on products of Riemann surfaces. J. Math. Phys. **51**(2), 022306 (2010). (10 pp)
20. Nishinou, T.: Global gauge fixing for connections with small curvature on T^2 . Int. J. Math. **18**(2), 165–177 (2007)
21. Nishinou, T.: Convergence of Hermitian–Yang–Mills Connections on Kähler Surfaces and mirror symmetry. [arXiv:math/0301324](https://arxiv.org/abs/math/0301324) [math.SG]
22. Fu, J.: Limiting behavior of a class of Hermitian–Yang–Mills metrics. Sci. China Math. **62**, 2155–2194 (2019)
23. Collins, T.C., Jacob, A., Yau, S.-T.: Poisson metrics on flat vector bundles over non-compact curves. Commun. Anal. Geom. **27**(3), 529–597 (2019)
24. Simpson, C.: Harmonic bundles on noncompact curves. J. Am. Math. Soc. **3**(3), 713–770 (1990)
25. Simpson, C.: Higgs bundles and local systems. Inst. Hautes Études Sci. Publ. Math. **75**, 5–95 (1992)
26. Datar, V., Jacob, A., Zhang, Y.: Adiabatic limits of anti-self-dual connections on collapsed K3 surfaces. J. Differ. Geom. **118**(2), 223–296 (2021)
27. Freed, D.: Special Kähler manifolds. Commun. Math. Phys. **203**(1), 31–52 (1999)
28. Hitchin, N.: Monopoles, Minimal Surfaces and Algebraic Curves. Séminaire de Mathématiques Supérieures, vol. 105, Les Presses de L'Université de Montréal (1987)
29. Friedman, R.: Algebraic Surfaces and Holomorphic Vector Bundles. Universitext, Springer, New York (1998)
30. Friedman, R., Morgan, J., Witten, E.: Vector bundles and F theory. Commun. Math. Phys. **187**(3), 679–743 (1997)
31. Morgan, J.: Holomorphic bundles over elliptic manifolds. School on Algebraic Geometry (Trieste, 1999), pp. 135–203, ICTP Lecture Notes, 1, Abdus Salam International Center for Theoretical Physics, Trieste (2000)
32. Donaldson, S.K., Kronheimer, P.B.: The Goemetry of Four-Manifolds. Oxford University Press, Oxford (1990)
33. Kobayashi, S.: Differential geometry of complex vector bundles. Publications of the Mathematical Society of Japan, 15. Kanô Memorial Lectures, 5. Princeton University Press, Princeton, NJ (1987)
34. Atiyah, M.F.: Vector bundles over an elliptic curve. Proc. Lond. Math. Soc. **7**, 414–452 (1957)
35. Uhlenbeck, K.: Removable singularities in Yang–Mills fields. Commun. Math. Phys. **83**, 11–29 (1982)
36. Tosatti, V., Zhang, Y.: Infinite time singularities of the Kähler–Ricci flow. Geom. Topol. **19**(5), 2925–2948 (2015)
37. Uhlenbeck, K.: Connections with L^p bounds on curvature. Commun. Math. Phys. **83**, 31–42 (1982)
38. Wehrheim, K.: Energy identity for anti-self-dual instantons on $\mathbb{C} \times \Sigma$. Math. Res. Lett. **13**(1), 161–166 (2006)
39. Jacob, A., Walpuski, T.: Hermitian Yang–Mills metrics on reflexive sheaves over asymptotically cylindrical Kähler manifolds. Commun. Partial Differ. Equ. **43**(11), 1566–1598 (2018)
40. Rade, J.: On the Yang–Mills heat equation in two and three dimensions. J. Reine Angew. Math. **431**, 123–163 (1992)
41. Trautwein, S.: A survey of the GIT picture for the Yang–Mills equation over Riemann surfaces. Enseign. Math. **63**, 63–153 (2017)

42. Georgoulas, V., Robbin, J., Salamon, D.: The moment-weight inequality and the Hilbert-Mumford criterion. (2013) [arXiv:1311.0410](#) [math.SG]
43. Feehan, P.: Morse theory for the Yang-Mills energy function near flat connections. [arXiv:1906.03954](#) [math.DG]

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.