



Geometry & Topology

Volume 26 (2022)

**Surface group representations in $SL_2(\mathbb{C})$
with finite mapping class orbits**

INDRANIL BISWAS

SUBHOJOY GUPTA

MAHAN MJ

JUNHO PETER WHANG



Surface group representations in $SL_2(\mathbb{C})$ with finite mapping class orbits

INDRANIL BISWAS
SUBHOJOY GUPTA
MAHAN MJ
JUNHO PETER WHANG

Given an oriented surface of positive genus with finitely many punctures, we classify the finite orbits of the mapping class group action on the moduli space of semisimple complex special linear two dimensional representations of the fundamental group of the surface. For surfaces of genus at least two, such orbits correspond to homomorphisms with finite image. For genus one, they correspond to the finite or special dihedral representations. We also obtain an analogous result for bounded orbits in the moduli space.

57M50; 57M05, 20E36, 20F29

1. Introduction	679
2. Background	683
3. Non-Zariski-dense representations	690
4. Analysis of Dehn twists	696
5. Proof of the main results	701
6. Applications	712
Appendix Case of a once-punctured torus	713
References	717

1 Introduction

Let Σ be an oriented surface of genus $g \geq 0$ with a finite set \mathcal{F} of punctures. The $SL_2(\mathbb{C})$ -character variety of Σ

$$X(\Sigma) = \text{Hom}(\pi_1(\Sigma), SL_2(\mathbb{C})) // SL_2(\mathbb{C})$$

is an affine algebraic variety whose complex points parametrize the conjugacy classes of semisimple representations $\pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ of the fundamental group of Σ . Let

$\text{Mod}(\Sigma)$ denote the pure mapping class group of Σ fixing \mathcal{F} pointwise. The group acts on the moduli space $X(\Sigma)$ by precomposition. This paper classifies the finite orbits of this action for surfaces of positive genus.

Our analysis divides into the cases of genus one and higher. For surfaces of genus at least two, we prove the following.

Theorem A *Let Σ be an oriented surface of genus $g \geq 2$ with $n \geq 0$ punctures. A semisimple representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ has finite mapping class group orbit in the character variety $X(\Sigma)$ if and only if ρ is finite.*

To describe the corresponding result for surfaces of genus 1, define an irreducible representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ to be *special dihedral* if

- its image lies in the infinite dihedral group D_∞ in $\text{SL}_2(\mathbb{C})$ (its definition is recalled in [Section 2](#)), and
- there is a nonseparating simple closed curve a in Σ such that the restriction of ρ to the complement $\Sigma \setminus a$ is diagonal.

Theorem B *Let Σ be an oriented surface of genus 1 with $n \geq 0$ punctures. A semisimple representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ has finite mapping class group orbit in $X(\Sigma)$ if and only if ρ is finite or special dihedral up to conjugacy.*

Many of the technical issues in the proofs of [Theorems A](#) and [B](#) above arise while dealing with punctures. For Σ closed (ie $n = 0$) the proofs become considerably simplified thanks to existing results, in particular results of Cooper and Manning [\[5\]](#), Gallo, Kapovich and Marden [\[11\]](#) and Previte and Xia [\[21\]](#). To illustrate this, we have included a short proof of [Theorem A](#) in [Section 5](#) for closed surfaces of genus at least 2 using [\[11\]](#) and [\[21\]](#).

For surfaces of genus zero with more than three punctures, the description of the finite mapping class group orbits in the $\text{SL}_2(\mathbb{C})$ -character variety is more complicated and in general unknown. Lisovsky and Tykhyy [\[18\]](#) completed the case of the four-punctured sphere as part of their classification of algebraic solutions to Painlevé VI differential equations, after earlier works including Dubrovin and Mazzocco [\[9\]](#), Iwasaki [\[15\]](#), Hitchin [\[13\]](#), Boalch [\[2\]](#) and Cantat and Loray [\[4\]](#). In the case of the four-punctured sphere, it is known that there exist finite mapping class group orbits corresponding to representations with Zariski dense image in $\text{SL}_2(\mathbb{C})$. In particular, the simplicity of

results in Theorems A and B contrasts with the complexity of finite orbits in genus zero. We remark that the once-punctured torus case of Theorem B was essentially proved by Dubrovin and Mazzocco [9] (also in connection with Painlevé VI); the derivation from [9] is recorded in the appendix. Our work gives a different proof in this case.

The relationship between algebraic solutions of Painlevé VI and finite mapping class group orbits in the character variety of the four-punctured sphere comes from R Fuchs' interpretation of Painlevé VI [10] as the equation for isomonodromic deformations of Fuchsian systems on the Riemann sphere with 4 singular points. More generally, as shown in the work of Cousin [6] and Cousin and Heu [7], the finite mapping class group orbits in the character variety $X(\Sigma)$ correspond to algebraic isomonodromic deformations of algebraic SL_2 -bundles with logarithmic connection on algebraic curves. Combined with their result, the above results give a complete classification of algebraic solutions to isomonodromy equations over curves of positive genus, in the case of semisimple SL_2 -monodromy, in terms of monodromy data.

This paper pursues the theme of characterizing points on the character variety $X(\Sigma)$ with special dynamical properties. In this spirit, we also prove the result below. Given a complex algebraic variety V , we shall say that a subset of $V(\mathbb{C})$ is *bounded* if it has compact closure in $V(\mathbb{C})$ with respect to the Euclidean topology.

Theorem C *Let Σ be an oriented surface of genus $g \geq 1$ with $n \geq 0$ punctures. A semisimple representation $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ has bounded mapping class group orbit in the character variety $X(\Sigma)$ if and only if*

- (a) ρ is unitary up to conjugacy, or
- (b) $g = 1$ and ρ is special dihedral up to conjugacy.

Our results and methods answer some previously raised basic questions. Theorems A and B imply (Corollary 6.1) that a faithful representation of a positive-genus hyperbolic surface group into $SL_2(\mathbb{C})$ (or $PSL_2(\mathbb{C})$) cannot have finite mapping class group orbit in the character variety, answering a question raised by Lubotzky. Theorems A and B also verify the $G = SL_2(\mathbb{C})$ case of the following conjecture of Kisin [26, Chapter 1]: For π the fundamental group of a closed surface or a free group of rank $r \geq 3$, the points with finite orbits for the action of the outer automorphism group $\text{Out}(\pi)$ on the character variety $\text{Hom}(\pi, G) // G$, for reductive algebraic groups G , correspond to representations $\pi \rightarrow G$ with virtually solvable image. However, there are counterexamples to this conjecture for general G ; see for instance Biswas, Koberda,

Mj and Santharoubane [1] and Koberda and Santharoubane [16]. Finally, we show that, given a closed hyperbolic surface S of genus at least two, the energy of the harmonic map associated to a representation $\pi_1(S) \rightarrow \mathrm{SL}_2(\mathbb{C})$ is bounded along the mapping class group orbit in the character variety if and only if the representation fixes a point of \mathbb{H}^3 (Theorem 6.2). This answers a question due to Goldman.

For a surface Σ of negative Euler characteristic with $n \geq 1$ marked punctures, the subvarieties $X_k(\Sigma)$ of $X(\Sigma)$ obtained by fixing the traces $k = (k_1, \dots, k_n) \in \mathbb{C}^n$ of local monodromy along the punctures form a family of log Calabi–Yau varieties with rich Diophantine structure; see Whang [30; 31]. Classifying the finite mapping class group orbits (and other invariant subvarieties) forms an important step in the study of strong approximation for these varieties, undertaken in the once-punctured torus case by Bourgain, Gamburd and Sarnak [3].

Finally, we note that Theorems A, B and C remain valid if the pure mapping class group is replaced by the full mapping class group (allowing for permutation of punctures). Similarly, since the fundamental groups of punctured surfaces are free, Theorems A and C imply the following. Given any finitely generated group π , let $X(\pi, \mathrm{SL}_2)$ be its SL_2 –character variety; see Section 2.2.

Corollary *Let $\pi = F_m$ be a free group of rank $m \geq 4$. A semisimple representation $\rho: \pi \rightarrow \mathrm{SL}_2(\mathbb{C})$ has finite (resp. bounded) $\mathrm{Out}(\pi)$ –orbit in $X(\pi, \mathrm{SL}_2)$ if and only if ρ has finite (resp. bounded) image.*

Organization of the paper In Section 2, we record background on algebraic subgroups of $\mathrm{SL}_2(\mathbb{C})$, character varieties, and mapping class groups. We also introduce the notion of loop configurations as a tool to keep track of subsurfaces of a given surface. In Section 3, we study representations of surface groups whose images are contained in proper algebraic subgroups of $\mathrm{SL}_2(\mathbb{C})$, and give a characterization of those with finite mapping class group orbits for surfaces of positive genus.

In Sections 4 and 5, we prove our main results. One of the ingredients in the proof of Theorems A and B is a theorem of Patel–Shankar–Whang [20, Theorem 1.2], which states that a semisimple $\mathrm{SL}_2(\mathbb{C})$ –representation of a positive-genus surface group with finite monodromy along every simple loop must in fact be finite. (For the proof of Theorem C which runs in parallel, there is an analogous result.) Along essential curves, the requisite finiteness of monodromy can be largely obtained by studying Dehn twists, as described in Section 4. Finiteness of local monodromy along the punctures is more

involved, and is achieved in [Section 5](#). In the special case where Σ is a closed surface, we also give a different (short) proof of [Theorem A](#) relying on some of the techniques developed by Gallo, Kapovich and Marden [[11](#)] and Previte and Xia [[21](#)].

In [Section 6](#), we provide applications of our work, answering earlier mentioned questions due to Lubotzky and Goldman. Finally, in the [appendix](#) we demonstrate a derivation of the once-punctured torus case of [Theorem B](#) from Dubrovin and Mazzocco [[9](#)].

2 Background

2.1 Subgroups of $SL_2(\mathbb{C})$

Let G be a proper algebraic subgroup of SL_2 defined over \mathbb{C} . Up to conjugation, G satisfies one of the following [[23](#), Theorem 4.29]:

- (1) G is a subgroup of the *standard Borel group*

$$B = \left\{ \begin{bmatrix} a & b \\ 0 & a^{-1} \end{bmatrix} \mid a \in \mathbb{C}^\times, b \in \mathbb{C} \right\}.$$

- (2) G is a subgroup of the *infinite dihedral group*

$$D_\infty = \left\{ \begin{bmatrix} c & 0 \\ 0 & c^{-1} \end{bmatrix} \mid c \in \mathbb{C}^\times \right\} \cup \left\{ \begin{bmatrix} 0 & c \\ -c^{-1} & 0 \end{bmatrix} \mid c \in \mathbb{C}^\times \right\}.$$

- (3) G is one of the finite groups BA_4 , BS_4 or BA_5 , which are the preimages in $SL_2(\mathbb{C})$ of the finite subgroups A_4 (tetrahedral group), S_4 (octahedral group) and A_5 (icosahedral group) of $PGL_2(\mathbb{C})$, respectively.

We refer to the [appendix](#) (after the statement of [Theorem A.3](#)) for an explicit description of the finite groups BA_4 , BS_4 and BA_5 .

Definition 2.1 A representation $\pi \rightarrow SL_2(\mathbb{C})$ of a group π is

- (1) *Zariski dense* if its image is not contained in a proper algebraic subgroup of SL_2 defined over \mathbb{C} ,
- (2) *diagonal* if it factors through the inclusion $i: \mathbb{C}^\times \rightarrow SL_2(\mathbb{C})$ of the maximal torus consisting of diagonal matrices,
- (3) *dihedral* if it factors through the inclusion $j: D_\infty \rightarrow SL_2(\mathbb{C})$,

- (4) *finite* if its image is finite,
- (5) *unitarizable* if its image is conjugate to a subgroup of $SU(2)$,
- (6) *reducible* if its image preserves a subspace of \mathbb{C}^2 ,
- (7) *irreducible* if it is not reducible,
- (8) *elementary* if it is unitary or reducible or dihedral,
- (9) *nonelementary* if it is not elementary, and
- (10) *semisimple* if \mathbb{C}^2 is a direct sum of π -modules under the representation (or, equivalently, the representation is irreducible or conjugate to a diagonal representation).

2.2 Character varieties

Given a finitely presented group π , let us define the SL_2 -representation variety $\text{Rep}(\pi)$ as the complex affine scheme determined by the functor

$$A \mapsto \text{Hom}(\pi, SL_2(A))$$

for every commutative \mathbb{C} -algebra A . Given a sequence of generators of π with m elements, we have a presentation of $\text{Rep}(\pi)$ as a closed subscheme of SL_2^m defined by equations coming from relations among the generators. For each $a \in \pi$, let tr_a be the regular function on $\text{Rep}(\pi)$ given by $\rho \mapsto \text{tr} \rho(a)$. The *character variety* of π over \mathbb{C} is the affine invariant theoretic quotient

$$X(\pi) = \text{Rep}(\pi) // SL_2 = \text{Spec } \mathbb{C}[\text{Rep}(\pi)]^{SL_2(\mathbb{C})}$$

under the conjugation action of SL_2 . The complex points of $X(\pi)$ parametrize the isomorphism classes of semisimple representations $\pi \rightarrow SL_2(\mathbb{C})$, or equivalently the Jordan equivalence classes of representations $\pi \rightarrow SL_2(\mathbb{C})$; see eg [25, Proposition 6.1]. For each $a \in \pi$ the regular function tr_a evidently descends to a regular function on $X(\pi)$. The scheme $X(\pi)$ has a natural model over \mathbb{Z} . We refer to [14; 22; 24] for details.

Example 2.2 We refer to Goldman [12] for details of the examples below. Let F_m denote the free group on $m \geq 1$ generators a_1, \dots, a_m .

- (1) We have $\text{tr}_{a_1}: X(F_1) \simeq \mathbb{A}^1$.
- (2) We have $(\text{tr}_{a_1}, \text{tr}_{a_2}, \text{tr}_{a_1 a_2}): X(F_2) \simeq \mathbb{A}^3$ by Fricke [12, Section 2.2].

(3) The coordinate ring $\mathbb{Z}[X(F_3)]$ is the quotient of the polynomial ring

$$\mathbb{Z}[\text{tr}_{a_1}, \text{tr}_{a_2}, \text{tr}_{a_3}, \text{tr}_{a_1 a_2}, \text{tr}_{a_2 a_3}, \text{tr}_{a_1 a_3}, \text{tr}_{a_1 a_2 a_3}, \text{tr}_{a_1 a_3 a_2}]$$

by the ideal generated by two elements

$$\begin{aligned} &\text{tr}_{a_1 a_2 a_3} + \text{tr}_{a_1 a_3 a_2} - (\text{tr}_{a_1 a_2} \text{tr}_{a_3} + \text{tr}_{a_1 a_3} \text{tr}_{a_2} + \text{tr}_{a_2 a_3} \text{tr}_{a_1} - \text{tr}_{a_1} \text{tr}_{a_2} \text{tr}_{a_3}), \\ &\text{tr}_{a_1 a_2 a_3} \text{tr}_{a_1 a_3 a_2} - \{(\text{tr}_{a_1}^2 + \text{tr}_{a_2}^2 + \text{tr}_{a_3}^2) + (\text{tr}_{a_1 a_2}^2 + \text{tr}_{a_2 a_3}^2 + \text{tr}_{a_1 a_3}^2) \\ &\quad - (\text{tr}_{a_1} \text{tr}_{a_2} \text{tr}_{a_1 a_2} + \text{tr}_{a_2} \text{tr}_{a_3} \text{tr}_{a_2 a_3} + \text{tr}_{a_1} \text{tr}_{a_3} \text{tr}_{a_1 a_3}) \\ &\quad + \text{tr}_{a_1 a_2} \text{tr}_{a_2 a_3} \text{tr}_{a_1 a_3} - 4\}. \end{aligned}$$

In particular, $\text{tr}_{a_1 a_2 a_3}$ and $\text{tr}_{a_1 a_3 a_2}$ are integral over the polynomial subring $\mathbb{Z}[\text{tr}_{a_1}, \text{tr}_{a_2}, \text{tr}_{a_3}, \text{tr}_{a_1 a_2}, \text{tr}_{a_2 a_3}, \text{tr}_{a_1 a_3}]$.

We record the following, which is attributed by Goldman [12] to Vogt [28].

Lemma 2.3 *Given a finitely generated group π and $a_1, a_2, a_3, a_4 \in \pi$, we have*

$$\begin{aligned} 2\text{tr}_{a_1 a_2 a_3 a_4} &= \text{tr}_{a_1} \text{tr}_{a_2} \text{tr}_{a_3} \text{tr}_{a_4} + \text{tr}_{a_1} \text{tr}_{a_2 a_3 a_4} + \text{tr}_{a_2} \text{tr}_{a_3 a_4 a_1} + \text{tr}_{a_3} \text{tr}_{a_4 a_1 a_2} \\ &\quad + \text{tr}_{a_4} \text{tr}_{a_1 a_2 a_3} + \text{tr}_{a_1 a_2} \text{tr}_{a_3 a_4} + \text{tr}_{a_4 a_1} \text{tr}_{a_2 a_3} - \text{tr}_{a_1 a_3} \text{tr}_{a_2 a_4} \\ &\quad - \text{tr}_{a_1} \text{tr}_{a_2} \text{tr}_{a_3 a_4} - \text{tr}_{a_3} \text{tr}_{a_4} \text{tr}_{a_1 a_2} - \text{tr}_{a_4} \text{tr}_{a_1} \text{tr}_{a_2 a_3} - \text{tr}_{a_2} \text{tr}_{a_3} \text{tr}_{a_4 a_1}. \end{aligned}$$

The above computations imply the following fact.

Fact 2.4 *If π is a group generated by a_1, \dots, a_m , then $\mathbb{Q}[X(\pi)]$ is generated as a \mathbb{Q} -algebra by the collection $\{\text{tr}_{a_{i_1} \dots a_{i_k}} \mid 1 \leq i_1 < \dots < i_k \leq m\}_{1 \leq k \leq 3}$.*

Remark Using Fact 2.4, we will show (Lemma 2.7) that an $SL_2(\mathbb{C})$ -representation ρ of a surface group has finite mapping class group orbit in the SL_2 -character variety if and only if the set of its traces along simple closed curves is finite.

The construction of $X(\pi)$ is functorial with respect to the group π . Given a homomorphism $f: \pi \rightarrow \pi'$ of finitely presented groups, the corresponding morphism $f^*: X(\pi') \rightarrow X(\pi)$ sends a representation ρ to the semisimplification of $\rho \circ f$. In particular, the automorphism group $\text{Aut}(\pi)$ of π naturally acts on $X(\pi)$. This action naturally factors through the outer automorphism group $\text{Out}(\pi)$, owing to the fact that $\text{tr}_{aba^{-1}} = \text{tr}_b$ for every $a, b \in \pi$.

Let $i : \mathbb{C}^\times \rightarrow \mathrm{SL}_2(\mathbb{C})$ and $j : D_\infty \rightarrow \mathrm{SL}_2(\mathbb{C})$ be the inclusion maps of the diagonal maximal torus and the infinite dihedral group, respectively. They induce $\mathrm{Out}(\pi)$ -equivariant maps

$$(*) \quad i_* : \mathrm{Hom}(\pi, \mathbb{C}^\times) \rightarrow X(\pi)(\mathbb{C}) \quad \text{and} \quad j_* : \mathrm{Hom}(\pi, D_\infty)/D_\infty \rightarrow X(\pi)(\mathbb{C}).$$

Lemma 2.5 *The following two statements hold:*

- (1) *The map $i_* : \mathrm{Hom}(\pi, \mathbb{C}^\times) \rightarrow X(\pi)(\mathbb{C})$ in (*) has finite fibers.*
- (2) *The map $j_* : \mathrm{Hom}(\pi, D_\infty)/D_\infty \rightarrow X(\pi)(\mathbb{C})$ in (*) has finite fibers.*

Proof (1) Let $\rho_1, \rho_2 : \pi \rightarrow \mathbb{C}^\times$ be characters such that $g i_*(\rho_1) g^{-1} = i_*(\rho_2)$ for some $g \in \mathrm{SL}_2(\mathbb{C})$. Without loss of generality, we may assume that $\rho_1(x) \neq \pm 1$ for some $x \in \pi$, since otherwise the image of ρ_1 is finite and we are done. Writing $g = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and $\rho_1(x) = \lambda$ with $\lambda \in \mathbb{C}^\times \setminus \{\pm 1\}$, we have

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \begin{bmatrix} \lambda ad - \lambda^{-1} bc & (\lambda^{-1} - \lambda) ab \\ (\lambda - \lambda^{-1}) cd & \lambda^{-1} ad - \lambda bc \end{bmatrix}.$$

For the matrix on the right-hand side to be diagonal, we must thus have $a = d = 0$ or $b = c = 0$ since $\lambda \neq \pm 1$. If $a = d = 0$, then $\rho_2 = \rho_1^{-1}$. If $b = c = 0$, then $\rho_1 = \rho_2$. This proves (1).

(2) Let $\rho_1, \rho_2 : \pi \rightarrow D_\infty$ be representations such that $g j_*(\rho_1) g^{-1} = j_*(\rho_2)$ for some $g \in \mathrm{SL}_2(\mathbb{C})$. Consider first the case where $\mathrm{tr} \rho_1(x) \in \{0, \pm 2\}$ for every $x \in \pi$. In this case, up to conjugation in D_∞ the image of ρ_1 must lie in the finite group

$$G = \left\langle \left[\begin{array}{cc} 0 & i^k \\ -i^{-k} & 0 \end{array} \right], \left[\begin{array}{cc} i^k & 0 \\ 0 & i^{-k} \end{array} \right] \mid k \in \mathbb{Z} \right\rangle.$$

To see this, note first that clearly the diagonal elements in the image of ρ_1 belong to G . Next, suppose we have (and fix) some $x \in \pi$ such that $\rho_1(x)$ is not diagonal. Up to conjugating ρ_1 by an element of D_∞ we may assume

$$\rho_1(x) = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

It follows that any other $y \in \pi$ with nondiagonal $\rho_1(y)$ must be in G , since otherwise $\mathrm{tr} \rho(xy) \notin \{0, \pm 2\}$. Thus, we conclude that if $\mathrm{tr} \rho_1(x) \in \{0, \pm 2\}$ for every $x \in \pi$ (and the same holds for ρ_2 since ρ_1 and ρ_2 are assumed to be $\mathrm{SL}_2(\mathbb{C})$ -conjugate), then ρ_1 and ρ_2 are each D_∞ -conjugate to a representation with image in the finite group G . Since π is finitely generated, there are only finitely many homomorphisms $\pi \rightarrow G$, and from this our claim follows.

It remains to treat the case where $\text{tr } \rho_1(x) \notin \{0, \pm 2\}$ for some $x \in \pi$. Now note that $\rho_1(x)$ must be diagonal. The equation $g\rho_1(x)g^{-1} = \rho_2(x)$ shows that, by the same computation as in the proof of part (1), we have $g \in D_\infty$. This proves (2). \square

2.3 Surfaces

Here, we set our notational conventions and terminology for various topological notions. Throughout this paper, a *surface* is the complement of a finite collection of interior points in a compact oriented topological manifold of dimension 2, with or without boundary.

A *simple closed curve* on a surface is an embedded copy of an unoriented circle. We shall often refer to simple closed curves simply as curves (since immersed curves will not be important in this paper). Given a surface Σ , a curve in Σ is *nondegenerate* if it does not bound a disk on Σ . A curve in the interior of Σ is *essential* if it is nondegenerate, does not bound a once-punctured disk on Σ , and is not isotopic to a boundary curve of Σ . Given a surface Σ and an essential curve $a \subset \Sigma$, we denote by $\Sigma|a$ the surface obtained by cutting Σ along a . An essential curve $a \subset \Sigma$ is *separating* if the two boundary curves of $\Sigma|a$ corresponding to a are in different connected components, and *nonseparating* otherwise.

Let Σ be a surface of genus g with n punctures or boundary curves. We shall denote by $\text{Mod}(\Sigma)$ the (pure) mapping class group of Σ . By definition, it is the group of isotopy classes of orientation-preserving homeomorphisms of Σ fixing the punctures and boundary points individually. Given a simple closed curve $a \subset \Sigma$, we shall denote by $\text{tw}_a \in \text{Mod}(\Sigma)$ the associated (left) Dehn twist.

We define the character variety of Σ (cf [Section 2.2](#)) to be

$$X(\Sigma) = X(\pi_1(\Sigma)).$$

The complex points of $X(\Sigma)$ can be seen as parametrizing the isomorphism classes of semisimple $SL_2(\mathbb{C})$ -local systems on Σ . Note that a simple closed curve $a \subset \Sigma$ unambiguously defines a function tr_a on $X(\Sigma)$, coinciding with tr_α for any loop $\alpha \in \pi_1(\Sigma)$ freely homotopic to a parametrization of a .

Given a continuous map $f: \Sigma' \rightarrow \Sigma$ of surfaces, we have an induced morphism of character varieties $f^*: X(\Sigma) \rightarrow X(\Sigma')$ depending only on the homotopy class of f . In particular, the mapping class group $\text{Mod}(\Sigma)$ acts naturally on $X(\Sigma)$ by precomposition: given $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ and $\gamma \in \text{Mod}(\Sigma)$, the class of ρ in $X(\Sigma)$ is mapped to the

class of $\rho \circ \gamma_*$, where γ_* is the outer automorphism of $\pi_1(\Sigma)$ determined by γ with any choice of basepoint. If $\Sigma' \subset \Sigma$ is a subsurface, the induced morphism on character varieties is $\text{Mod}(\Sigma')$ -equivariant for the induced morphism $\text{Mod}(\Sigma') \rightarrow \text{Mod}(\Sigma)$ of mapping class groups. In particular, if a semisimple $\text{SL}_2(\mathbb{C})$ -representation of $\pi_1(\Sigma)$ has a finite $\text{Mod}(\Sigma)$ -orbit in $X(\Sigma)$, then its restriction to any subsurface $\Sigma' \subset \Sigma$ has a finite $\text{Mod}(\Sigma')$ -orbit in $X(\Sigma')$.

2.4 Loop configurations

Let Σ be a surface of genus g with n punctures. We fix a basepoint in Σ . For convenience, we shall say that a sequence $\ell = (\ell_1, \dots, \ell_m)$ of based loops on Σ is *clean* if each loop is simple and the loops pairwise intersect only at the basepoint.

Example 2.6 Recall the standard presentation of the fundamental group

$$\pi_1(\Sigma) = \langle a_1, d_1, \dots, a_g, d_g, c_1, \dots, c_n \mid [a_1, d_1] \cdots [a_g, d_g] c_1 \cdots c_n \rangle.$$

We can choose (the based loops representing) the generators so that the sequence of loops $(a_1, d_1, \dots, a_g, d_g, c_1, \dots, c_n)$ is clean. For $i = 1, \dots, g$, let b_i be the based simple loop parametrizing the curve underlying d_i with the opposite orientation. Note that $(a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_n)$ is a clean sequence with the property that any product of distinct elements preserving the cyclic ordering on the sequence, such as $a_1 b_g$ or $a_1 a_2 b_2 b_g$ or $b_g c_n a_1$, can be represented by a simple loop in Σ . We shall refer to $(a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_n)$ as an *optimal sequence of generators* of $\pi_1(\Sigma)$. See Figure 1 for an illustration of the optimal generators for $(g, n) = (2, 1)$.

Lemma 2.7 Let $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ be a representation. Then ρ has finite $\text{Mod}(\Sigma)$ -orbit $X(\Sigma)$ if and only if

$$S = \{ \text{tr}_a(\rho) \mid a \subset \Sigma \text{ is a simple closed curve} \}$$

is a finite set.




Figure 1: Optimal generators for $(g, n) = (2, 1)$.

Proof Let (a_1, \dots, a_{2g+n}) be an optimal sequence of generators for $\pi_1(\Sigma)$. By [Fact 2.4](#), the coordinate ring $\mathbb{Q}[X(\Sigma)]$ of $X(\Sigma)$ is generated over \mathbb{Q} by the collection of trace functions

$$\{\text{tr}_{a_{i_1} \dots a_{i_k}} \mid 1 \leq i_1 < \dots < i_k \leq 2g+n\}_{1 \leq k \leq 2g+n},$$

giving us a closed immersion

$$j = (\text{tr}_{a_{i_1} \dots a_{i_k}}): X(\Sigma) \hookrightarrow \mathbb{A}^{2^{2g+n}-1}.$$

By our observation about optimal sequences of generators in [Example 2.6](#), every $a_{i_1} \dots a_{i_k}$ appearing above is homotopic to a simple loop. Since $\text{tr}_a(\gamma^* \rho) = \text{tr}_{\gamma(a)}(\rho)$ for every simple closed curve $a \subset \Sigma$ and $\gamma \in \text{Mod}(\Sigma)$, we see that if \mathcal{S} is finite then the coordinates (under the immersion j) of the points in the $\text{Mod}(\Sigma)$ -orbit of ρ belong to a finite set, and hence ρ has finite $\text{Mod}(\Sigma)$ -orbit in $X(\Sigma)$. Conversely, if ρ has finite $\text{Mod}(\Sigma)$ -orbit in $X(\Sigma)$, then $\{b_1, \dots, b_r\}$ is a collection of representatives of mapping class group equivalence classes of simple closed curves on Σ (considered up to isotopy), and \mathcal{S}' is the finite set of coordinates of the images of the points in the orbit under the morphism $(\text{tr}_{b_i}): X(\Sigma) \rightarrow \mathbb{A}^r$, then we have $\mathcal{S} = \mathcal{S}'$ which is therefore finite. □

Definition 2.8 A *loop configuration* is a planar graph consisting of a single vertex v and a finite cyclically ordered sequence of directed rays, equipped with a bijection between the set of rays departing from v and the set of rays arriving at v . We denote by $L_{g,n}$ the loop configuration whose sequence of rays is of the form

$$(a_1, b_1, \bar{a}_1, \bar{b}_1, \dots, a_g, b_g, \bar{a}_g, \bar{b}_g, c_1, \bar{c}_1, \dots, c_n, \bar{c}_n),$$

where a_i, b_i and c_i are the rays directed away from v , corresponding respectively to the rays \bar{a}_i, \bar{b}_i and \bar{c}_i directed towards v . An isomorphism of loop configurations is an isomorphism of planar graphs respecting the bijections between the departing and arriving rays. See [Figure 2](#) for an illustration of $L_{2,1}$.




Figure 2: Loop configuration $L_{2,1}$.

Given a clean sequence $\ell = (\ell_1, \dots, \ell_m)$ of loops on Σ , we have an associated loop configuration $L(\ell)$, obtained by taking a sufficiently small open neighborhood of the basepoint and setting the departing and arriving ends of the loops ℓ_i to correspond to each other. For example, if $(a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_n)$ is a sequence of optimal generators for $\pi_1(\Sigma)$, then

$$L(a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_n) \simeq L_{g,n}.$$

Definition 2.9 Let h and m be nonnegative integers. A sequence of based loops $\ell = (\ell_1, \dots, \ell_{2h+m})$ on Σ is said to be in $(h, m+1)$ -position if it is homotopic termwise to a clean sequence $\ell' = (\ell'_1, \dots, \ell'_{2h+m})$ such that $L(\ell') \simeq L_{h,m}$. We denote by $\Sigma(\ell) \subset \Sigma$ the (isotopy class of a) subsurface of genus h with $m+1$ boundary curves obtained by taking a small closed tubular neighborhood of the union of the simple curves underlying $\ell'_1, \dots, \ell'_{2h+m}$ in Σ .

Remark It is worth emphasizing that the terminology above is intended to evoke the topological type of the surface $\Sigma(\ell)$, and not the isomorphism type of the loop configuration.

3 Non-Zariski-dense representations

Let Σ be a surface of genus $g \geq 1$ with $n \geq 0$ punctures. The purpose of this section is to characterize representations $\pi_1(\Sigma)$ with non-Zariski-dense image in $\mathrm{SL}_2(\mathbb{C})$ that have finite mapping class group orbit in the character variety $X(\Sigma)$. An irreducible representation $\rho: \pi_1(\Sigma) \rightarrow \mathrm{SL}_2(\mathbb{C})$ will be called *special dihedral* if it factors through D_∞ and there is a nonseparating essential curve a in Σ such that the restriction $\rho|_{(\Sigma \setminus a)}$ is diagonal. The main result of this section is the following.

Proposition 3.1 *Let Σ be a surface of genus $g \geq 1$ with $n \geq 0$ punctures. Let $\rho: \pi_1(\Sigma) \rightarrow \mathrm{SL}_2(\mathbb{C})$ be a semisimple representation whose image is not Zariski-dense in $\mathrm{SL}_2(\mathbb{C})$. Then ρ has a finite mapping class group orbit in $X(\Sigma)$ if and only if one of the following holds:*

- (1) ρ is a finite representation.
- (2) $g = 1$ and ρ is special dihedral up to conjugation.

It is evident that, if the image of $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ belongs to one of the finite groups BA_4 , BS_4 or BA_5 , then its orbit in $X(\Sigma)$ is finite. From the discussion in Section 2.1 and Lemma 2.5, to prove Proposition 3.1 it remains to understand the finite mapping class group orbits on $\text{Hom}(\pi_1(\Sigma), \mathbb{C}^\times)$ and $\text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty$. Proposition 3.1 thus follows by combining Lemmas 3.2, 3.3 and 3.5 below.

Case 1 (diagonal representations)

Lemma 3.2 *Assume Σ is a surface of genus $g \geq 1$ with $n \geq 0$ punctures. A representation $\rho: \pi_1(\Sigma) \rightarrow \mathbb{C}^\times$ has finite (resp. bounded) mapping class group orbit in $\text{Hom}(\pi_1(\Sigma), \mathbb{C}^\times)$ if and only if it has finite (resp. bounded) image.*

Proof Let $\rho: \pi_1(\Sigma) \rightarrow \mathbb{C}^\times$ be a representation with finite (resp. bounded) mapping class group orbit in $\text{Hom}(\pi_1(\Sigma), \mathbb{C}^\times)$. Let $(a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_n)$ be the optimal generators of $\pi_1(\Sigma)$. Assume first that $n = 0$ or 1. By considering the effect of the Dehn twist along the curve underlying a_1 on the curve underlying b_1 , we conclude that $\rho(a_1)$ must be torsion (resp. have absolute value 1). Applying the same argument to the other loops in the sequence of optimal generators, we conclude that ρ is finite (resp. bounded) if $n \leq 1$, as desired.

Thus, only the case $n \geq 2$ remains. Since $(a_1, b_1, \dots, a_g, b_g)$ is in $(g, 1)$ -position, the above analysis shows that $\rho(a_i)$ and $\rho(b_i)$ are roots of unity (resp. have absolute value 1) for $i = 1, \dots, g$. Similarly, we see that the sequence

$$L_i = (a_1, b_1, \dots, a_{g-1}, b_{g-1}, a_g, b_g c_i)$$

is in $(g, 1)$ -position for every $i = 1, \dots, n$. Since ρ restricted to the surface $\Sigma(L_i)$ must have finite (resp. bounded) $\text{Mod}(\Sigma(L_i))$ -orbit, it follows that $\rho(c_i)$ is a root of unity (resp. has absolute value 1) for $i = 1, \dots, n$ as well. This shows that ρ has finite (resp. bounded) image, as desired. □

Case 2 (dihedral representations) We first prove Proposition 3.1 for surfaces of genus $g \geq 2$ in Lemma 3.3 below.

Lemma 3.3 *Assume Σ is a surface of genus $g \geq 2$ with $n \geq 0$ punctures. A representation $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ has finite mapping class group orbit in $\text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty$ if and only if it has finite image.*

Proof The “if” direction is clear, and we now prove the converse. Let $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ be a representation with finite mapping class group orbit in $\text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty$. We

have a short exact sequence $1 \rightarrow \mathbb{C}^\times \rightarrow D_\infty \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 0$, where the homomorphism $D_\infty \rightarrow \mathbb{Z}/2\mathbb{Z}$ is given by

$$\begin{bmatrix} c & 0 \\ 0 & c^{-1} \end{bmatrix} \mapsto 0 \quad \text{and} \quad \begin{bmatrix} 0 & c \\ -c^{-1} & 0 \end{bmatrix} \mapsto 1 \quad \text{for all } c \in \mathbb{C}^\times.$$

This gives us a $\text{Mod}(\Sigma)$ -equivariant map

$$\text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty \rightarrow \text{Hom}(\pi_1(\Sigma), \mathbb{Z}/2\mathbb{Z}).$$

The fiber of this map above the zero homomorphism consists of those points given by diagonal representations, to which [Lemma 3.2](#) applies, noting that the map

$$\text{Hom}(\pi_1(\Sigma), \mathbb{C}^\times) \rightarrow \text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty$$

has finite fibers. It suffices to consider the case where $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ is not in the fiber over the zero homomorphism. This assumption implies that there is a nonseparating loop a_1 on Σ such that $\rho(a_1)$ is not diagonal. Let us extend such a_1 to an optimal sequence of generators

$$(a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_n)$$

for $\pi_1(\Sigma)$; see [Section 2.4](#) for the definition. Note that $L = (a_1, b_1, \dots, a_g, b_g)$ is in $(g, 1)$ -position, and we have a $\text{Mod}(\Sigma(L))$ -equivariant homomorphism

$$\text{Hom}(\pi_1(\Sigma), \mathbb{Z}/2\mathbb{Z}) \rightarrow \text{Hom}(\pi_1(\Sigma(L)), \mathbb{Z}/2\mathbb{Z}).$$

The action of $\text{Mod}(\Sigma(L))$ on $\text{Hom}(\pi_1(\Sigma(L)), \mathbb{Z}/2\mathbb{Z})$ factors through the projection

$$\text{Mod}(\Sigma(L)) \twoheadrightarrow \text{Sp}(2g, \mathbb{Z}/2\mathbb{Z}).$$

From the transitivity of $\text{Sp}(2g, \mathbb{Z}/2\mathbb{Z})$ on $(\mathbb{Z}/2\mathbb{Z})^{2g}$ away from the origin, it follows that, up to $\text{Mod}(\Sigma)$ -action, we may assume that

$$\bar{\rho}(a_1) = 1 \quad \text{and} \quad \bar{\rho}(b_1) = \bar{\rho}(a_2) = \dots = \bar{\rho}(b_g) = 0,$$

where $\bar{\rho}$ is the image of ρ in $\text{Hom}(\pi_1(\Sigma), \mathbb{Z}/2\mathbb{Z})$. Up to conjugation by an element in D_∞ , we may moreover assume that

$$\rho(a_1) = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

It suffices to show that the entries of the matrices $\rho(b_1), \rho(a_2), \rho(b_2), \dots, \rho(a_g), \rho(b_g), \rho(c_1), \dots, \rho(c_n)$ are roots of unity. Let $i_1 < \dots < i_q$ be precisely the indices in $\{1, \dots, n\}$ such that $\bar{\rho}(c_{i_j}) = 0$. Since

$$L' = (a_2, b_2, \dots, a_g, b_g, c_{i_1}, \dots, c_{i_q}, b_1)$$

is in $(g-1, q+2)$ -position, and the restriction of ρ to $\Sigma(L')$ is diagonal, we see that $\rho(b_1), \rho(a_2), \rho(b_2), \dots, \rho(a_g), \rho(b_g), \rho(c_{i_1}), \dots, \rho(c_{i_q})$ are torsion by Lemma 3.2. Let us now take $i \in \{1, \dots, n\} \setminus \{i_1, \dots, i_q\}$. The restriction of ρ to the subsurface $\Sigma(c_i a_1, b_1)$ of genus 1 with one boundary curve is diagonal. Writing

$$\rho(c_i) = \begin{bmatrix} 0 & \lambda_i \\ -\lambda_i^{-1} & 0 \end{bmatrix}$$

with $\lambda_i \in \mathbb{C}^\times$, we have

$$\rho(c_i a_1) = \begin{bmatrix} 0 & \lambda_i \\ -\lambda_i^{-1} & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} -\lambda_i & 0 \\ 0 & -\lambda_i^{-1} \end{bmatrix}.$$

It follows from Lemma 3.2 that λ_i is a root of unity. This completes the proof that the entries of $\rho(b_1), \rho(a_2), \rho(b_2), \dots, \rho(a_g), \rho(b_g), \rho(c_1), \dots, \rho(c_n)$ are roots of unity, and hence the image of ρ is finite, as desired. \square

Before proving the case $g = 1$ of Proposition 3.1 in Lemma 3.5 below, we record the following elementary lemma.

Lemma 3.4 *Let Σ be a surface of genus 1 with $n \geq 0$ punctures. Suppose that $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ is special dihedral. Given a pair of loops (a, b) in $(1, 1)$ -position on Σ , at least one of $\rho(a)$ and $\rho(b)$ is not diagonal.*

Proof Let $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ be special dihedral. We argue by contradiction. Assume that (a_1, b_1) is a pair of loops in $(1, 1)$ -position on Σ with both $\rho(a_1)$ and $\rho(b_1)$ diagonal. We can complete (a_1, b_1) to a sequence of optimal generators $(a_1, b_1, c_1, \dots, c_n)$ of $\pi_1(\Sigma)$. Since ρ is special dihedral, the matrices $\rho(c_1), \dots, \rho(c_n)$ must be diagonal (noting that the property of a matrix being diagonal is not changed under conjugation by an element in D_∞). This implies that ρ is in fact diagonal, contradicting the hypothesis that ρ is irreducible. \square

Lemma 3.5 *Let Σ be a surface of genus 1 with $n \geq 0$ punctures. A representation $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ has finite mapping class group orbit in $\text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty$ if and only if it is finite or special dihedral.*

Proof The same argument as in Lemma 3.3 shows that if $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ has finite $\text{Mod}(\Sigma)$ -orbit in $\text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty$, then ρ is finite or special dihedral. Indeed, by Lemma 3.2 we are done if ρ is diagonal, so let us assume otherwise. This implies

there is a nonseparating loop a_1 such that $\rho(a_1)$ is not diagonal. We extend a_1 to an optimal sequence of generators for $\pi_1(\Sigma)$,

$$(a_1, b_1, c_1, \dots, c_n).$$

Without loss of generality, we may assume $\rho(b_1)$ is diagonal — otherwise, replace b_1 by a simple loop homotopic to $b_1 a_1$ and complete $(a_1, b_1 a_1)$ to an optimal sequence of generators for $\pi_1(\Sigma)$. Up to conjugation in D_∞ , we can also assume

$$\rho(a_1) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

If $\rho(c_1), \dots, \rho(c_n)$ are all diagonal, then ρ is special dihedral as desired. So assume otherwise; we need to show that ρ is finite. Suppose $\rho(c_i)$ is not diagonal, and say

$$\rho(c_i) = \begin{bmatrix} 0 & \lambda_i \\ -\lambda_i^{-1} & 0 \end{bmatrix}$$

for some $\lambda_i \in \mathbb{C}^\times$. Note that $(c_i a_1, b_1)$ is in $(1, 1)$ -position and the restriction of ρ to $\Sigma(c_i a_1, b_1)$ is diagonal, so λ_i as well as the eigenvalues of b_1 are roots of unity. This holds true for all c_i with nondiagonal $\rho(c_i)$. In what follows, fix one such c_i with $\rho(c_i)$ nondiagonal. For any c_j with $j < i$ such that $\rho(c_j)$ is diagonal, we see that $(c_i a_1, b_1 c_j)$ is in $(1, 1)$ -position and the restriction of ρ to $\Sigma(c_i a_1, b_1 c_j)$ is diagonal, so the eigenvalues of $\rho(c_j)$ are roots of unity. Similarly, for any c_j with $i < j$ such that $\rho(c_j)$ is diagonal, we see that $(c_i c_j a_1, b_1)$ is in $(1, 1)$ -position and the restriction of ρ to $\Sigma(c_i c_j a_1, b_1)$ is diagonal, so the eigenvalues of $\rho(c_j)$ are roots of unity. Combining these observations, it follows that ρ has finite image, as desired. Thus, we have prove the “only if” direction of the lemma.

Now let $\rho: \pi_1(\Sigma) \rightarrow D_\infty$ be a special dihedral representation. We shall show that ρ has finite mapping class group orbit in $\text{Hom}(\pi_1(\Sigma), D_\infty)/D_\infty$, or equivalently in the character variety $X(\Sigma)$. By [Lemma 2.7](#), it suffices to show that the set

$$\{\text{tr } \rho(a) \mid a \subset \Sigma \text{ essential curve}\} \subseteq \mathbb{C}$$

is finite.

Let $(a_1, b_1, c_1, \dots, c_n)$ be optimal generators of $\pi_1(\Sigma)$. Since ρ is special dihedral, $\rho(c_1), \dots, \rho(c_n)$ are diagonal matrices. Suppose a is a separating essential curve in Σ . Since one of the connected components of $\Sigma|_a$ is a surface of genus zero (with one boundary curve corresponding to a and finitely many punctures), the monodromy of ρ along a is a product of conjugates of a subset of $\{\rho(c_1), \dots, \rho(c_n)\}$. But since ρ is




Figure 3: Building new loops out of old ones.

dihedral, the conjugate of each $\rho(c_i)$ is $\rho(c_i)$ or $\rho(c_i)^{-1}$. It follows that $\text{tr } \rho(a)$ belongs to a finite set depending only on the boundary traces $\text{tr } \rho(c_i)$. Thus, we conclude that $\{\text{tr } \rho(a) \mid a \subset \Sigma \text{ is a separating curve}\}$ is finite.

It remains to show that $\{\text{tr } \rho(a) \mid a \subset \Sigma \text{ is a nonseparating curve}\}$ is finite. Let a_0 be a nonseparating curve in Σ such that the restriction $\rho|(\Sigma \setminus a_0)$ is diagonal; such a curve exists since ρ is special dihedral. Let b be a nonseparating curve in Σ . Up to isotopy, we may assume that b intersects a_0 only finitely many times. If b does not intersect a_0 , then $b \subset \Sigma \setminus a_0$ and hence $\text{tr } \rho(b)$ can take only finitely many values as $\rho|(\Sigma \setminus a_0)$ is diagonal.

If b intersects a_0 exactly once, then we must have $\text{tr } \rho(b) = 0$ by Lemma 3.4. Let us now assume that b intersects a_0 more than once. Let us choose a parametrization of b . Since b is nonseparating, there must be two neighboring points of intersection of a_0 and b where the two segments of b have the same orientation, as in Figure 3. The operations as in Figure 3 produce for us a new simple closed curve b' which is also nonseparating, as well as a pair (c, c') of simple loops in $(1, 1)$ -position on Σ . Note that b' as well as c and c' are constructed so that b is homotopic to cc' (here the implicit basepoint is the intersection point of c and c'), and b' is homotopic to $c'c^{-1}$. The fact that

$$\text{tr}(AB) + \text{tr}(AB^{-1}) = \text{tr}(A) \text{tr}(B)$$

for any $A, B \in SL_2(\mathbb{C})$ shows, upon choosing $A = \rho(c)$ and $B = \rho(c')$, that we have

$$\text{tr } \rho(b) = \text{tr } \rho(c) \text{tr } \rho(c') - \text{tr } \rho(b').$$

By Lemma 3.4, we have $\text{tr } \rho(c) \text{tr } \rho(c') = 0$, and therefore $\text{tr } \rho(b) = -\text{tr } \rho(b')$. Note furthermore that b' intersects a_0 in a smaller number of points than b does. Applying induction on the number of intersection points, we thus conclude that

$$\{\text{tr } \rho(a) \mid a \subset \Sigma \text{ is a nonseparating curve}\}$$

is finite. This completes the proof that special dihedral representations have finite mapping class group orbits in X . \square

4 Analysis of Dehn twists

Throughout this section, let Σ be a surface of genus $g \geq 1$ with $n \geq 0$ punctures. For convenience of exposition, we shall denote by (\dagger) the following condition on a representation $\pi_1(\Sigma) \rightarrow \mathrm{SL}_2(\mathbb{C})$:

- (\dagger) The representation is semisimple, and moreover Σ has genus at least 2 or the representation is not special dihedral.

Given $\rho: \pi_1(\Sigma) \rightarrow \mathrm{SL}_2(\mathbb{C})$ satisfying (\dagger) , Theorems A and B (resp. Theorem C) state that ρ has finite (resp. bounded) image in $\mathrm{SL}_2(\mathbb{C})$ if its mapping class group orbit in the character variety $X(\Sigma)$ is finite (resp. bounded). Let us recall the following.

Theorem 4.1 [20, Theorem 1.2] *Let Σ be a surface of positive genus $g \geq 1$ with $n \geq 0$ punctures. If a semisimple representation $\pi_1(\Sigma) \rightarrow \mathrm{SL}_2(\mathbb{C})$ has finite monodromy along all simple loops on Σ , then it has finite image.*

Proposition 4.2 [20, Lemma 2.2] *Let Σ be a surface of positive genus $g \geq 1$ with $n \geq 0$ punctures. If a semisimple representation $\pi_1(\Sigma) \rightarrow \mathrm{SL}_2(\mathbb{C})$ has elliptic or central monodromy along all simple loops on Σ , then it is unitarizable.*

Consequently, to prove Theorems A and B (resp. Theorem C) it suffices to prove that a representation $\rho: \pi_1(\Sigma) \rightarrow \mathrm{SL}_2(\mathbb{C})$ with finite (resp. bounded) mapping class group orbit in the character variety has finite (resp. elliptic or central) monodromy along all nondegenerate simple closed curves on Σ . Let us divide up the curves into four types:

- **Type I** Nonseparating essential curves.
- **Type II** Separating essential curves $a \subset \Sigma$ with each component of $\Sigma|a$ (defined in Section 2.3) having genus at least one.
- **Type III** Separating essential curves $a \subset \Sigma$ with one component of $\Sigma|a$ having genus zero.
- **Type IV** Boundary curves.

The purpose of this section is to show that, if $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ is a representation satisfying the hypotheses (\dagger) mentioned above with finite (resp. bounded) mapping class group orbit in $X(\Sigma)$, then ρ must have finite (resp. elliptic or central) monodromy along all curves of type I and II. We shall also show that ρ has finite (resp. elliptic or central) monodromy along all curves of type III provided the same holds for all curves of type IV. In particular, this is enough for us to prove the following special cases of Theorems A, B and C.

Proposition 4.3 *Let Σ be a surface of genus $g \geq 1$ with $n \geq 1$ punctures. Suppose that $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ is a semisimple representation with finite (resp. elliptic or central) local monodromy around the punctures, and finite (resp. bounded) mapping class group orbit in the character variety $X(\Sigma)$. Then one of the following holds:*

- (1) ρ is finite (resp. unitarizable).
- (2) $g = 1$ and ρ is special dihedral, up to conjugacy.

The rest of this section proves Proposition 4.3 by dealing with curves of types I, II and III. We shall complete the proof of our main results in Section 5 by treating the curves of type IV.

Type I (nonseparating essential curves)

Lemma 4.4 *Let $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ be a representation satisfying (\dagger) , and let $a \subset \Sigma$ be a curve of type I. If ρ has finite (resp. bounded) orbit under $\text{Mod}(\Sigma)$ in $X(\Sigma)$, then ρ has finite (resp. elliptic or central) monodromy along a .*

Proof Let us choose a basepoint x_0 on Σ lying on a , and let α be a simple based loop parametrizing a . Suppose β is another simple loop such that the pair (α, β) is in $(1, 1)$ -position, and let b denote the underlying curve. For each integer $k \in \mathbb{Z}$, the loop $\alpha^k \beta$ is homotopic to a simple loop whose underlying curve is isotopic to $\text{tw}_a^k(b)$. In particular, by our hypothesis the set $\{\text{tr } \rho(\alpha^k \beta) \mid k \in \mathbb{Z}\}$ is a finite (resp. bounded) subset of \mathbb{C} . Up to global conjugation of ρ by an element of $SL_2(\mathbb{C})$, we may consider two cases.

(a) Suppose first that

$$\rho(\alpha) = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix} \quad \text{with } \lambda \in \mathbb{C}^\times.$$

Let β be a simple loop on Σ such that (α, β) is in $(1, 1)$ -position, and write

$$\rho(\beta) = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}.$$

For each $k \in \mathbb{Z}$, we have

$$\text{tr } \rho(\alpha^k \beta) = \text{tr} \left(\begin{bmatrix} \lambda^k & 0 \\ 0 & \lambda^{-k} \end{bmatrix} \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \right) = \lambda^k b_1 + \lambda^{-k} b_4.$$

The fact that $\{\text{tr } \rho(\alpha^k \beta) \mid k \in \mathbb{Z}\}$ is a finite (resp. bounded) subset of \mathbb{C} then implies that λ is a root of unity (resp. has absolute value 1) so that $\rho(\alpha)$ is torsion (resp. elliptic or central), or that $b_1 = b_4 = 0$. Since the argument applies to any loop β such that (α, β) is in $(1, 1)$ -position, it remains only to consider the case where $\rho(\beta)$ has both diagonal entries zero for every such β . In this case, we see upon reflection that the restriction $\rho|(\Sigma|a)$ must be diagonal. This shows that ρ is special dihedral. Since ρ satisfies (\dagger) this means that Σ moreover has genus at least 2, so $\Sigma|a$ has genus at least 1 and it follows from [Lemma 3.2](#) that $\rho(\alpha)$ is torsion (resp. elliptic or central).

(b) Suppose that

$$\rho(\alpha) = s \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

for some $s \in \{\pm 1\}$ and $x \in \mathbb{C}$. Let us assume $s = +1$; the case $s = -1$ will follow similarly. Let β be a simple loop on Σ such that (α, β) is in $(1, 1)$ -position, and let us follow the notation for $\rho(\beta)$ from the previous case. For each $k \in \mathbb{Z}$, we have

$$\text{tr } \rho(\alpha^k \beta) = \text{tr} \left(\begin{bmatrix} 1 & kx \\ 0 & 1 \end{bmatrix} \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \right) = b_1 + b_4 + kx b_3.$$

The fact that $\{\text{tr } \rho(\alpha^k \beta) \mid k \in \mathbb{Z}\}$ is a finite (resp. bounded) subset of \mathbb{C} then implies that $x = 0$ or $b_3 = 0$. Since the argument applies to any loops β such that (α, β) is in $(1, 1)$ -position, it remains only to consider the case where $\rho(\beta)$ is upper triangular for every such β . In this case, ρ is upper triangular (hence diagonal). It follows from [Lemma 3.2](#) that $\rho(\alpha)$ is torsion (resp. elliptic or central).

The above arguments show that $\rho(\alpha)$ is torsion (resp. elliptic or central) unless $g = 1$ and ρ is special dihedral, as desired. □

Types II and III (separating essential curves)

Lemma 4.5 *Let $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ be a representation satisfying (\dagger) , and let $a \subset \Sigma$ be a curve of type II. If ρ has finite (resp. bounded) orbit under $\text{Mod}(\Sigma)$ in $X(\Sigma)$, then ρ has finite (resp. elliptic or central) monodromy along a .*

Lemma 4.6 *Let $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ be a representation satisfying (\dagger) , and let $a \subset \Sigma$ be a curve of type III. If ρ has finite (resp. bounded) orbit under $\text{Mod}(\Sigma)$ in $X(\Sigma)$, and moreover ρ has finite (resp. elliptic or central) monodromy along all punctures of Σ , then ρ has finite (resp. elliptic or central) monodromy along a .*

Proof of Lemmas 4.5 and 4.6 Let us choose a basepoint x_0 on Σ lying on a , and let α be a simple based loop parametrizing a . Let Σ_1 and Σ_2 be the connected components of $\Sigma|a$, and lift the basepoint on Σ to Σ_1 and Σ_2 . Suppose $\beta \in \pi_1(\Sigma_1)$ and $\gamma \in \pi_1(\Sigma_2)$ are simple loops such that their product $\beta\gamma$ on Σ is homotopic to a simple loop (with underlying curve denoted by d , say) transversely intersecting a exactly twice. For convenience, in this paragraph we shall call such a pair (β, γ) *good*. The loop $\beta\alpha^k\gamma\alpha^{-k}$ for each $k \in \mathbb{Z}$ is freely homotopic to a simple loop whose underlying curve belongs to the orbit $\langle \text{tw}_a \rangle \cdot d$ (all curves considered up to isotopy). In particular, by our hypothesis the set $\{\text{tr } \rho(\beta\alpha^k\gamma\alpha^{-k}) \mid k \in \mathbb{Z}\}$ is a finite (resp. bounded) subset of \mathbb{C} . Up to global conjugation of ρ by an element of $SL_2(\mathbb{C})$, we may consider two cases.

(a) Suppose first that

$$\rho(\alpha) = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix} \quad \text{with } \lambda \in \mathbb{C}^\times.$$

Let (β, γ) be a good pair, and let us write

$$\rho(\beta) = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \quad \text{and} \quad \rho(\gamma) = \begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix}.$$

For each $k \in \mathbb{Z}$, we have

$$\begin{aligned} \text{tr } \rho(\beta\alpha^k\gamma\alpha^{-k}) &= \text{tr} \left(\begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \begin{bmatrix} \lambda^k & 0 \\ 0 & \lambda^{-k} \end{bmatrix} \begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} \begin{bmatrix} \lambda^{-k} & 0 \\ 0 & \lambda^k \end{bmatrix} \right) \\ &= b_1c_1 + \lambda^{-2k}b_2c_3 + \lambda^{2k}c_2b_3 + b_4c_4. \end{aligned}$$

The fact that $\{\text{tr } \rho(\beta\alpha^k\gamma\alpha^{-k}) \mid k \in \mathbb{Z}\}$ is a finite (resp. bounded) subset of \mathbb{C} then implies that λ is a root of unity (resp. has absolute value 1) so that $\rho(\alpha)$ is torsion (resp. elliptic or central), or that $b_2c_3 = c_2b_3 = 0$. If the former happens, then we are done.

Suppose that the latter happens, and that at least one of b_2, b_3, c_2, c_3 is nonzero. We shall assume $b_2 \neq 0$; the other cases will follow similarly. As $b_2c_3 = 0$, we must have $c_3 = 0$. Applying the same argument with γ replaced by any simple loop $\gamma' \in \Sigma_2$ such that (β, γ') is good, we are reduced to the case where $\rho|_{\Sigma_2}$ is upper triangular. If $\rho|_{\Sigma_1}$ is also upper triangular, then ρ must be upper triangular (whence diagonal),

and from [Lemma 3.2](#) $\rho(\alpha)$ is torsion (resp. elliptic or central). So suppose there is a simple loop $\beta' \in \pi_1(\Sigma)$ such that $\rho(\beta')$ is not upper triangular. By repeating the above argument with β replaced by β' , we are reduced to the case where $\rho|_{\Sigma_2}$ must be diagonal. If Σ_2 has genus at least 1, then from [Lemma 3.2](#) it follows that $\rho(\alpha)$ is torsion (resp. is elliptic or central). If Σ_2 has genus 0, then the fact that ρ has finite (resp. elliptic or central) local monodromy along the punctures implies that $\rho(\alpha)$ has finite (resp. elliptic or central) monodromy.

It remains to consider the case where $b_2 = b_3 = c_2 = c_3 = 0$. Running through all the good pairs (β, γ) and repeating the above argument, we are left with the case where ρ is diagonal; [Lemma 3.2](#) then implies that $\rho(\alpha)$ is torsion (resp. elliptic or central).

The second case:

(b) Suppose that

$$\rho(\alpha) = s \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

for some $s \in \{\pm 1\}$ and $x \in \mathbb{C}$. We assume $s = +1$; the case $s = -1$ will follow similarly. Let (β, γ) be a good pair, and let us follow the notation for $\rho(\beta)$ and $\rho(\gamma)$ from the previous case. For each $k \in \mathbb{Z}$, we have

$$\begin{aligned} & \text{tr } \rho(\beta \alpha^k \gamma \alpha^{-k}) \\ &= \text{tr} \left(\begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \begin{bmatrix} 1 & kx \\ 0 & 1 \end{bmatrix} \begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} \begin{bmatrix} 1 & -kx \\ 0 & 1 \end{bmatrix} \right) \\ &= b_1 c_1 + b_2 c_3 + b_3 c_2 + b_4 c_4 + (b_1 c_3 - b_3 c_1 - b_4 c_3 + b_3 c_4) kx - b_3 c_3 k^2 x^2. \end{aligned}$$

The fact that $\{\text{tr } \rho(\beta \alpha^k \gamma \alpha^{-k}) \mid k \in \mathbb{Z}\}$ is a finite (resp. bounded) subset of \mathbb{C} then implies that $x = 0$ or $b_3 c_3 = ((b_1 - b_4) c_3 - b_3 (c_1 - c_4)) = 0$. If the former happens, then we are done.

Suppose now that $x \neq 0$ and that the latter happens, and that $b_3 \neq 0$ or $c_3 \neq 0$. We shall consider the case $b_3 \neq 0$; the other case will follow similarly. Note that we must then have $c_3 = 0$ and $c_1 = c_4$. Applying the same argument with γ replaced by any simple loop $\gamma' \in \Sigma_2$ such that (β, γ') is good, we conclude that $\rho|_{\Sigma_2}$ is upper triangular with image consisting of parabolic elements in $\text{SL}_2(\mathbb{C})$. If Σ_2 has genus at least one, then by considering $\text{tr } \rho(\beta \gamma')$ for good pairs (β, γ') with γ' nonseparating, we see that in fact $\rho|_{\Sigma_2}$ must have image in $\{\pm \mathbf{1}\}$. If Σ_2 has genus zero, then by our hypothesis on ρ we again conclude that $\rho|_{\Sigma_2}$ must have image in $\{\pm \mathbf{1}\}$. A fortiori, $\rho(\alpha)$ is central in both cases.

It only remains to consider the case where $b_3 = c_3 = 0$. Running through all the possible good pairs (β, γ) and repeating the above argument, we are left with the case where ρ is upper triangular (whence diagonal); then $\rho(\alpha)$ is torsion (resp. elliptic or central). This completes the proof. \square

5 Proof of the main results

The goal of this section is to complete the proof of Theorems A, B and C. This section is organized as follows. In Section 5.1 we record an irreducibility criterion for $SL_2(\mathbb{C})$ -representations of positive genus surface groups, to be used in subsequent parts of this section. In Section 5.2, we establish Theorem C for surfaces of genus one with at most two punctures. In Sections 5.3–5.5 we complete the proofs of our main theorems. Finally, in Section 5.6 we give an alternative proof of Theorem A for closed surfaces using [11; 21].

5.1 An irreducibility criterion

Lemma 5.1 *Let (a, b, c) be a sequence of loops on a surface in $(1, 2)$ -position. The following pairs are in $(1, 1)$ -position:*

$$(a, b), (a, bc), (ca, b), (ab, bc), (ca, cb), (ac, bc), (ca, ab).$$

Proof This is seen by drawing the corresponding loop configurations of homotopic clean sequences. See Figure 4, noting that any segments not passing the central basepoint are not to be considered as part of each loop configuration. \square

Let Σ be a surface of genus $g \geq 1$ with $n \geq 0$ punctures. Given a pair (a, b) of based loops in $(1, 1)$ -position on Σ , there is an embedding $\Sigma(a, b) \subset \Sigma$ of a surface of genus 1 with one boundary curve, ie a one-holed torus. Up to isotopy, every embedding of a one-holed torus is of the form $\Sigma(a, b)$ for some choice of (a, b) . The notion of loop configuration facilitates the proof of the following result, which will be used in the proof of our main theorems but may be of independent interest. (See [5] for a proof when $n = 0$.)

Proposition 5.2 *Let Σ be a surface of genus $g \geq 1$ with $n \geq 0$ punctures. A representation $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ is irreducible if and only if there is a one-holed torus subsurface $\Sigma' \subset \Sigma$ such that the restriction $\rho|_{\Sigma'}$ is irreducible.*




Figure 4: Loop configurations for Lemma 5.1.

Proof The “if” direction is clear. To prove the converse, let us begin by fixing a representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ whose restriction to every one-holed torus subsurface is reducible. In this proof, given $a \in \pi_1(\Sigma)$ we shall also denote by a the matrix $\rho(a) \in \text{SL}_2(\mathbb{C})$, for simplicity. The statement that the restriction $\rho|_{\Sigma(a, b)}$ is reducible for an embedding $\Sigma(a, b) \subset \Sigma$ associated to a pair (a, b) of loops in $(1, 1)$ -position is equivalent to saying that the pair $(\rho(a), \rho(b))$ of matrices in $\text{SL}_2(\mathbb{C})$ has a common eigenvector in \mathbb{C}^2 .

Throughout, we shall be using the following observation: if $a \in \text{SL}_2(\mathbb{C}) \setminus \{\pm \mathbf{1}\}$, and if $x, y, z \in \mathbb{C}^2$ are eigenvectors of a , then at least two of them are proportional; in notation, $x \sim y, x \sim z$ or $y \sim z$. First, we prove the following claim.

Claim Any triple (a, b, c) of loops on Σ in $(1, 2)$ -position has a common eigenvector under the representation ρ .

Proof of claim Let (a, b, c) be in $(1, 2)$ -position. Each of the pairs

$$(a, b), (a, bc), (ca, b), (ab, bc), (ca, cb), (ac, bc), (ca, ab)$$

is in $(1, 1)$ -position by Lemma 5.1, and has a common eigenvector by our hypothesis on ρ . If any of the elements a, b, ab, ca, bc are $\pm \mathbf{1}$ then (a, b, c) clearly has a common eigenvector, so we will assume otherwise in what follows. The two steps below run in parallel.

(1) Let x be a common eigenvector of (ca, b) , and let y be a common eigenvector of (ca, cb) . If $x \sim y$ then x is a common eigenvector for (a, b, c) , so let us assume otherwise. In particular, $\{x, y\}$ is an eigenbasis for ca . Since (ca, ab) has a common

eigenvector, it follows that ab has x or y as an eigenvector. If ab has x as an eigenvector, then x is a common eigenvector for (a, b, c) . So assume ab has y as an eigenvector.

Note that y is then also an eigenvector of $a^2 = ab(cb)^{-1}ca$. We may assume that $a^2 = -\mathbf{1}$, since otherwise y is also an eigenvector of a , and hence is a common eigenvector of (a, b, c) .

(2) Let x' be a common eigenvector of (a, bc) , and let y' be a common eigenvector of (ac, bc) . If $x' \sim y'$ then x' is a common eigenvector for (a, b, c) , so let us assume otherwise. In particular, $\{x', y'\}$ is an eigenbasis for bc . Since (ab, bc) has a common eigenvector, it follows that ab has x' or y' as an eigenvector. If ab has x' as an eigenvector, then x' is a common eigenvector for (a, b, c) . So assume ab has y' as an eigenvector.

Note that y' is then also an eigenvector of $b^2 = bc(ac)^{-1}ab$. We may assume that $b^2 = -\mathbf{1}$, since otherwise y' is also an eigenvector of b , and hence is a common eigenvector of (a, b, c) .

Let z be a common eigenvector of (a, b) . Since $a^2 = b^2 = -\mathbf{1}$ by the above, the eigenvalue of ab for the eigenvector z is ± 1 . Since $ab \neq \pm \mathbf{1}$ by assumption, it follows that z must be a unique eigenvector of ab , and hence $z \sim y \sim y'$. Then z is a common eigenvector of (a, b, c) . This completes the proof of our claim. □

To prove the proposition, we use the following inductive argument. First, let us assume that the image of ρ is not contained in $\{\pm \mathbf{1}\}$, since otherwise the proposition is clear. We thus can, and will, choose a nonseparating loop a_1 such that $a_1 \neq \pm \mathbf{1}$. Let us complete a_1 to an optimal sequence

$$(a_1, a_2, \dots, a_{2g-1}, a_{2g}, a_{2g+1}, \dots, a_{2g+n})$$

of generators of $\pi_1(\Sigma)$. We show that (a_1, \dots, a_{2g+n}) has a common eigenvector. For simplicity of arguments we may assume that at least one element in each pair

$$(a_1, a_2), \dots, (a_{2g-1}, a_{2g})$$

is not equal to $\pm \mathbf{1}$; for if some pair is of the form (a, b) with $a, b \in \{\pm \mathbf{1}\}$, we may simply skip over that pair in the considerations below.

If $(g, n) = (1, 0)$, then we are done since every representation $\pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ is abelian. We thus assume $(g, n) \neq (1, 0)$. By our claim, (a_1, a_2, a_3) has a common eigenvector. Assume next that $4 \leq k \leq 2g + n$ and (a_1, \dots, a_{k-1}) has a common

eigenvector $x \in \mathbb{C}^2$. We show that (a_1, \dots, a_k) has a common eigenvector. We consider the following cases.

(1) ($k = 5, 7, \dots, 2g - 1$ or $k = 2g + 1, 2g + 2, \dots, 2g + n$) The triple (a_1, a_2, a_k) is in $(1, 2)$ -position, and hence has a common eigenvector y by our claim.

Given any $3 \leq j < k$, note that $(a_1, a_2 a_j, a_k)$ is in $(1, 2)$ -position and hence has a common eigenvector by our claim, say y_j . Since $a_1 \neq \pm 1$ by assumption, we must have

$$x \sim y, \quad x \sim y_j \quad \text{or} \quad y_j \sim y.$$

If one of the first two occurs, then we conclude that (a_1, \dots, a_k) has a common eigenvector x , as required. Thus, we are left with the case $y_j \sim y$ for every $3 \leq j < k$. But this implies that $(a_1, a_2, a_2 a_j, a_k)$ has a common eigenvector y , which is thus shared also by a_j . Thus, y is a common eigenvector of (a_1, \dots, a_k) , as desired.

(2) ($k = 4, 6, \dots, 2g$) First, consider the case where $a_{k-1} = \pm 1$. It then suffices to show that the sequence $(a_1, \dots, a_{k-2}, a_k)$ has a common eigenvector. This can be shown by repeating the argument of the previous case (1) above. Thus, we may assume that $a_{k-1} \neq \pm 1$. Note that, for each integer m with $2m < k$, by our claim we have

- a common eigenvector w_m of (a_{k-1}, a_k, a_{2m}) , and
- a common eigenvector z_m of (a_{k-1}, a_k, a_{2m-1}) .

Since $a_{k-1} \neq \pm 1$, we must have

$$x \sim w_m, \quad x \sim z_m \quad \text{or} \quad w_m \sim z_m.$$

If one of the first two occurs, then we conclude that (a_1, \dots, a_k) has a common eigenvector x , and we are done. Thus, we are left with the case where $w_m \sim z_m$ for every $2m < k$. Note in this case that w_m is a common eigenvector of $(a_{2m-1}, a_{2m}, a_{k-1}, a_k)$. Comparing the vectors x , w_m and $w_{m'}$ for different m and m' , we are in turn reduced to the case $w_m \sim w_{m'}$ for all m and m' , in which case (a_1, \dots, a_k) has a common eigenvector w_1 , as desired.

Thus, (a_1, \dots, a_k) has a common eigenvector. This completes the induction, and shows that (a_1, \dots, a_{2g+n}) has a common eigenvector, proving the proposition. \square

5.2 Genus 1 with one or two punctures

We now give a separate proof of [Theorem C](#) for surfaces of genus 1 with one or two punctures. Let $\Sigma_{1,1}$ denote a surface of genus 1 with one puncture. We begin with the following:




Figure 5: Preferred generators for $\Sigma_{1,2}$.

Lemma 5.3 *Let $\rho: \pi_1(\Sigma_{1,1}) \rightarrow SL_2(\mathbb{C})$ be a semisimple representation such that each nonseparating simple loop in $\Sigma_{1,1}$ maps to an elliptic or central element of $SL_2(\mathbb{C})$. Then ρ is unitarizable, ie $\rho(\pi_1(\Sigma_{1,1}))$ has a global fixed point in \mathbb{H}^3 .*

Proof This is known in the literature; we cite two sources below.

First, the lemma can be obtained as an immediate consequence of [27, Theorem 1.2]. We sketch the connection of the setup in the current paper with that in [27]. Since every simple closed nonperipheral curve in $\Sigma_{1,1}$ maps to an elliptic element of $SL_2(\mathbb{C})$, it follows that we have $\text{tr } \rho(a) \in [-2, 2]$ for every a in the curve complex of $\Sigma_{1,1}$. Hence, in the terminology of [27], the set of end-invariants is given by the set of all projective measured laminations on $\Sigma_{1,1}$. It now follows from [27, Theorem 1.2] that $\rho(\pi_1(\Sigma_{1,1}))$ is either unitarizable or dihedral. We finish by observing that a dihedral representation $\rho: \pi_1(\Sigma) \rightarrow D_\infty \subset SL_2(\mathbb{C})$ sending every simple nonseparating simple loop to an elliptic or central element is unitarizable.

Second, a direct proof of what is essentially the contrapositive, based on an explicit presentation of the character variety $X(\Sigma_{1,1})$ (see the appendix), is given as the Algebraic Lemma in [9, Section 1.4.2]. □

We now turn to the twice-punctured torus $\Sigma_{1,2}$. We start with the following suggestive presentation of its fundamental group:

$$\pi_1(\Sigma_{1,2}) = \langle u, x, y, p_1, p_2 \mid uxy = p_1, uyx = p_2 \rangle,$$

where p_1 and p_2 denote loops around the two different punctures of $\Sigma_{1,2}$. See Figure 5 below and [12, Section 5.3].

Proposition 5.4 *Let $\rho: \pi_1(\Sigma_{1,2}) \rightarrow SL_2(\mathbb{C})$ be a semisimple representation such that each nonseparating simple loop in $\Sigma_{1,2}$ maps to an elliptic or central element of $SL_2(\mathbb{C})$. Then ρ is unitarizable.*

Combining with Lemma 4.4, we note that Lemma 5.3 and Proposition 5.4 prove Theorem C for surfaces of genus 1 with at most two punctures. Our proof of Proposition 5.4 shall rely on the following observation. In what follows, R_L denotes a hyperbolic reflection on the totally geodesic plane L in \mathbb{H}^3 .

Lemma 5.5 *Let A and B be two nontrivial elliptic rotations with distinct but intersecting axes lying on a common plane K . Then AB is an elliptic rotation that has axis lying on a plane Q that is at equal angles from K and the plane K' obtained as the image of K under A .*

Proof We note that A is a composition of two hyperbolic reflections, that is, we can write $A = R_Q \circ R_K$, where Q is a totally geodesic plane containing the axis of A , and is at half the rotation angle (of A) from K . Similarly, $B = R_K \circ R_S$, where S is a totally geodesic plane that is at half the rotation angle of B from K . Hence the composition is $AB = R_Q \circ R_S$, and its axis is the intersection of the planes Q and S . In particular, the axis lies on the plane Q , proving the lemma. \square

Proof of Proposition 5.4 Suppose that $\rho: \pi_1(\Sigma_{1,2}) \rightarrow \mathrm{SL}_2(\mathbb{C})$ is a representation satisfying the hypothesis of the proposition. We may assume that ρ is irreducible, since otherwise the result is clear.

Moreover, we may assume that the restriction of ρ to any one-holed torus subsurface of $\Sigma_{1,2}$ is semisimple. Indeed, otherwise, we may choose an optimal sequence (a_1, b_1, c_1, c_2) of generators of $\pi_1(\Sigma)$ such that the restriction of ρ to the one-holed torus $\Sigma' \subset \Sigma$ associated to the pair (a_1, b_1) is upper triangular, with the boundary loop $c = c_1 c_2$ having noncentral parabolic monodromy. The irreducibility of ρ then implies that $\rho(c_1)$ and $\rho(c_2)$ cannot be upper triangular. But then an argument as in part (b) of the proof of Lemma 4.6 shows that the restriction $\rho|_{\Sigma'}$ is reducible and moreover $\rho(a_1)$ and $\rho(b_1)$ are parabolic, whence they must both be central and $\rho(c)$ is also central; a contradiction.

Thus, in what follows, $\rho: \pi_1(\Sigma_{1,2}) \rightarrow \mathrm{SL}_2(\mathbb{C})$ is an irreducible representation satisfying the hypothesis of the proposition, with the property that its restriction to every one-holed torus subsurface is semisimple. We may further assume that none of the nonseparating simple loops have central monodromy. Indeed, suppose we have a nonseparating simple loop u such that $\rho(u)$ is central. Completing u to a sequence (u, x, y) giving a preferred presentation of $\pi_1(\Sigma_{1,2})$ above, it then suffices to show that the restriction of ρ to the subsurface $\Sigma(x, y) \subset \Sigma$ of genus 1 with one boundary curve is unitarizable, which is treated in Lemma 5.3.

Let us write $U = \rho(u)$, $X = \rho(x)$ and $Y = \rho(y)$ for the presentation of $\pi_1(\Sigma_{1,2})$ given above. By Lemma 5.3, the elements U , X and Y have coplanar, pairwise intersecting axes. We call the common plane P . Let p be the intersection point of the axes of X and Y . It suffices to prove that U fixes p as well, that is, the axis of U also passes through p . Assume that the axis of U does not pass through p ; we shall eventually reach a contradiction.

Consider the element $UXYX = UT$, say. Let $Z = X^{-1}YX$ so that $T = X^2Z$. Note that T is elliptic as it fixes the intersection of the axes of X and Y . Moreover, it can be easily checked using the loop configuration diagram that the curve represented by the element $uxyx$ is simple, closed and essential. Hence UT is also an elliptic element. However, we also have:

Claim 1 *The axis of $T = X^2Z$ does not lie on the plane P unless X is a π -rotation.*

Proof of claim Note that since $Z = X^{-1}YX$, the axis of Z is the image of the axis of Y under X^{-1} . Choose the plane K to be the one containing the axes of X and Z . Then $K = X^{-1}(P)$, ie the image of K under X is the plane P . Since X^2 and Z have the same axes as X and Z respectively, Lemma 5.5 shows that the axis of X^2Z will be on a plane that is at equal angles from K and P . In particular, the axis does not lie on the plane P unless X^2 is the identity element, that is, unless X is a π -rotation. \square

Since we have assumed that the axis of U does not pass through p , the plane P is the unique plane that contains both p and the axis of U . Note that the axis of T contains p ; hence if it does not lie in the plane P , the axes of U and T cannot lie on a common plane. By Lemma 3.4.1 of [11], we have a contradiction to the fact that UT is elliptic. Hence, X must be a π -rotation. Repeating the same argument for other presentations of $\pi_1(\Sigma_{1,2})$, we are thus reduced to the case where the monodromy trace of every nonseparating simple loop on $\Sigma_{1,2}$ under ρ is 0.

We claim that such ρ must be dihedral up to conjugacy. Indeed, let (a_1, b_1, c_1, c_2) be an optimal sequence of generators for $\pi_1(\Sigma_{1,2})$. Since

$$(\text{tr } \rho(a_1), \text{tr } \rho(b_1), \text{tr } \rho(a_1 b_1)) = (0, 0, 0),$$

which determines the restriction of ρ to $\langle a_1, b_1 \rangle$ up to conjugacy, up to global conjugation we may assume that

$$\rho(a_1) = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} \quad \text{and} \quad \rho(b_1) = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Writing

$$\rho(c_1) = \begin{bmatrix} x & y \\ z & w \end{bmatrix},$$

we have

$$\rho(b_1 c_1) = \begin{bmatrix} z & w \\ -x & -y \end{bmatrix} \quad \text{and} \quad \rho(a_1 b_1 c_1) = \begin{bmatrix} iz & iw \\ ix & iy \end{bmatrix}.$$

Since $\text{tr } \rho(b_1 c_1) = \text{tr } \rho(a_1 b_1 c_1) = 0$, the above expressions show that $y = z = 0$, ie $\rho(c_1)$ is diagonal. This shows that ρ is dihedral up to conjugacy as desired. So the hypothesis of the proposition implies that ρ is unitarizable, a contradiction. \square

5.3 Proof of Theorem A

We restate and complete the proof of Theorem A.

Theorem A *Let Σ be an oriented surface of genus $g \geq 2$ with $n \geq 0$ punctures. A semisimple representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}(2, \mathbb{C})$ has finite mapping class group orbit in the character variety $X(\Sigma)$ if and only if ρ is finite.*

Proof Suppose that $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ is a semisimple representation with finite mapping class group orbit in $X(\Sigma)$. By Lemma 3.2, we are done if ρ is reducible, so we may assume that ρ is irreducible. It suffices to show that ρ has finite monodromy along every simple closed curve around a puncture of Σ , for then we reduce to Proposition 4.3. By Lemmas 4.4 and 4.5, we see that ρ must have finite monodromy along any essential curve $a \subset \Sigma$ which is either

- nonseparating, or
- separating with each component of $\Sigma|a = \Sigma_1 \sqcup \Sigma_2$ having positive genus.

By Proposition 5.2, there is a one-holed torus subsurface $\Sigma' \subset \Sigma$ such that $\rho|_{\Sigma'}$ is irreducible. Let c be a simple closed curve around a puncture of Σ . Let $\Sigma'' \subset \Sigma$ be a two-holed torus containing Σ' and having c as one of its boundary curves; let c' be the other boundary curve of Σ'' . (Note here that each component of $\Sigma|c'$ has positive genus, by design.) We shall prove that $\rho|_{\Sigma''}$ is finite, so a fortiori ρ has finite monodromy along c . For this, we follow the strategy of [20] below.

First, we shall show that the restriction of ρ to Σ'' is unitarizable. This follows from Proposition 5.4, but for the benefit of the reader we also give another proof. First, we know from above that ρ has finite monodromy along every essential curve of Σ'' ,

as well as along the boundary curve c' . In particular, the restriction of ρ to the one-holed torus Σ' has image that is conjugate to a subgroup of $SU(2)$. Let $A \subset \mathbb{R}$ be the \mathbb{Z} -algebra generated by the set of traces of ρ along the essential curves of Σ'' . By considering the preferred generators for $\pi_1(\Sigma'')$ introduced in [Section 5.2](#), it follows from the trace relations given in [Example 2.2](#) — cf [\[12, Section 5.3\]](#) — that $\text{tr } \rho(c)$ satisfies a monic quadratic equation over the ring A , with the other root being $\text{tr } \rho(c')$. Since $\text{tr } \rho(c') \in \mathbb{R}$, it follows that $\text{tr } \rho(c_1) \in \mathbb{R}$ as well. Applying [Fact 2.4](#) to a sequence of optimal generators for $\pi_1(\Sigma'')$, we deduce that the character of $\rho|_{\Sigma''}$ is real, and since $\rho|_{\Sigma''}$ is semisimple its image is conjugate to a subgroup of $SU(2)$ or $SL_2(\mathbb{R})$; see eg [\[19, Proposition III.1.1\]](#). The latter cannot occur, since otherwise the restriction of ρ to Σ' has image conjugate to a subgroup of $SO(2)$, contradicting the fact that $\rho|_{\Sigma'}$ is irreducible, hence nonabelian. Thus, the restriction of ρ to Σ'' has image conjugate to a subgroup of $SU(2)$.

It also follows from the above analysis that the character of $\rho|_{\Sigma''}$ takes values in the ring of algebraic integers in $\overline{\mathbb{Q}}$. In particular, we may assume without loss of generality that the image of $\rho|_{\Sigma''}$ lies in $SL_2(\overline{\mathbb{Q}})$. By considering conjugates of $\rho|_{\Sigma''}$ by elements of the absolute Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ of \mathbb{Q} and noting that the above analysis goes through for all of these conjugates, we conclude that the eigenvalues of monodromy of ρ along c are algebraic integers all of whose Galois conjugates have absolute value 1. By Kronecker's theorem, it follows that the eigenvalues are roots of unity, ie ρ has finite monodromy along c (note that monodromy along ρ cannot be unipotent). This is the desired result. \square

5.4 Proof of [Theorem B](#)

We restate and complete the proof of [Theorem B](#).

Theorem B *Let Σ be an oriented surface of genus 1 with $n \geq 0$ punctures. A semisimple representation $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ has finite mapping class group orbit in the character variety $X(\Sigma)$ if and only if ρ is finite or special dihedral.*

Proof Our proof proceeds as in the proof of [Theorem A](#), with minor modifications. Suppose that $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ is a semisimple representation with finite mapping class group orbit in $X(\Sigma)$. By [Lemma 3.2](#), we are done if ρ is reducible, so we may assume that ρ is irreducible. We may also assume that ρ is not special dihedral. It suffices to show that ρ has finite monodromy along every simple closed curve around a puncture of Σ , for then we reduce to [Proposition 4.3](#).

By [Lemma 4.4](#), ρ has finite monodromy along any nonseparating essential curve $a \subset \Sigma$. Given any c which is a simple closed curve around a puncture of Σ or a separating essential curve on Σ , there is a two-holed torus subsurface of Σ having c as one of its boundary components. By the trace relations in [Example 2.2](#) (cf [[12](#), Section 5.3]), we see that $\text{tr } \rho(c)$ is an algebraic integer. It follows that $\text{tr } \rho(a)$ is an algebraic integer for every simple closed curve a on Σ . This easily implies — see eg [[20](#), Section 2.1] — that $\text{tr } \rho(\alpha)$ is an algebraic integer for every $\alpha \in \pi_1(\Sigma)$. We may in particular assume that ρ is a representation of $\pi_1(\Sigma)$ into $\text{SL}_2(\overline{\mathbb{Q}})$.

Now, ρ is unitarizable by [Theorem C](#) proved below, and in particular the monodromy eigenvalues of ρ along any curve c around a puncture of Σ have absolute value 1. Applying this observation to every conjugate of ρ by an element of the absolute Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, we see that the eigenvalues of $\rho(c)$ for any curve around a puncture of Σ are algebraic integers all of whose conjugates have absolute value 1. By Kronecker's theorem, it follows that the eigenvalues are roots of unity, ie ρ has finite monodromy along c (note that monodromy along ρ cannot be unipotent). This is the desired result. \square

5.5 Proof of [Theorem C](#)

We restate and complete the proof of [Theorem C](#).

Theorem C *Let Σ be an oriented surface of genus $g \geq 1$ with $n \geq 0$ punctures. A semisimple representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}(2, \mathbb{C})$ has bounded mapping class group orbit in the character variety $X(\Sigma)$ if and only if*

- (1) ρ is unitary up to conjugacy, or
- (2) $g = 1$ and ρ is special dihedral up to conjugacy.

Proof In the case where Σ is a surface of genus $g \geq 2$, a minor modification of the argument in the proof of [Theorem A](#) proves that ρ has elliptic or central local monodromy around the punctures. [Proposition 4.3](#) then shows that ρ is unitarizable, as desired.

Let us now assume that Σ has genus 1 with $n \geq 0$ punctures. We know the case $n \leq 2$ of [Theorem C](#) by our work in [Section 5.2](#); we shall deduce the general case from it. So suppose we have a semisimple representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ whose monodromy is central or elliptic along every nonseparating curve on Σ . It is clear that ρ is unitarizable if it is moreover reducible, so let us assume that ρ is irreducible in what follows.

By [Proposition 5.2](#), there is a one-holed torus subsurface $\Sigma' \subset \Sigma$ such that $\rho|_{\Sigma'}$ is irreducible. For any two-holed torus $\Sigma'' \subset \Sigma$ containing Σ' , [Proposition 5.4](#) shows that $\rho|_{\Sigma''}$ is unitarizable. (A fortiori, $\rho|_{\Sigma'}$ is unitarizable.) In particular, the monodromy of ρ along each of the two boundary curves of such Σ'' is central or elliptic; and hence has real trace. By considering an optimal sequence of generators for $\pi_1(\Sigma')$ and extending it to an optimal sequence of generators for $\pi_1(\Sigma)$, and applying [Fact 2.4](#), we see that the coordinate ring of the character variety $X(\Sigma)$ is generated by the monodromy traces tr_a around a finite collection of curves a each of which is either a nonseparating curve, the boundary curve of the one-holed torus Σ' , or a boundary curve of one of the two-holed tori Σ'' mentioned above. In particular, it follows that ρ has real character, and therefore the image of ρ is conjugate to a subgroup of $SU(2)$ or $SL_2(\mathbb{R})$. We claim that the latter cannot occur. Indeed, if the image of ρ lies in $SL_2(\mathbb{R})$, then unitarizability of $\rho|_{\Sigma'}$ implies that the image of $\rho|_{\Sigma'}$ is conjugate to a subgroup of $SO(2)$, contradicting the irreducibility of $\rho|_{\Sigma'}$. Thus, the image of ρ must be conjugate to a subgroup of $SU(2)$, as desired. \square

5.6 Alternative proof of [Theorem A](#) for closed surfaces

We give a different short proof of [Theorem A](#) in the case where Σ is a closed surface of genus $g \geq 2$, using the following result of Gallo, Kapovich and Marden.

Theorem 5.6 [[11](#), Section 3] *Let Σ be a closed oriented surface of genus greater than one. If $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ is nonelementary, then there exist simple loops a and b on Σ such that:*

- (1) *Their intersection number is $i(a, b) = 1$.*
- (2) *The images $\rho(a), \rho(b) \in PSL_2(\mathbb{C})$ are loxodromic and distinct, and generate a Schottky group.*

Theorem 5.7 *Let Σ be a closed orientable surface of genus $g \geq 2$. Given a semisimple representation $\pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$, the orbit of ρ in $X(\Sigma)$, under the action of $\text{Mod}(\Sigma)$, is finite if and only if the image of ρ is a finite group.*

Proof Suppose first that ρ is nonelementary. By [Theorem 5.6](#), there exist simple loops a, b on Σ with $i(a, b) = 1$ such that $\rho(a), \rho(b) \in PSL_2(\mathbb{C})$ are loxodromic and distinct. Since $\rho(a), \rho(b) \in PSL_2(\mathbb{C})$ are loxodromic, $\text{tw}_a^n(b)$ gives an infinite sequence of curves in Σ with ρ -images $\rho(a)^n \rho(b)$, whose translation length in \mathbb{H}^3 tends to infinity as $n \rightarrow \infty$, while $\text{tw}_a^n(a)$ remains fixed. It follows that the tw_a^n -orbit of ρ is infinite in $X(\Sigma)$, and hence so is the $\text{Mod}(\Sigma)$ -orbit.

Suppose that ρ is elementary. In view of the results in [Section 3](#), it remains only to treat the case where ρ is a representation whose image is a dense subgroup of $SU(2)$ in the Euclidean topology. In this case, the main theorem of [\[21\]](#) states that the $\text{Mod}(\Sigma)$ -orbit of ρ is dense in $\text{Hom}(\pi_1(\Sigma), SU(2))/SU(2)$ in the Euclidean topology. Hence it is infinite. \square

6 Applications

In this section, we collect applications of our results and methods developed in the previous sections. The following immediate corollary of [Theorem A](#) answers a question that was posed to us by Lubotzky:

Corollary 6.1 *Given a surface Σ of genus $g \geq 1$ with $n \geq 0$ punctures, a faithful representation $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ (or $\text{PSL}_2(\mathbb{C})$) cannot have finite $\text{Mod}(\Sigma)$ -orbit in the character variety.*

Proof It follows from [Theorems A and B](#) that if $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ has a finite $\text{Mod}(\Sigma)$ -orbit in $X(\Sigma)$, then ρ cannot be faithful. The same holds when $\text{SL}_2(\mathbb{C})$ is replaced by $\text{PSL}_2(\mathbb{C})$. \square

Let Σ be a closed surface of genus greater than one; we fix a hyperbolic metric on Σ such that $\Sigma = \mathbb{H}^2/\Gamma$, where Γ is a Fuchsian group. Recall that for any semisimple representation $\rho: \Gamma \rightarrow \text{PSL}_2(\mathbb{C})$, there exists a ρ -equivariant harmonic map $\tilde{h}: \mathbb{H}^2 \rightarrow \mathbb{H}^3$ from the universal cover of Σ to the symmetric space for $\text{PSL}_2(\mathbb{C})$; see eg [\[8\]](#). The *equivariant energy* of this harmonic map is the energy of its restriction to a fundamental domain of the Γ -action on \mathbb{H}^2 . The following answers a question due to Goldman.

Theorem 6.2 *Fix a Riemann surface Σ of genus greater than one with $\Gamma = \pi_1(\Sigma)$, and let $\rho: \Gamma \rightarrow \text{PSL}_2(\mathbb{C})$ be a semisimple representation. Suppose that the equivariant energies of the harmonic maps corresponding to the mapping class group orbit of ρ in $\text{Hom}(\pi_1(\Sigma), \text{PSL}_2(\mathbb{C})) // \text{PSL}_2(\mathbb{C})$ is uniformly bounded. Then $\rho(\Gamma)$ fixes a point of \mathbb{H}^3 ; in particular, $\rho(\Gamma)$ can be conjugated to lie in $\text{PSU}(2)$.*

Proof If $\rho(\Gamma)$ is elementary, but not unitary, then $\rho(\Gamma)$ fixes a geodesic in \mathbb{H}^3 , and thus the image of any equivariant harmonic map coincides with this geodesic. Let a be a simple closed curve mapped to an infinite order hyperbolic element in $\rho(\Gamma)$.

Let b be a simple closed curve on Σ with $i(a, b) > 0$. Then the translation length of $\rho(\text{tw}_a^n b)$ increases to infinity as $n \rightarrow \infty$. Hence the $\text{Mod}(\Sigma)$ -orbit of ρ cannot have bounded energy, since a uniform energy bound on the harmonic maps implies that they are uniformly Lipschitz.

Otherwise, suppose $\rho(\Gamma)$ is nonelementary. By [Theorem 5.6](#), there exist simple closed curves a, b on S such that $i(a, b) = 1$ and $\rho(a), \rho(b) \in \text{PSL}_2(\mathbb{C})$ are loxodromic and distinct. It follows as in the proof of [Theorem 5.7](#) that the translation lengths in \mathbb{H}^3 of $\rho(\text{tw}_a^n(b))$ tend to infinity, while $\text{tw}_a^n(a)$ remains fixed. Hence the \mathbb{Z} -orbit $\rho \circ \text{tw}_a^n$ of ρ under $\langle \text{tw} \rangle$ cannot have bounded energy. Thus the $\text{Mod}(\Sigma)$ -orbit of ρ cannot have bounded energy, as before.

The possibility that remains is that $\rho(\Gamma)$ is unitary. □

Appendix Case of a once-punctured torus

In this appendix, we describe how the work of Dubrovin and Mazzocco [\[9\]](#) can be used to prove the once-punctured torus case of our [Theorem B](#). We begin with the following well-known observation.

Lemma A.1 *A pair (a, b) of elements in $SL_2(\mathbb{C})$ has a common eigenvector in \mathbb{C}^2 , or in other words lies in the standard Borel B up to simultaneous conjugation, if and only if $\text{tr}([a, b]) = 2$, where $[a, b] = aba^{-1}b^{-1}$.*

Let Σ be a surface of genus 1 with one puncture. Let (a, b, c) be an optimal sequence of generators for $\pi_1(\Sigma)$, as defined in [Example 2.6](#). Let $X = X(\Sigma)$ be the character variety of Σ . Note that $\pi_1(\Sigma)$ is freely generated by a and b . The trace functions on $X(\Sigma)$ furnish an isomorphism

$$(x_1, x_2, x_3) = (\text{tr}_a, \text{tr}_b, \text{tr}_{ab}): X(\Sigma) \xrightarrow{\sim} \mathbb{A}^3$$

by Fricke; see [\[12\]](#) for details. Observing that $\text{tr}(\mathbf{1}) = 2$ for the identity $\mathbf{1} \in SL_2(\mathbb{C})$ and that $\text{tr}(A) \text{tr}(B) = \text{tr}(AB) + \text{tr}(AB^{-1})$ for any $A, B \in SL_2(\mathbb{C})$, we have

$$\begin{aligned} \text{tr}_c &= \text{tr}_{aba^{-1}b^{-1}} \\ &= \text{tr}_{aba^{-1}} \text{tr}_{b^{-1}} - \text{tr}_{aba^{-1}b} \\ &= \text{tr}_b^2 - \text{tr}_{ab} \text{tr}_{a^{-1}b} + \text{tr}_{aa} \\ &= \text{tr}_b^2 - \text{tr}_{ab}(\text{tr}_{a^{-1}} \text{tr}_b - \text{tr}_{ab}) + \text{tr}_a^2 - \text{tr}_1 \\ &= \text{tr}_a^2 + \text{tr}_b^2 + \text{tr}_{ab}^2 - \text{tr}_a \text{tr}_b \text{tr}_{ab} - 2. \end{aligned}$$

In particular, under the identification $(x_1, x_2, x_3) = (\text{tr}_a, \text{tr}_b, \text{tr}_{ab})$ of the coordinate functions above and Lemma A.1, we see that the locus of reducible representations in $X(\Sigma) = \mathbb{A}^3$ is the cubic algebraic surface cut out by the equation

$$x_1^2 + x_2^2 + x_3^2 - x_1x_2x_3 - 2 = 2.$$

The mapping class group $\text{Mod}(\Sigma)$ acts on $X(\Sigma)$ via polynomial transformations. For convenience, we shall denote the isotopy classes of simple closed curves lying in the free homotopy classes of a , b and ab by the same letters. We have the following descriptions of the associated Dehn twist actions.

Lemma A.2 *The Dehn twist actions tw_a , tw_b and tw_{ab} on $X(\Sigma)$ are given by*

$$\begin{aligned} \text{tw}_a^* &: (x_1, x_2, x_3) \mapsto (x_1, x_3, x_1x_3 - x_2), \\ \text{tw}_b^* &: (x_1, x_2, x_3) \mapsto (x_1x_2 - x_3, x_2, x_1), \\ \text{tw}_{ab}^* &: (x_1, x_2, x_3) \mapsto (x_2, x_2x_3 - x_1, x_3), \end{aligned}$$

in terms of the above coordinates.

Proof Note that $\text{tw}_a(a)$ has the homotopy class of α , $\text{tw}_a(b)$ has the homotopy class of $\alpha\beta$, and $\text{tw}_a(ab)$ has the homotopy class of $\alpha\alpha\beta$. Noting that $\text{tr}_{\alpha\alpha\beta} = \text{tr}_\alpha \text{tr}_{\alpha\beta} - \text{tr}_\beta$, we obtain the desired expression for tw_a^* . The other Dehn twists are similar. \square

Let Π be the group of polynomial automorphisms of \mathbb{A}^3 generated by tw_a^* , tw_b^* and tw_{ab}^* . It is precisely the image of the mapping class group $\text{Mod}(\Sigma)$ in the group of polynomial automorphisms of $X(\Sigma) = \mathbb{A}^3$. Let Π' be the group generated by Π together with the following transformations:

$$\begin{aligned} \sigma_{12} &: (x_1, x_2, x_3) \mapsto (-x_1, -x_2, x_3), \\ \sigma_{23} &: (x_1, x_2, x_3) \mapsto (x_1, -x_2, -x_3), \\ \sigma_{13} &: (x_1, x_2, x_3) \mapsto (-x_1, x_2, -x_3). \end{aligned}$$

It is easy to see that $[\Pi' : \Pi] < \infty$. Hence, a point in \mathbb{A}^3 has finite Π -orbit if and only if it has finite Π' -orbit. Now, the group Π' contains a group generated by transformations

$$\begin{aligned} \beta_1 &= \sigma_{12} \text{tw}_{ab}^* (\text{tw}_b^* \text{tw}_a^*)^{-1} : (x_1, x_2, x_3) \mapsto (-x_1, x_3 - x_1x_2, x_2), \\ \beta_2 &= \sigma_{23} \text{tw}_a^* (\text{tw}_b^* \text{tw}_a^*)^{-1} : (x_1, x_2, x_3) \mapsto (x_3, -x_2, x_1 - x_2x_3), \end{aligned}$$

whose finite orbits in \mathbb{A}^3 were studied by Dubrovin and Mazzocco [9, Theorem 1.6] in connection with algebraic solutions of special Painlevé VI equations. They defined

a triple $(x_1, x_2, x_3) \in \mathbb{A}^3(\mathbb{C})$ to be *admissible* if it has at most one coordinate zero and $x_1^2 + x_2^2 + x_3^2 - x_1x_2x_3 - 2 \neq 2$. It is easy to verify that the admissible points are precisely those which do not correspond to reducible or special dihedral representations. The result of [9] we shall use is the following.

Theorem A.3 (Dubrovin–Mazzocco) *A complete set of representatives for the finite (β_1, β_2) -orbits of admissible triples in \mathbb{A}^3 is given by*

$$(0, -1, -1), (0, -1, -\sqrt{2}), (0, -1, -\varphi), (0, -1, -\varphi^{-1}), (0, -\varphi, -\varphi^{-1}),$$

where $\varphi = \frac{1}{2}(1 + \sqrt{5})$ is the golden ratio.

To deduce **Theorem B** in the once-punctured torus case from the above, we recall the following explicit description of the finite subgroups BA_4, BS_4, BA_5 of $SL_2(\mathbb{C})$. First, let us identify the group of unit quaternions

$$Sp(1) = \{z = (a, b, c, d) = a + bi + cj + dk \in \mathbb{H} \mid |z| = a^2 + b^2 + c^2 + d^2 = 1\}$$

as a subgroup of $SL_2(\mathbb{C})$ by the map

$$z = (a, b, c, d) \mapsto \begin{bmatrix} a + bi & c + di \\ -c + di & a - bi \end{bmatrix}.$$

Under the identification, the *binary tetrahedral group* BA_4 is given by

$$BA_4 = \{\pm 1, \pm i, \pm j, \pm k, \frac{1}{2}(\pm 1 \pm i \pm j \pm k)\},$$

with all sign combinations taken in the above. The *binary octahedral group* BS_4 is the union of BA_4 with all quaternions obtained from $(\pm 1, \pm 1, 0, 0)/\sqrt{2}$ by all permutations of coordinates and all sign combinations. The *binary icosahedral group* BA_5 is the union of BA_4 with all quaternions obtained from $\frac{1}{2}(0, \pm 1, \pm\varphi^{-1}, \pm\varphi)$ by an even permutation of coordinates and all possible sign combinations, where $\varphi = \frac{1}{2}(1 + \sqrt{5})$ is the golden ratio.

Corollary A.4 *If $(x_1, x_2, x_3) \in \mathbb{A}^3(\mathbb{C})$ is an admissible triple with finite $\text{Mod}(\Sigma)$ -orbit, then it corresponds to a representation $\rho: \pi_1(\Sigma) \rightarrow SL_2(\mathbb{C})$ with finite image.*

Proof of the corollary Replacing (x_1, x_2, x_3) by another triple within its $\text{Mod}(\Sigma)$ -orbit if necessary, we may assume that (x_1, x_2, x_3) is one of the triples in **Theorem A.3** or its image under one of the transformations σ_{12}, σ_{23} or σ_{13} . We shall show that

$$(x_1, x_2, x_3) = (\text{tr } A, \text{tr } B, \text{tr}(AB)),$$

where $A, B \in \text{SL}_2(\mathbb{C})$ are elements that both lie in one of the finite subgroups BA_4 , BS_4 or BA_5 of $\text{SL}_2(\mathbb{C})$. Since the matrix -1 is contained in every one of these groups, it suffices to treat the case where (x_1, x_2, x_3) is one of the triples in [Theorem A.3](#). By explicit computation, we find that the triples in [Theorem A.3](#) respectively correspond to traces of the triples of matrices

$$\begin{aligned} & \left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -(1+i) & 1-i \\ -(1+i) & -(1-i) \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -(1+i) & -(1-i) \\ 1+i & -(1-i) \end{bmatrix} \right), \\ & \left(\frac{1}{\sqrt{2}} \begin{bmatrix} 0 & 1-i \\ 1+i & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -(1+i) & 1-i \\ -(1+i) & -(1-i) \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & i \\ i & -1 \end{bmatrix} \right), \\ & \left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -1 & (\varphi + \varphi^{-1}i) \\ -(\varphi - \varphi^{-1}i) & -1 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -(\varphi - \varphi^{-1}i) & -1 \\ 1 & -(\varphi + \varphi^{-1}i) \end{bmatrix} \right), \\ & \left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} 1-\varphi i & \varphi^{-1} \\ -\varphi^{-1} & -(1+\varphi i) \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -\varphi^{-1} & -(1+\varphi i) \\ -(1-\varphi i) & -\varphi^{-1} \end{bmatrix} \right), \\ & \left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -\varphi & \varphi^{-1} + i \\ -\varphi^{-1} + i & -\varphi \end{bmatrix}, \frac{1}{2} \begin{bmatrix} -\varphi^{-1} + i & -\varphi \\ \varphi & -\varphi^{-1} - i \end{bmatrix} \right), \end{aligned}$$

where $\varphi = \frac{1}{2}(1 + \sqrt{5})$ is the golden ratio. The matrices for the first triple all lie in the binary tetrahedral group BA_4 , the matrices for the second triple all lie in the binary octahedral group BS_4 , and the matrices for the remaining three triples all lie in the binary icosahedral group BA_5 . In each triple, the third matrix is the product of the first two. Thus, each of the triples in [Theorem A.3](#) correspond to representations $\pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$ with finite image, proving the corollary. □

Remark In [\[9\]](#), two proofs of [Theorem A.3](#) are given. The first proof is based on an explicit analysis of certain relevant trigonometric Diophantine equations. General equations of this type are effectively solvable by Lang’s \mathbb{G}_m conjecture (proved by Laurent [\[17\]](#)), as noted in [\[3\]](#). The second proof in [\[9\]](#), based on a suggestion of Vinberg, uses consideration of certain representations of Coxeter groups of reflections associated to admissible triples. Both methods use special features present in the once-punctured torus case which do not seem to generalize easily to the case of general surfaces treated in our work.

Acknowledgements Theorems [A](#) and [B](#) along with parts of this paper originally appeared in Chapter 6 of Whang’s PhD thesis [\[29\]](#), but the proof had a gap which has been filled in this paper. Whang thanks Peter Sarnak and Phillip Griffiths for encouragement, and Anand Patel and Ananth Shankar for collaborative work in [\[20\]](#),

which led to one of the key ingredients in this paper. Biswas and Mj thank Bill Goldman for useful email correspondence, in particular for alerting us to dihedral representations. We thank the anonymous referee for a careful reading and useful comments. Biswas and Mj are supported in part by the Department of Atomic Energy, Government of India, under project 12-R&D-TFR-5.01-0500. Research of Biswas partly supported by a DST JC Bose Fellowship. Gupta acknowledges the SERB, DST (grant MT/2017/000706) and the Infosys Foundation for their support. Mj is supported in part by an endowment of the Infosys Foundation, a DST JC Bose Fellowship, Matrices research project grant MTR/2017/000005, and CEFIPRA project 5801-1. Mj was also partially supported by the grant 346300 for IMPAN from the Simons Foundation and the matching 2015–2019 Polish MNiSW fund.

References

- [1] **I Biswas, T Koberda, M Mj, R Santharoubane**, *Representations of surface groups with finite mapping class group orbits*, New York J. Math. 24 (2018) 241–250 [MR](#) [Zbl](#)
- [2] **P Boalch**, *The fifty-two icosahedral solutions to Painlevé VI*, J. Reine Angew. Math. 596 (2006) 183–214 [MR](#) [Zbl](#)
- [3] **J Bourgain, A Gamburd, P Sarnak**, *Markoff triples and strong approximation*, C. R. Math. Acad. Sci. Paris 354 (2016) 131–135 [MR](#) [Zbl](#)
- [4] **S Cantat, F Loray**, *Dynamics on character varieties and Malgrange irreducibility of Painlevé VI equation*, Ann. Inst. Fourier (Grenoble) 59 (2009) 2927–2978 [MR](#) [Zbl](#)
- [5] **D Cooper, J F Manning**, *Nonfaithful representations of surface groups into $SL(2, \mathbb{C})$ which kill no simple closed curve*, Geom. Dedicata 177 (2015) 165–187 [MR](#) [Zbl](#)
- [6] **G Cousin**, *Algebraic isomonodromic deformations of logarithmic connections on the Riemann sphere and finite braid group orbits on character varieties*, Math. Ann. 367 (2017) 965–1005 [MR](#) [Zbl](#)
- [7] **G Cousin, V Heu**, *Algebraic isomonodromic deformations and the mapping class group*, J. Inst. Math. Jussieu 20 (2021) 1497–1545 [MR](#) [Zbl](#)
- [8] **S K Donaldson**, *Twisted harmonic maps and the self-duality equations*, Proc. Lond. Math. Soc. 55 (1987) 127–131 [MR](#) [Zbl](#)
- [9] **B Dubrovin, M Mazzocco**, *Monodromy of certain Painlevé-VI transcendents and reflection groups*, Invent. Math. 141 (2000) 55–147 [MR](#) [Zbl](#)
- [10] **R Fuchs**, *Über lineare homogene Differentialgleichungen zweiter Ordnung mit drei im Endlichen gelegenen wesentlich singulären Stellen*, Math. Ann. 63 (1907) 301–321 [MR](#) [Zbl](#)

- [11] **D Gallo, M Kapovich, A Marden**, *The monodromy groups of Schwarzian equations on closed Riemann surfaces*, Ann. of Math. 151 (2000) 625–704 MR Zbl
- [12] **W M Goldman**, *Trace coordinates on Fricke spaces of some simple hyperbolic surfaces*, from “Handbook of Teichmüller theory, II” (A Papadopoulos, editor), IRMA Lect. Math. Theor. Phys. 13, Eur. Math. Soc., Zürich (2009) 611–684 MR Zbl
- [13] **N J Hitchin**, *Poncelet polygons and the Painlevé equations*, from “Geometry and analysis” (S S Abhyankar, editor), Tata Inst. Fund. Res., Bombay (1995) 151–185 MR Zbl
- [14] **R D Horowitz**, *Characters of free groups represented in the two-dimensional special linear group*, Comm. Pure Appl. Math. 25 (1972) 635–649 MR Zbl
- [15] **K Iwasaki**, *A modular group action on cubic surfaces and the monodromy of the Painlevé VI equation*, Proc. Japan Acad. Ser. A Math. Sci. 78 (2002) 131–135 MR Zbl
- [16] **T Koberda, R Santharoubane**, *Quotients of surface groups and homology of finite covers via quantum representations*, Invent. Math. 206 (2016) 269–292 MR Zbl
- [17] **M Laurent**, *Équations diophantiennes exponentielles*, C. R. Acad. Sci. Paris Sér. I Math. 296 (1983) 945–947 MR Zbl
- [18] **O Lisovyy, Y Tykhyy**, *Algebraic solutions of the sixth Painlevé equation*, J. Geom. Phys. 85 (2014) 124–163 MR Zbl
- [19] **J W Morgan, P B Shalen**, *Valuations, trees, and degenerations of hyperbolic structures, I*, Ann. of Math. 120 (1984) 401–476 MR Zbl
- [20] **A Patel, A N Shankar, J P Whang**, *The rank two p -curvature conjecture on generic curves*, Adv. Math. 386 (2021) art. id. 107800 MR Zbl
- [21] **J P Previte, E Z Xia**, *Topological dynamics on moduli spaces, II*, Trans. Amer. Math. Soc. 354 (2002) 2475–2494 MR Zbl
- [22] **J H Przytycki, A S Sikora**, *On skein algebras and $SL_2(\mathbb{C})$ -character varieties*, Topology 39 (2000) 115–148 MR Zbl
- [23] **M van der Put, M F Singer**, *Galois theory of linear differential equations*, Grundlehr. Math. Wissen. 328, Springer (2003) MR Zbl
- [24] **K Saito**, *Character variety of representations of a finitely generated group in SL_2* , from “Topology and Teichmüller spaces” (S Kojima, Y Matsumoto, K Saito, M Seppälä, editors), World Sci., River Edge, NJ (1996) 253–264 MR Zbl
- [25] **C T Simpson**, *Moduli of representations of the fundamental group of a smooth projective variety, II*, Inst. Hautes Études Sci. Publ. Math. 80 (1994) 5–79 MR Zbl
- [26] **J C Sinz**, *Some results in representation dynamics*, PhD thesis, University of Chicago (2010) MR Available at <https://www.proquest.com/docview/825726397>
- [27] **SP Tan, Y L Wong, Y Zhang**, *End invariants for $SL(2, \mathbb{C})$ characters of the one-holed torus*, Amer. J. Math. 130 (2008) 385–412 MR Zbl

- [28] **H Vogt**, *Sur les invariants fondamentaux des équations différentielles linéaires du second ordre*, Ann. Sci. École Norm. Sup. 6 (1889) 3–71 [MR](#) [Zbl](#)
- [29] **JP Whang**, *Diophantine analysis on moduli of local systems*, PhD thesis, Princeton University (2018) [MR](#) Available at <https://www.proquest.com/docview/2057298827>
- [30] **JP Whang**, *Global geometry on moduli of local systems for surfaces with boundary*, Compos. Math. 156 (2020) 1517–1559 [MR](#) [Zbl](#)
- [31] **JP Whang**, *Nonlinear descent on moduli of local systems*, Israel J. Math. 240 (2020) 935–1004 [MR](#) [Zbl](#)

*School of Mathematics, Tata Institute of Fundamental Research
Mumbai, India*

*Department of Mathematics, Indian Institute of Science
Bangalore, India*

*School of Mathematics, Tata Institute of Fundamental Research
Mumbai, India*

*Department of Mathematics, Massachusetts Institute of Technology
Cambridge, MA, United States*

indranil@math.tifr.res.in, subhojoy@gmail.com, mahan@math.tifr.res.in,
jwhang@mit.edu

Proposed: David M Fisher

Received: 17 April 2020

Seconded: Ulrike Tillmann, Benson Farb

Revised: 3 March 2021

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITOR

András I. Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Dan Abramovich	Brown University dan_abramovich@brown.edu	Mark Gross	University of Cambridge mgross@dpnms.cam.ac.uk
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Mark Behrens	Massachusetts Institute of Technology mbehrens@math.mit.edu	Frances Kirwan	University of Oxford frances.kirwan@balliol.oxford.ac.uk
Mladen Bestvina	Imperial College, London bestvina@math.utah.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Fedor A. Bogomolov	NYU, Courant Inst., and HSE Univ., Moscow bogomolo@cims.nyu.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Martin R. Bridson	Imperial College, London m.bridson@ic.ac.uk	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	John Lott	University of California, Berkeley lott@math.berkeley.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Ralph Cohen	Stanford University ralph@math.stanford.edu	Haynes Miller	Massachusetts Institute of Technology hrm@math.mit.edu
Tobias H. Colding	Massachusetts Institute of Technology colding@math.mit.edu	Tom Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Walter Neumann	Columbia University neumann@math.columbia.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Jean-Pierre Otal	Université d'Orléans jean-pierre.otal@univ-orleans.fr
Benson Farb	University of Chicago farb@math.uchicago.edu	Peter Ozsváth	Columbia University ozsvath@math.columbia.edu
Steve Ferry	Rutgers University sferry@math.rutgers.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Ron Fintushel	Michigan State University ronfint@math.msu.edu	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David M. Fisher	Indiana University - Bloomington fisherdm@indiana.edu	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
Mike Freedman	Microsoft Research michaelf@microsoft.com	Paul Seidel	Massachusetts Institute of Technology pseidel@mit.edu
David Gabai	Princeton University gabai@princeton.edu	Peter Teichner	University of California, Berkeley teichner@math.berkeley.edu
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Richard P. Thomas	Imperial College, London richard.thomas@imperial.ac.uk
Cameron Gordon	University of Texas gordon@math.utexas.edu	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Lothar Göttsche	Abdus Salam Int. Centre for Th. Physics gottsche@ictp.trieste.it	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2022 is US \$685/year for the electronic version, and \$960/year (+ \$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 7 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2022 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 26

Issue 2 (pages 477–936)

2022

- Algebraic cobordism and étale cohomology 477
ELDEN ELMANTO, MARC LEVINE, MARKUS
SPITZWECK and PAUL ARNE ØSTVÆR
- Hilbert schemes and y -ification of Khovanov–Rozansky
homology 587
EUGENE GORSKY and MATTHEW HOGANCAMP
- Surface group representations in $SL_2(\mathbb{C})$ with finite mapping
class orbits 679
INDRANIL BISWAS, SUBHOJOY GUPTA, MAHAN MJ
and JUNHO PETER WHANG
- Deformed dimensional reduction 721
BEN DAVISON and TUDOR PĂDURARIU
- A universal Hochschild–Kostant–Rosenberg theorem 777
TASOS MOULINOS, MARCO ROBALO and BERTRAND
TOËN
- From flops to diffeomorphism groups 875
GLEB SMIRNOV
- Homogeneous Einstein metrics on Euclidean spaces are
Einstein solvmanifolds 899
CHRISTOPH BÖHM and RAMIRO A LAFUENTE