

Toeplitz operators and Hilbert modules on the symmetrized polydisc

Tirthankar Bhattacharyya*

*Department of Mathematics, Indian Institute of Science
Bengaluru 560012, Karnataka, India
tirtha@iisc.ac.in*

B. Krishna Das

*Department of Mathematics
Indian Institute of Technology Bombay
Powai, Mumbai 400076, Maharashtra, India
dasb@math.iitb.ac.in
bata436@gmail.com*

Haripada Sau

*Department of Mathematics
Indian Institute of Science Education and Research Pune
Pashan, Pune 411008, Maharashtra, India
haripadasau215@gmail.com
hsau@iiserpune.ac.in*

Received 23 July 2022

Revised 17 August 2022

Accepted 25 August 2022

Published 22 October 2022

Communicated by Yasuyuki Kawahigashi

When is the collection of S -Toeplitz operators with respect to a tuple of commuting bounded operators $S = (S_1, S_2, \dots, S_{d-1}, P)$, which has the symmetrized polydisc as a spectral set, nontrivial? The answer is in terms of powers of P as well as in terms of a unitary extension. *En route*, the Brown–Halmos relations are investigated. A commutant lifting theorem is established. Finally, we establish a general result connecting the C^* -algebra generated by the commutant of S and the commutant of its unitary extension R .

Keywords: Symmetrized polydisc; polydisc; Toeplitz operator; contractive Hilbert modules; contractive embeddings.

Mathematics Subject Classification 2020: 47A13, 47A20, 47B35, 46L07

*Corresponding author.

1. Introduction

A famous result of Brown and Halmos shows that a bounded linear operator A on the Hardy space H^2 is a Toeplitz operator if and only if $M_z^*AM_z = A$, where M_z denotes the unilateral shift. This led to the study of operators A on general Hilbert spaces satisfying an operator equation of the form $T^*AT = A$ by many authors.

Recently, the study of Toeplitz operators has been extended to the Hardy space of the symmetrized bidisc, see [2]. This motivates our study. The characterizations obtained in this paper are in terms of Hilbert modules which are the natural settings for operator theory in several variables. Completely positive maps were used by Prunaru [15] because they were well suited for the Euclidean unit ball. We use both. It is this interplay of completely positive maps with Hilbert modules which allows the results to be presented in their most general and natural form.

Let \mathbb{D} be the open unit disc while \mathbb{D}^d , $\overline{\mathbb{D}}^d$ and \mathbb{T}^d denote the open polydisc, the closed polydisc and the d -torus, respectively, in d -dimensional complex space for $d \geq 2$.

If T_1, T_2, \dots, T_d are commuting contractions on a Hilbert space \mathcal{H} and P denotes the product $T_1T_2 \dots T_d$, then an application of the arguments in [11] shows that there is a nontrivial bounded operator A on \mathcal{H} satisfying $T_i^*AT_i = A$ for all $i = 1, 2, \dots, d$ if and only if P^n does not converge to the zero operator strongly. This may remind an astute reader of [15, Corollary 2.2] where the author showed that a completely positive, completely contractive and ultraweakly continuous linear map on $\mathcal{B}(\mathcal{H})$ has a nonzero operator as a fixed point if and only if its powers at the identity operator do not converge strongly to the zero operator. Prunaru’s results were attuned to the Euclidean unit ball whereas the case of commuting contractions is attuned to the polydisc. In this paper, we discuss a third domain which has gained prominence over the last two decades.

Consider the elementary symmetric polynomials e_k for $k = 1, 2, \dots, d$ in d variables:

$$e_k(z_1, z_2, \dots, z_d) = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq d} z_{j_1} z_{j_2} \dots z_{j_k}.$$

By convention, e_0 is the constant polynomial 1. The closed symmetrized polydisc is the polynomially convex set

$$\Gamma_d = \{(e_1(\mathbf{z}), e_2(\mathbf{z}), \dots, e_d(\mathbf{z})) : \mathbf{z} = (z_1, z_2, \dots, z_d) \in \overline{\mathbb{D}}^d\}.$$

A point of Γ_d will usually be denoted by (s_1, s_2, \dots, p) . The number $d \geq 2$ will be fixed for this paper.

Let $\mathbf{S} = (S_1, \dots, S_{d-1}, P)$ be a commuting d -tuple of bounded operators on a Hilbert space \mathcal{H} . Let $\mathcal{A} = \mathbb{C}[z_1, z_2, \dots, z_d]$ be the algebra of polynomials in d commuting variables. Consider the \mathcal{A} -module structure induced on \mathcal{H} as follows:

$$f \cdot h = f(S_1, \dots, S_{d-1}, P)h \quad \text{for } f \in \mathcal{A} \quad \text{and} \quad h \in \mathcal{H}.$$

Let $\|f\|$ be the supremum norm of f over Γ_d . The \mathcal{A} -module \mathcal{H} is called a *contractive Hilbert module* if

$$\|f(S_1, \dots, S_{d-1}, P)h\| \leq \|f\| \|h\| \quad \text{for } f \in \mathcal{A} \quad \text{and} \quad h \in \mathcal{H}. \tag{1.1}$$

The polynomial convexity of Γ_d implies that (1.1) holds for every function holomorphic in a neighborhood of Γ_d . We could call such an \mathcal{H} a Γ_d -contractive Hilbert module because Γ_d is a spectral set for \mathbf{S} . However, in this paper, the set Γ_d is fixed. Hence the brevity. The commuting tuple \mathbf{S} satisfying (1.1) is also known as a Γ_d -contraction, see [4]. We shall write a Hilbert module as above as $(\mathcal{H}, \mathbf{S})$ because we shall often need the tuple \mathbf{S} . For an account of why the language of Hilbert modules is a natural one for nontrivial extension of operator theory results to several variables, see [8]. In the present context, see [16].

If $(\mathcal{K}, \mathbf{R})$ is a contractive module as above where $\mathbf{R} = (R_1, \dots, R_{d-1}, U)$ consists of normal operators and the Taylor joint spectrum $\sigma(\mathbf{R})$ is contained in the distinguished boundary $b\Gamma_d = \{(s_1, \dots, s_{d-1}, p) \in \Gamma_d : |p| = 1\}$ (see [4, Theorem 2.4(iii)]), $(\mathcal{K}, \mathbf{R})$ will be called a *unitary Hilbert module*. Let $(\mathcal{M}, \mathbf{T})$ with $\mathbf{T} = (T_1, \dots, T_{d-1}, V)$ be a submodule of a unitary Hilbert module $(\mathcal{K}, \mathbf{R})$. Then it is called an *isometric Hilbert module*. The operator tuples \mathbf{R} and \mathbf{T} are called Γ_d -unitary and Γ_d -isometry, respectively.

If $\mathbf{S}^* \stackrel{\text{def}}{=} (S_1^*, \dots, S_{d-1}^*, P^*)$, then *the adjoint module* $(\mathcal{H}, \mathbf{S}^*)$ is a contractive module when $(\mathcal{H}, \mathbf{S})$ is so. This is easy to see from the definition. Clearly, the adjoint module of a unitary module is again a unitary module. Obviously, the adjoint of an isometric module need not be an isometric module.

Given two Hilbert modules $(\mathcal{H}, \mathbf{S})$ and $(\mathcal{K}, \mathbf{R})$, a *Hilbert module homomorphism* $\mathfrak{J} : \mathcal{H} \rightarrow \mathcal{K}$ is a bounded operator \mathfrak{J} such that

$$\mathfrak{J}(f \cdot h) = f \cdot \mathfrak{J}h \quad \text{for all } f \in \mathcal{A} \quad \text{and} \quad h \in \mathcal{H}.$$

Such a homomorphism will often be called a *module map*. If, moreover, \mathfrak{J} is a contraction, we say that the module $(\mathcal{H}, \mathbf{S})$ is contractively embedded as a submodule of $(\mathcal{K}, \mathbf{R})$. A contractive module map $\mathfrak{J} : \mathcal{H} \rightarrow \mathcal{K}$ is called a *canonical module map* if $(\mathcal{K}, \mathbf{R})$ is *minimal in the sense that there is no proper submodule of $(\mathcal{K}, \mathbf{R})$ containing $(\mathcal{H}, \mathbf{S})$ and reducing \mathbf{R} and*

$$\mathfrak{J}^* \mathfrak{J} = \text{SOT-} \lim P^{*n} P^n. \tag{1.2}$$

Definition 1.1. Let $(\mathcal{H}, \mathbf{S})$ be a contractive Hilbert module. A bounded operator A on \mathcal{H} is said to be an *S-Toeplitz operator* if it satisfies the *Brown–Halmos relations* with respect to \mathbf{S} :

$$S_i^* A P = A S_{d-i} \quad \text{for each } 1 \leq i \leq d-1 \quad \text{and} \quad P^* A P = P. \tag{1.3}$$

The $*$ -closed and norm-closed vector space of all S-Toeplitz operators is denoted by $\mathcal{T}(\mathbf{S})$.

See [2, 6] for the motivation of the definition above.

Theorem 1. *Let $(\mathcal{H}, \mathcal{S})$ be a contractive Hilbert module. Then the following are equivalent:*

- (1) $\mathcal{T}(\mathcal{S})$ is nontrivial.
- (2) $Q = \text{SOT-lim } P^{*n} P^n \neq 0$.
- (3) $(\mathcal{H}, \mathcal{S})$ can be canonically embedded as a submodule of a unitary module $(\mathcal{K}, \mathcal{R})$, which is unique up to unitary isomorphism.
- (4) $(\mathcal{H}, \mathcal{S})$ can be contractively embedded as a submodule of an isometric Hilbert module.

Moreover, in this case, for any unitary module $(\mathcal{K}', \mathcal{R}')$ and any contractive embedding $\mathfrak{J}' : \mathcal{H} \rightarrow \mathcal{K}'$, the following are true:

- (i) $\mathfrak{J}'^* \mathfrak{J}' \leq \text{SOT-lim } P^{*n} P^n = \mathfrak{J}^* \mathfrak{J}$;
- (ii) $(\mathcal{K}, \mathcal{R})$ is contractively embedded into $(\mathcal{K}', \mathcal{R}')$ through a contraction $\mathfrak{T} : \mathcal{K} \rightarrow \mathcal{K}'$ such that $\mathfrak{T} \mathfrak{J} = \mathfrak{J}'$.

This theorem is proved in Sec. 2. In Theorem 1 above, $\mathcal{T}(\mathcal{S})$ depends on S_i and P whereas the condition (2) is in terms of P alone. That is the surprising strength.

The Toeplitz C^* -algebra, denoted by $C^*(I_{\mathcal{H}}, \mathcal{T}(\mathcal{S}))$, is the C^* -algebra generated by $I_{\mathcal{H}}$ and the vector space $\mathcal{T}(\mathcal{S})$ of \mathcal{S} -Toeplitz operators.

Theorem 2. *Let $(\mathcal{H}, \mathcal{S})$ be a contractive Hilbert module satisfying*

$$Q = \text{SOT-lim } P^{*n} P^n \neq 0.$$

Then $(\mathcal{H}, \mathcal{S})$ can be canonically embedded as a submodule of a unitary module $(\mathcal{K}, \mathcal{R})$ by a module map \mathfrak{J} such that:

- (1) *the map ρ defined on the commutant algebra $\{R_1, \dots, R_{d-1}, U\}'$ by*

$$\rho(Y) = \mathfrak{J}^* Y \mathfrak{J}$$

is a complete isometry onto $\mathcal{T}(\mathcal{S})$;

- (2) *there exists a surjective unital $*$ -representation*

$$\pi : C^*\{I_{\mathcal{H}}, \mathcal{T}(\mathcal{S})\} \rightarrow \{R_1, \dots, R_{d-1}, U\}'$$

such that $\pi \circ \rho = I$;

- (3) *there exists a completely contractive, unital and multiplicative mapping*

$$\Theta : \{S_1, \dots, S_{d-1}, P\}' \rightarrow \{R_1, \dots, R_{d-1}, U\}'$$

defined by $\Theta(X) = \pi(\mathfrak{J}^ \mathfrak{J} X)$ which satisfies*

$$\Theta(X) \mathfrak{J} = \mathfrak{J} X.$$

When the contractive module above is an isometric one, the following stronger version holds.

Theorem 3. *Let (\mathcal{H}, S) be an isometric Hilbert module. Then we have the following:*

- (1) *There exists a unitary module (\mathcal{K}, R) containing (\mathcal{H}, S) as an isometrically embedded submodule such that R is the minimal extension of S . In fact, \mathcal{K} is the span closure of the following elements:*

$$\{U^m h : h \in \mathcal{H} \text{ and } m \in \mathbb{Z}\}.$$

Moreover, any operator X acting on \mathcal{H} commutes with S if and only if X has a unique norm-preserving extension Y acting on \mathcal{K} commuting with R .

- (2) *An operator X is in $\mathcal{T}(S)$ if and only if there exists a unique operator Y in the commutant of the von Neumann algebra generated by $\{R_1, \dots, R_{d-1}, U\}$ such that $\|X\| = \|Y\|$ and $X = P_{\mathcal{H}}Y|_{\mathcal{H}}$.*
- (3) *Let $\mathcal{C}^*(S)$ and $\mathcal{C}^*(R)$ denote the unital \mathcal{C}^* -algebras generated by S and R , respectively, and $\mathcal{I}(S)$ denote the closed ideal of $\mathcal{C}^*(S)$ generated by all the commutators $XY - YX$ for $X, Y \in \mathcal{C}^*(S) \cap \mathcal{T}(S)$. Then, there exists a short exact sequence*

$$0 \rightarrow \mathcal{I}(S) \hookrightarrow \mathcal{C}^*(S) \xrightarrow{\pi_0} \mathcal{C}^*(R) \rightarrow 0$$

with a completely isometric cross-section, where $\pi_0 : \mathcal{C}^(S) \rightarrow \mathcal{C}^*(R)$ is the canonical unital $*$ -homomorphism which sends the generating set S to the corresponding generating set R , i.e. $\pi_0(P) = U$ and $\pi_0(S_i) = R_i$ for all $1 \leq i \leq d-1$.*

The proof of this is in Sec. 3. As an application of the above, we obtain a version of the commutant lifting theorem in Sec. 4.

Theorem 4. *Let (\mathcal{H}, S) be a contractive Hilbert module. Let $\mathfrak{J} : \mathcal{H} \rightarrow \mathcal{K}$ be a canonical embedding of (\mathcal{H}, S) as a submodule of a unitary Hilbert module (\mathcal{K}, R) . Let X be in the commutant of S . Then the module (\mathcal{H}, X) can be contractively embedded as a submodule of (\mathcal{K}, Y) for a Y in the commutant of R and $\|Y\| \leq \|X\|$.*

2. Proof of Theorem 1 and Relation to Quotient Modules

Fix a contractive Hilbert module (\mathcal{H}, S) or, in other words, a Γ_d -contraction $S = (S_1, \dots, S_{d-1}, P)$. It is well known that

$$S_i - S_{d-i}^* P = D_P F_i D_P, \quad i = 1, 2, \dots, d-1,$$

for a certain operator tuple $(F_1, F_2, \dots, F_{d-1})$. The operators F_i are called the *fundamental operators*. The existence was discovered for $d = 2$ in [3]. For a proof for $d > 2$, see [12] or [13]. Alternatively, it is easily deducible from [14, Proposition 2.5(3)].

Lemma 2.1. *For each $i = 1, 2, \dots, d-1$, we have*

$$P^{*j}(S_{d-i} - S_i^* P)P^j \rightarrow 0 \quad \text{strongly as } j \rightarrow \infty. \tag{2.1}$$

Proof. For every $h \in \mathcal{H}$, we have

$$\begin{aligned} \|P^{*j}(S_{d-i} - S_i^*P)P^j h\|^2 &= \|P^{*j}(D_P F_{d-i} D_P)P^j h\|^2 \\ &\leq \|F_{d-i}\|^2 \|D_P P^j h\|^2 \\ &= \|F_{d-i}\|^2 (\|P^j h\|^2 - \|P^{j+1} h\|^2). \end{aligned}$$

The proof follows because the last term in bracket converges to zero as $j \rightarrow \infty$. □

To prove Theorem 1, we shall take the path $(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) \Rightarrow (1)$.

Proof of $(1) \Rightarrow (2)$. Let there be a nonzero S-Toeplitz operator A . This, in particular, implies that for all $n \geq 0$ we have $A = P^{*n} A P^n$ and hence $\|Ah\| \leq \|A\| \|P^n h\|$ for every vector h . So, if P^n strongly converges to 0, then $A = 0$ which is a contradiction.

Proof of $(2) \Rightarrow (3)$. Assume $P^n \rightarrow 0$ strongly. As P is a contraction,

$$I_{\mathcal{H}} \succeq P^* P \succeq P^{*2} P^2 \succeq \dots \succeq P^{*n} P^n \succeq \dots \succeq 0.$$

This guarantees a positive nonzero contraction Q such that

$$Q = \text{SOT-}\lim P^{*n} P^n. \tag{2.2}$$

Clearly

$$P^* Q P = Q.$$

Hence, we can define an isometry $V : \overline{\text{Ran}}Q \rightarrow \overline{\text{Ran}}Q$ satisfying

$$V : Q^{\frac{1}{2}} h \mapsto Q^{\frac{1}{2}} P h \quad \text{for each } h \in \mathcal{H}. \tag{2.3}$$

We now prove that Q is an S-Toeplitz operator. Indeed

$$S_i^* Q P - Q S_{d-i} = \lim_j (S_i^* P^{*j} P^j P - P^{*j} P^j S_{d-i}) = \lim_j P^{*j} (S_i^* P - S_{d-i}) P^j.$$

By (2.1), the limit above is zero and hence $S_i^* Q P = Q S_{d-i}$. Define operators $T_j : \overline{\text{Ran}}Q \rightarrow \overline{\text{Ran}}Q$ for $j = 1, 2, \dots, d-1$ as

$$T_j : Q^{\frac{1}{2}} h \mapsto Q^{\frac{1}{2}} S_j h. \tag{2.4}$$

It is straightforward to see that each T_j is well defined. The tuple (T_1, \dots, T_{d-1}, V) is a commuting tuple too. The computation below establishes the identities $T_i = T_{d-i}^* V$ for each $i = 1, 2, \dots, d-1$:

$$\langle T_{d-i}^* V Q^{\frac{1}{2}} h, Q^{\frac{1}{2}} h' \rangle = \langle Q^{\frac{1}{2}} P h, Q^{\frac{1}{2}} S_{d-i} h' \rangle = \langle Q S_i h, h' \rangle = \langle T_i Q^{\frac{1}{2}} h, Q^{\frac{1}{2}} h' \rangle,$$

where to obtain the third equality, we use the fact that Q is an S-Toeplitz operator. We have already noted that V is an isometry. Thus, by [4, Theorem 4.12], all we need to show to conclude that (T_1, \dots, T_{d-1}, V) is a Γ_d -isometry is that the $d-1$

tuple $(\gamma_1 T_1, \dots, \gamma_{d-1} T_{d-1})$ is a Γ_{d-1} -contraction, where $\gamma_j = (d - j)/d$ for each $j = 1, 2, \dots, d - 1$. This readily follows from the identity

$$\xi(\gamma_1 T_1, \dots, \gamma_{d-1} T_{d-1}) Q^{\frac{1}{2}} = Q^{\frac{1}{2}} \xi(\gamma_1 S_1, \dots, \gamma_{d-1} S_{d-1})$$

for every polynomial ξ in $\mathbb{C}[z_1, z_2, \dots, z_{d-1}]$, and the fact that $(\gamma_1 S_1, \dots, \gamma_{d-1} S_{d-1})$ is a Γ_{d-1} -contraction where $\gamma_j = (d - j)/d$.

Let $R = (R_1, \dots, R_{d-1}, U)$ acting on \mathcal{K} be a minimal Γ_d -unitary extension of the Γ_d -isometry $T = (T_1, \dots, T_{d-1}, V)$. Define a contraction $\mathfrak{J} : \mathcal{H} \rightarrow \mathcal{K}$ as

$$\mathfrak{J} : h \mapsto Q^{\frac{1}{2}} h \quad \text{for every } h \in \mathcal{H}.$$

This is a module homomorphism because

$$R_j \mathfrak{J} h = R_j Q^{\frac{1}{2}} h = T_j Q^{\frac{1}{2}} h = Q^{\frac{1}{2}} S_j h = \mathfrak{J} S_j h. \tag{2.5}$$

Similarly $U \mathfrak{J} = \mathfrak{J} P$. Finally, by the definition of \mathfrak{J} and Q , it follows that $\mathfrak{J}^* \mathfrak{J}$ is the limit of $P^{*n} P^n$ in the strong operator topology.

For the uniqueness part, consider

$$(\mathfrak{J}, \mathcal{K}, R) = (R_1, \dots, R_{d-1}, U)$$

as above and

$$(\mathfrak{J}', \mathcal{K}', R') = (R'_1, \dots, R'_{d-1}, U')$$

with similar properties. Define the operator $\tau : \mathcal{K} \rightarrow \mathcal{K}'$ densely by

$$\tau : f(R, R^*) \mathfrak{J} h \mapsto f(R', R'^*) \mathfrak{J}' h$$

for every $h \in \mathcal{H}$ and polynomial f in z and \bar{z} . The map τ is surjective by minimality. Note that τ clearly satisfies $\tau \mathfrak{J} = \mathfrak{J}'$. We will be done if we can show that τ is an isometry. Let f be a polynomial in z and $\bar{f} f = \sum a_{\mathbf{n}, \mathbf{m}} z^{\mathbf{n}} \bar{z}^{\mathbf{m}}$, where \mathbf{n} and \mathbf{m} are multi-indices.

$$\begin{aligned} \|f(R, R^*) \mathfrak{J} h\|^2 &= \sum a_{\mathbf{n}, \mathbf{m}} \langle \mathfrak{J}^* R^{*\mathbf{m}} R^{\mathbf{n}} \mathfrak{J} h, h \rangle \\ &= \sum a_{\mathbf{n}, \mathbf{m}} \langle S^{*\mathbf{m}} \mathfrak{J}^* \mathfrak{J} S^{\mathbf{n}} h, h \rangle \\ &= \sum a_{\mathbf{n}, \mathbf{m}} \langle S^{*\mathbf{m}} Q S^{\mathbf{n}} h, h \rangle. \end{aligned} \tag{2.6}$$

Since the last term depends on S only, τ is an isometry.

Proof of (3) \Rightarrow (4). Obvious.

Proof of (4) \Rightarrow (1). Let (\mathcal{M}, T) be the isometric module with $T = (T_1, \dots, T_{d-1}, V)$. In this case, we have

$$P^* \mathfrak{J}^* \mathfrak{J} P = \mathfrak{J}^* V^* V \mathfrak{J} = \mathfrak{J}^* \mathfrak{J}$$

and

$$S_{d-j}^* \mathfrak{J}^* \mathfrak{J} P = \mathfrak{J}^* T_{d-j}^* V \mathfrak{J} = \mathfrak{J}^* T_j \mathfrak{J} = \mathfrak{J}^* \mathfrak{J} S_j.$$

This proves that the nonzero operator $\mathfrak{J}^*\mathfrak{J}$ belongs to $\mathcal{T}(\mathcal{S})$. This in particular establishes that (4) implies (1).

To complete the proof, first note that the proof of (3) \Rightarrow (1) above implies that $\mathfrak{J}'^*\mathfrak{J}'$ is an \mathcal{S} -Toeplitz operator. In particular, $\mathfrak{J}'^*\mathfrak{J}'$ is in $\mathcal{T}(P)$. This implies

$$\mathfrak{J}'^*\mathfrak{J}' = P^{*n}\mathfrak{J}'^*\mathfrak{J}'P^n \leq P^{*n}P^n \quad \text{for every } n.$$

This proves part (i).

For part (ii), we define the operator $\mathfrak{T} : \mathcal{K} \rightarrow \mathcal{K}'$ densely by

$$\mathfrak{T} : f(\mathcal{R}, \mathcal{R}^*)\mathfrak{J}h \mapsto f(\mathcal{R}', \mathcal{R}'^*)\mathfrak{J}'h$$

for every $h \in \mathcal{H}$ and polynomial f in z and \bar{z} . Using part (i), a similar computation as done in (2.6) yields

$$\|f(\mathcal{R}', \mathcal{R}'^*)\mathfrak{J}'h\| \leq \|f(\mathcal{R}, \mathcal{R}^*)\mathfrak{J}h\| \quad \text{for every } h \in \mathcal{H}.$$

This shows that \mathfrak{T} is not only well defined but also a contraction. Finally, it readily follows from the definition of \mathfrak{T} that it intertwines \mathcal{R} and \mathcal{R}' and that $\mathfrak{T}\mathfrak{J} = \mathfrak{J}'$. \square

Remark 2.2. It is known that the minimal unitary (or isometric) dilation space of a contraction acting on a Hilbert space is always infinite-dimensional even in the case of matrices. In contrast, if \mathcal{H} is finite-dimensional, the isometric module (T_1, \dots, T_{d-1}, V) defined in (2.4) acts on a finite dimensional space, viz. $\overline{\text{Ran}Q}$, and hence is Γ_d -unitary.

Biswas and Shyam Roy [4] showed that an isometric Hilbert module $(\mathcal{M}, \mathcal{T})$ always decomposes (up to unitary equivalence) as the direct sum of a pure isometric Hilbert module and a unitary Hilbert module, i.e. $\mathcal{M} = H^2(\mathcal{E}) \oplus \mathcal{K}$ where \mathcal{E} is a Hilbert space, $T_i = M_{A_i + A_{n-i}^*z} \oplus R_i$ for $i = 1, 2, \dots, d-1$ and $V = M_z \oplus U$ for certain $A_i \in \mathcal{B}(\mathcal{E})$ and $(\mathcal{K}, \mathcal{R})$ is a unitary Hilbert module, called the *unitary part* of $(\mathcal{M}, \mathcal{T})$.

If a Hilbert module $(\mathcal{H}, \mathcal{S})$ is isomorphic to $(\mathcal{M}, \mathcal{T})$ quotiented by a submodule $(\mathcal{M}', \mathcal{T}|_{\mathcal{M}'})$, then $(\mathcal{H}, \mathcal{S})$ is said to be realized as a *quotient module* or equivalently $(\mathcal{H}, \mathcal{S})$ is said to have a *co-extension* $(\mathcal{M}, \mathcal{T})$ because there is a co-isometric module map $L : \mathcal{M} \rightarrow \mathcal{H}$; see [7]. The module $(\mathcal{M}, \mathcal{T})$ is said to be *minimal* over $(\mathcal{H}, \mathcal{S})$ if there is no proper submodule of $(\mathcal{M}, \mathcal{T})$ which contains $(\mathcal{H}, \mathcal{S})$ and reduces \mathcal{T} .

Lemma 2.3. *If a contractive Hilbert module $(\mathcal{H}, \mathcal{S})$ can be realized as a quotient module of a minimal isometric module $(\mathcal{M}, \mathcal{T})$ which has a nontrivial unitary part $(\mathcal{K}, \mathcal{R})$, then there is a (not necessarily canonical) contractive embedding of the adjoint module $(\mathcal{H}, \mathcal{S}^*)$ in $(\mathcal{K}, \mathcal{R})$.*

Proof. The proof consists of defining $\mathfrak{J} : \mathcal{H} \rightarrow \mathcal{K}$ as

$$\mathfrak{J} : h \rightarrow P_{\mathcal{K}}h \quad (h \in \mathcal{H}),$$

where $P_{\mathcal{K}}$ denotes the orthogonal projection of \mathcal{M} onto \mathcal{K} . \square

Lemma 2.3 has an interesting, but not a very unexpected, consequence whose proof is easy and hence omitted.

Corollary 2.4. *If a pure contractive Hilbert module $(\mathcal{H}, \mathcal{S})$ can be realized as a quotient module of an isometric module $(\mathcal{M}, \mathcal{T})$ which is minimal over $(\mathcal{H}, \mathcal{S})$, then the unitary part in the Biswas–Shyam Roy decomposition of $(\mathcal{M}, \mathcal{T})$ is absent.*

3. Algebraic Structure of the Toeplitz C^* -Algebra

3.1. Proof of Theorem 2

We begin with a few lemmas. The first lemma below gives us the existence of an important completely positive map. This is a particular case of [15, Lemma 2.1]. The central idea of the proof goes back to Arveson, see [1, Proposition 5.2]. For a subnormal operator tuple, in the multivariable situation, Eschmeier and Everard have proven a similar result by direct construction, see [9, Sec. 3].

Lemma 3.1. *Let P be a contraction on the Hilbert space \mathcal{H} . Then there exists a completely positive, completely contractive, idempotent linear map $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ such that $\text{Ran } \Phi = \mathcal{T}(P)$. Moreover, if $A, B \in \mathcal{B}(\mathcal{H})$ satisfy $P^*(AXB)P = AP^*XPB$ for all $X \in \mathcal{B}(\mathcal{H})$ then $\Phi(AXB) = A\Phi(X)B$. In addition,*

$$\Phi(I_{\mathcal{H}}) = Q = \lim_{n \rightarrow \infty} P^{*n}P^n,$$

where the limit is in the strong operator topology.

Part of [5, Theorem 3.1] gives us more properties of Φ .

Lemma 3.2 ([5]). *Let $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ be a completely positive and completely contractive map such that $\Phi \circ \Phi = \Phi$. Then, for all X and Y in $\mathcal{B}(\mathcal{H})$, we have*

$$\Phi(\Phi(X)Y) = \Phi(X\Phi(Y)) = \Phi(\Phi(X)\Phi(Y)). \tag{3.1}$$

The last lemma that we need will play a crucial role. Its proof follows from [15, Theorem 3.1]. For someone not familiar with [15], this might create some opacity and hence a simplified proof in our context is supplied below.

Lemma 3.3. *There is an idempotent, completely positive and completely contractive map $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ such that*

$$\text{Ran } \Phi = \{X \in \mathcal{B}(\mathcal{H}) : P^*XP = X\} = \mathcal{T}(P). \tag{3.2}$$

If $(\mathcal{K}, \pi, \mathfrak{J})$ is the minimal Stinespring dilation of the restriction of Φ to the unital C^ -algebra $C^*(I_{\mathcal{H}}, \mathcal{T}(P))$ generated by $\mathcal{T}(P)$, and if $Q = \text{SOT-lim } P^{*n}P^n$, then the following properties are satisfied:*

- (P₁) $U := \pi(QP)$ is a unitary operator. Moreover, $\mathfrak{J}P = U\mathfrak{J}$ and \mathcal{K} is the smallest reducing subspace for U containing $\mathfrak{J}\mathcal{H}$.
- (P₂) The map $\rho : \{U\}' \rightarrow \mathcal{T}(P)$ defined by $\rho(Y) = \mathfrak{J}^*Y\mathfrak{J}$, for all $Y \in \{U\}'$, is surjective and a complete isometry.

(P₃) The Stinespring triple $(\mathcal{K}, \pi, \mathfrak{J})$ satisfies $\pi \circ \rho = I$. In particular,

$$\pi(C^*(I_{\mathcal{H}}, \mathcal{T}(P))) = \{U\}'.$$

(P₄) The linear map $\Theta : \{P\}' \rightarrow \{U\}'$ defined by $\Theta(X) = \pi(QX)$ is completely contractive, unital and multiplicative.

Proof. We restrict Φ to $C^*(I_{\mathcal{H}}, \mathcal{T}(P))$ and continue to call it Φ .

By definition of minimal Stinespring dilation,

$$\Phi(X) = \mathfrak{J}^* \pi(X) \mathfrak{J} \quad \text{for every } X \in C^*(I_{\mathcal{H}}, \mathcal{T}(P)). \tag{3.3}$$

Note that $Q = \Phi(I_{\mathcal{H}}) = \mathfrak{J}^* \mathfrak{J} = \text{SOT-}\lim_{n \rightarrow \infty} P^{*n} P^n$.

The kernel of Φ is an ideal in $C^*(I_{\mathcal{H}}, \mathcal{T}(P))$ by Lemma 3.2 (when Φ is allowed as a map on the whole of $\mathcal{B}(\mathcal{H})$, its kernel may not be an ideal) and hence it follows from the construction of the minimal Stinespring dilation that $\text{Ker } \Phi = \text{Ker } \pi$. Thus

$$\pi(X) = \pi(\Phi(X)) \quad \text{for any } X \in C^*(I, \mathcal{T}(P)). \tag{3.4}$$

Since π is a representation, we get $U^* \pi(X) U = \pi(X)$ for any $X \in C^*(I, \mathcal{T}(P))$.

Since π is unital, we get that U is an isometry. If P' is a projection in the weak* closure of $\pi(C^*(I, \mathcal{T}(P)))$, then we also have $U^* P' U = P'$ and $U^* P'^{\perp} U = P'^{\perp}$. This shows that $U P' = P' U$ and therefore $\pi(X) U = U \pi(X)$ for all $X \in C^*(I, \mathcal{T}(P))$. In particular, it follows that U is unitary and $\pi(C^*(I_{\mathcal{H}}, \mathcal{T}(P))) \subseteq \{U\}'$.

We now prove an identity which will be used in this proof as well as later. The identity is

$$\pi(QX) \mathfrak{J} = \mathfrak{J} X \tag{3.5}$$

for any $X \in \mathcal{B}(\mathcal{H})$ that commutes with P . The proof of (3.5) follows from two computations. For every $h, h' \in \mathcal{H}$, we have

$$\begin{aligned} \langle \pi(QX) \mathfrak{J} h, \mathfrak{J} h' \rangle &= \langle \mathfrak{J}^* \pi(QX) \mathfrak{J} h, k \rangle \\ &= \langle \Phi(QX) h, h' \rangle \\ &= \langle QX h, h' \rangle \quad (\text{because } \mathcal{T}(P) \text{ is fixed by } \Phi) \\ &= \langle \mathfrak{J} X h, \mathfrak{J} h' \rangle, \end{aligned}$$

showing that $P_{\overline{\text{Ran } \mathfrak{J}}} \pi(QX) \mathfrak{J} = \mathfrak{J} X$. On the other hand,

$$\begin{aligned} \|\pi(QX) \mathfrak{J} h\|^2 &= \langle \mathfrak{J}^* \pi(X^* Q^2 X) \mathfrak{J} h, h \rangle \\ &= \langle \Phi(X^* Q^2 X) h, h \rangle \\ &= \langle X^* \Phi(Q^2) X h, h \rangle \quad (\text{by Lemma 3.1}) \\ &= \langle X^* Q X h, h \rangle \quad (\text{by Lemma 3.2}) \\ &= \|\mathfrak{J} X h\|^2. \end{aligned}$$

Hence, (3.5) is proved.

A trivial consequence of (3.5) is that $U\mathfrak{J} = \mathfrak{J}P$. To complete the proof of **(P₁)**, it is required to establish that \mathcal{K} is the smallest reducing subspace for U containing $\mathfrak{J}\mathcal{H}$. To that end, we consider a map δ from $\text{Ran } \pi$ into $\mathcal{T}(P)$ given by

$$\delta(\pi(X)) = \mathfrak{J}^* \pi(X) \mathfrak{J} = \Phi(X) \quad \text{for all } X \in C^*(I, \mathcal{T}(P)).$$

It is injective because $\text{Ker } \Phi = \text{Ker } \pi$.

Since $\delta \circ \pi = \Phi$, we have $\delta \circ \pi$ to be idempotent and this coupled with the injectivity of δ gives us $\pi \circ \delta = I$ on $\pi\{C^*(I, \mathcal{T}(P))\}$. This immediately implies that δ is a complete isometry.

Let $\mathcal{K}_0 \subseteq \mathcal{K}$ be the smallest reducing subspace for U containing $\mathfrak{J}\mathcal{H}$. Let $P_{\mathcal{K}_0}$ be the projection in $\mathcal{B}(\mathcal{K})$ onto the space \mathcal{K}_0 . Consider the vector space

$$P_{\mathcal{K}_0}\{U\}'P_{\mathcal{K}_0} := \{P_{\mathcal{K}_0}XP_{\mathcal{K}_0} : X \in \{U\}'\} = \{P_{\mathcal{K}_0}X|_{\mathcal{K}_0} \oplus 0_{\mathcal{K}_0^\perp} : X \in \{U\}'\}$$

and the map $\delta' : P_{\mathcal{K}_0}\{U\}'P_{\mathcal{K}_0} \rightarrow \mathcal{T}(P) \subseteq \mathcal{B}(\mathcal{H})$ defined by $X \mapsto \mathfrak{J}^*X\mathfrak{J}$. This is injective.

Indeed, it is easy to check that $\mathfrak{J}^*X\mathfrak{J} \in \mathcal{T}(P)$ for $X \in \{U\}'$. Now if $\mathfrak{J}^*X\mathfrak{J} = 0$ for some $X \in \{U\}'$ then using the identity $\mathfrak{J}P = U\mathfrak{J}$, we get that

$$\langle Xf(U, U^*)\mathfrak{J}h, g(U, U^*)\mathfrak{J}k \rangle = 0$$

for any two variable polynomials f and g and $h, k \in \mathcal{H}$. This shows that $P_{\mathcal{K}_0}XP_{\mathcal{K}_0} = 0$ and therefore, δ' is injective. For any $Y \in P_{\mathcal{K}_0}\{U\}'P_{\mathcal{K}_0}$,

$$\delta'(P_{\mathcal{K}_0}\pi(\mathfrak{J}^*Y\mathfrak{J})P_{\mathcal{K}_0} - Y) = \mathfrak{J}^*\pi(\mathfrak{J}^*Y\mathfrak{J})\mathfrak{J} - \mathfrak{J}^*Y\mathfrak{J} = \Phi(\mathfrak{J}^*Y\mathfrak{J}) - \mathfrak{J}^*Y\mathfrak{J} = 0.$$

Thus, by the injectivity of δ' , we have $P_{\mathcal{K}_0}\pi(C^*(I, \mathcal{T}(P)))P_{\mathcal{K}_0} = P_{\mathcal{K}_0}\{U\}'P_{\mathcal{K}_0}$. In other words, we have a surjective complete contraction

$$\tilde{C}_{\mathcal{K}_0} : \pi(C^*(I, \mathcal{T}(P))) \rightarrow P_{\mathcal{K}_0}\{U\}'P_{\mathcal{K}_0} = \{P_{\mathcal{K}_0}X|_{\mathcal{K}_0} \oplus 0_{\mathcal{K}_0^\perp} : X \in \{U\}'\},$$

defined by $X \mapsto P_{\mathcal{K}_0}XP_{\mathcal{K}_0}$. Since $\delta = \delta' \circ \tilde{C}_{\mathcal{K}_0}$ and δ is a complete isometry, $\tilde{C}_{\mathcal{K}_0}$ is a complete isometry. Then the induced compression map

$$C_{\mathcal{K}_0} : \pi(C^*(I, \mathcal{T}(P))) \rightarrow \{P_{\mathcal{K}_0}U|_{\mathcal{K}_0}\}' \subseteq \mathcal{B}(\mathcal{K}_0), \quad X \mapsto P_{\mathcal{K}_0}X|_{\mathcal{K}_0},$$

is a unital complete isometry and therefore a C^* -isomorphism by a result of Kadison [10]. Hence, by the minimality of the Stinespring representation π we have $\mathcal{K} = \mathcal{K}_0$ and therefore $\pi(C^*(I, \mathcal{T}(P))) = \{U\}'$. This completes the proofs of **(P₁)**, **(P₂)** and **(P₃)**.

To prove **(P₄)**, first note that Θ is completely contractive and unital as $\pi(Q) = I$. We have also proved that $\Theta(X)\mathfrak{J} = \mathfrak{J}X$ for all $X \in \{P\}'$. Since, for $X, Y \in \{P\}'$,

$$\delta(\Theta(XY) - \Theta(X)\Theta(Y)) = \mathfrak{J}^*\mathfrak{J}XY - \mathfrak{J}^*\Theta(X)\Theta(Y)\mathfrak{J} = 0,$$

then by injectivity of δ , we have Θ to be multiplicative. Hence **(P₄)** is proved. \square

With these properties of the Stinespring dilation of Φ at hand, we start the proof of the theorem. Define

$$R_i := \pi(QS_i) \quad \text{for } 1 \leq i \leq d-1 \quad \text{and} \quad U = \pi(QP).$$

We first show that $(\mathcal{K}, \mathcal{R})$ is a unitary module. To that end, we shall use [4, Theorem 4.2]. Let $\gamma_i = (d-i)/d$ for $i = 1, 2, \dots, d-1$. Since Θ in (\mathbf{P}_4) is multiplicative, the tuple (R_1, \dots, R_{d-1}, U) is commuting. Note that $\pi \circ \Phi(X) = \pi(X)$, for every $X \in \mathcal{C}^*(I_{\mathcal{H}}, \mathcal{T}(P))$, which follows from the facts that $\rho \circ \pi(X) = \Phi(X)$ for all $X \in \mathcal{C}^*(I_{\mathcal{H}}, \mathcal{T}(P))$ and $\pi \circ \rho = I$.

Also note that for each $1 \leq i \leq d-1$,

$$\begin{aligned} R_i^*U &= \pi(S_i^*Q^2P) = \pi(\Phi(S_i^*Q^2P)) = \pi(S_i^*\Phi(Q^2)P) = \pi(S_i^*QP) = \pi(QS_{d-i}) \\ &= R_{d-i}. \end{aligned}$$

It remains to show that the tuple $(\gamma_1 R_1, \dots, \gamma_{d-1} R_{d-1})$ is a Γ_{d-1} -contraction. Since (S_1, \dots, S_{d-1}, P) is a Γ_d -contraction, by [4, Lemma 2.7], $(\gamma_1 S_1, \dots, \gamma_{d-1} S_{d-1})$ is a Γ_{d-1} -contraction. Since Θ in (\mathbf{P}_4) is multiplicative, $(\gamma_1 R_1, \dots, \gamma_{d-1} R_{d-1})$ is also a Γ_{d-1} -contraction. It follows that the tuple (R_1, \dots, R_{d-1}, U) is Γ_d -unitary.

That $(\mathcal{H}, \mathcal{S})$ is canonically embedded as a submodule of the unitary module $(\mathcal{K}, \mathcal{R})$ by \mathfrak{J} follows from the operator identity $\pi(QX)\mathfrak{J} = \mathfrak{J}X$ for every $X \in \{P\}'$ which was proved in the paragraphs following (3.5).

Minimality follows from (\mathbf{P}_1) , which says that \mathcal{K} is actually equal to

$$\overline{\text{span}}\{U^m \mathfrak{J}h : h \in \mathcal{H} \text{ and } m \in \mathbb{Z}\}.$$

Let ρ be as in (\mathbf{P}_2) above. Consider the restriction of ρ to $\{R_1, \dots, R_{d-1}, U\}'$ and continue to denote it by ρ . Since complete isometry is a hereditary property, to prove part (1), all we have to show is that $\rho(Y)$ lands in $\mathcal{T}(\mathcal{S})$, whenever Y is in $\{R_1, \dots, R_{d-1}, U\}'$ and ρ is surjective. To that end, let $Y \in \{R_1, \dots, R_{d-1}, U\}'$. Then for each $i = 1, 2, \dots, d-1$, we see that

$$\begin{aligned} S_i^* \rho(Y)P &= S_i^* \mathfrak{J}^* Y \mathfrak{J} P = \mathfrak{J}^* R_i^* Y U \mathfrak{J} = \mathfrak{J}^* R_{d-i} Y \mathfrak{J} \\ &= \mathfrak{J}^* Y R_{d-i} \mathfrak{J} = \mathfrak{J}^* Y \mathfrak{J} S_{d-i} = \rho(Y) S_{d-i}. \end{aligned}$$

Thus ρ maps $\{R_1, \dots, R_{d-1}, U\}'$ into $\mathcal{T}(\mathcal{S})$. For proving surjectivity of ρ , pick $X \in \mathcal{T}(\mathcal{S})$. Then by (\mathbf{P}_2) above, there exists a Y in $\{U\}'$ such that $\rho(Y) = \mathfrak{J}^* Y \mathfrak{J} = X$. We have to show that Y commutes with each R_i . Since X is in $\mathcal{T}(\mathcal{S})$, we have $S_i^* \mathfrak{J}^* Y \mathfrak{J} P = \mathfrak{J}^* Y \mathfrak{J} S_{d-i}$. Therefore, by the intertwining property of \mathfrak{J} we have $\mathfrak{J}^* R_i^* Y U \mathfrak{J} = \mathfrak{J}^* Y R_{d-i} \mathfrak{J}$, which is the same as $\mathfrak{J}^* R_{d-i}^* Y \mathfrak{J} = \mathfrak{J}^* Y R_{d-i} \mathfrak{J}$. This implies that for each $i = 1, 2, \dots, d-1$,

$$\rho(Y R_{d-i} - R_{d-i} Y) = \mathfrak{J}^* (Y R_{d-i} - R_{d-i} Y) \mathfrak{J} = 0.$$

This establishes the commutativity of Y with each R_i , since ρ is an isometry. This completes the proof of part (1).

Part (2) of the theorem follows from the content of (\mathbf{P}_3) if we restrict π to $C^*(I, \mathcal{T}(\mathbf{S}))$ and continue to call it π .

For the last part of theorem, let us take Θ as in (\mathbf{P}_4) , i.e.

$$\Theta(X) = \pi(QX)$$

for every X in $\{P\}'$. Restrict Θ to $\{S_1, \dots, S_{d-1}, P\}'$ and continue to call it Θ . The aim is to show that $\Theta(X) \in \{R_1, \dots, R_{d-1}, U\}'$ if $X \in \{S_1, \dots, S_{d-1}, P\}'$. For this, we first observe that if X commutes with each S_j , then QX is in $\mathcal{T}(\mathbf{S})$. Now the rest of the proof follows from part (2) of the theorem and (3.5).

3.2. Proof of Theorem 3

Let $Q, \mathfrak{J}, \pi, \rho$ and $\mathbf{R} = (R_1, \dots, R_{d-1}, U)$ be as in Theorem 2. We first note that \mathfrak{J} is an isometry because $\mathfrak{J}^*\mathfrak{J} = Q = \text{SOT} - \lim_j P^{*j}P^j = I$. We shall identify the module $(\mathcal{H}, \mathbf{S})$ with the submodule $(\mathfrak{J}\mathcal{H}, \mathfrak{J}\mathbf{S}\mathfrak{J}^*)$ of $(\mathcal{K}, \mathbf{R})$. Thus, we get from part (1) of Theorem 2 that $(\mathcal{K}, \mathbf{R})$ is a minimal unitary extension of $(\mathcal{H}, \mathbf{S})$. Now let X be an operator on \mathcal{H} which commutes with \mathbf{S} . Set $Y := \pi(X)$. Then by part (3) of Theorem 2, it follows that Y commutes with \mathbf{R} and $Y|_{\mathcal{H}} = X$, i.e. Y is an extension of X . Also since the map ρ , as in part (2) of Theorem 2, is an isometry, Y is a unique norm-preserving extension of X . Thus part (1) follows.

Part (2) follows straightforward by setting $Y := \pi(X)$ and remembering that π is the minimal Stinespring dilation of the CP map $\Phi_0 = \Phi|_{C^*(I_{\mathcal{H}}, \mathcal{T}(P))} : C^*(I_{\mathcal{H}}, \mathcal{T}(P)) \rightarrow \mathcal{B}(\mathcal{H})$ where $\Phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ is an idempotent CP map with range $\mathcal{T}(P)$.

To prove part (3), set π_0 to be the restriction of π to $C^*(\mathbf{S})$. The representation π maps the generating set \mathbf{S} of $C^*(\mathbf{S})$ to the generating set \mathbf{R} of $C^*(\mathbf{R})$. Since $\pi_0(\mathbf{S}) = \mathbf{R}$, the range of π_0 is $C^*(\mathbf{R})$. Therefore, to prove that the following sequence

$$0 \rightarrow \mathcal{I}(\mathbf{S}) \hookrightarrow C^*(\mathbf{S}) \xrightarrow{\pi_0} C^*(\mathbf{R}) \rightarrow 0$$

is a short exact sequence, all we need to show is that $\ker \pi_0 = \mathcal{I}(\mathbf{S})$.

Since $\pi_0(C^*(\mathbf{S}))$ is abelian, we have $XY - YX$ in the kernel of π_0 , for any $X, Y \in C^*(\mathbf{S}) \cap \mathcal{T}(\mathbf{S})$. Hence $\mathcal{I}(\mathbf{S}) \subseteq \ker \pi_0$. To prove the other inclusion, let Z_1 be a finite product of members of \mathbf{S}^* and Z_2 be a finite product of members of \mathbf{S} and call $Z = Z_1Z_2$. Since $Z \in \mathcal{T}(\mathbf{S}) \subseteq \mathcal{T}(P)$, we have $\Phi_0(Z) = Z$. Note that $\Phi_0(Z) = P_{\mathcal{H}}\pi_0(Z)|_{\mathcal{H}}$, for every $Z \in C^*(\mathbf{S})$. Now let Z be any arbitrary finite product of members from \mathbf{S} and \mathbf{S}^* . Since $\pi_0(\mathbf{S}) = \mathbf{R}$, which is a family of normal operators, we obtain, by the Fuglede–Putnam Theorem, that action of Φ_0 on Z has all the members from \mathbf{S}^* at the left and all the members from \mathbf{S} at the right. It follows from $\ker \pi = \ker \Phi_0$ and idempotence of Φ_0 that $\ker \pi_0 = \{Z - \Phi_0(Z) : Z \in C^*(\mathbf{S})\}$.

Because of the above action of Φ_0 , if Z is a finite product of elements from \mathbf{S} and \mathbf{S}^* then a simple commutator manipulation shows that $Z - \Phi_0(Z)$ belongs to the ideal generated by all the commutators $XY - YX$, where $X, Y \in C^*(\mathbf{S}) \cap \mathcal{T}(\mathbf{S})$. This shows that $\ker \pi_0 = \mathcal{I}(\mathbf{S})$.

In order to find a completely isometric cross-section, set $\rho_0 := \rho|_{\pi(\mathcal{C}^*(\mathcal{S}))}$. Then by the definition of ρ and the action of Φ_0 , it follows that $\rho_0(\pi(X)) = \mathfrak{J}^* \pi(X) \mathfrak{J} = \Phi_0(X) \in \mathcal{C}^*(\mathcal{S})$ for all $X \in \mathcal{C}^*(\mathcal{S})$. Thus, $\text{Ran } \rho_0 \subseteq \mathcal{C}^*(\mathcal{S})$ and therefore is a completely isometric cross-section. This completes the proof of the theorem.

4. An Application

In this section, we prove Theorem 4. Let Q be the limit as in (2.2). Consider the isometric module $(\overline{\text{Ran}Q}, \mathbb{T})$ constructed in the proof of Theorem 1, see Eqs. (2.3) and (2.4). We shall obtain a bounded operator \tilde{X} acting on $\overline{\text{Ran}Q}$ with the following properties:

- (1) \tilde{X} would commute with $\mathbb{T} = (T_1, \dots, T_{d-1}, V)$; and
- (2) $\|\tilde{X}\| \leq \|X\|$.

We shall then apply the commutant extension theorem established in part (1) of Theorem 3.

To that end, we first do a simple inner product computation. For every $h \in \mathcal{H}$,

$$\|Q^{\frac{1}{2}}Xh\|^2 = \langle X^*QXh, h \rangle = \lim_n \langle P^{*n}X^*XP^n h, h \rangle \leq \|X\|^2 \langle Qh, h \rangle.$$

Thus, there is a bounded operator $\tilde{X} : \overline{\text{Ran}Q} \rightarrow \overline{\text{Ran}Q}$ such that

$$\tilde{X} : Q^{\frac{1}{2}}h \mapsto Q^{\frac{1}{2}}Xh,$$

with norm at most $\|X\|$. Let $j = 1, 2, \dots, d-1$ and let T_j be the operators defined in (2.4). Then for each $h \in \mathcal{H}$, we have

$$\tilde{X}T_jQ^{\frac{1}{2}}h = \tilde{X}Q^{\frac{1}{2}}S_jh = Q^{\frac{1}{2}}XS_jh = Q^{\frac{1}{2}}S_jXh = T_jQ^{\frac{1}{2}}Xh = T_j\tilde{X}Q^{\frac{1}{2}}h,$$

showing that \tilde{X} commutes with T_j for all $j = 1, \dots, d-1$. A similar computation also establishes that $\tilde{X}V = V\tilde{X}$. We are now ready to apply the technique of commutant extension mentioned above.

Consider $\mathbb{R} = (R_1, \dots, R_{d-1}, U)$ acting on \mathcal{K} as in (2.5). Recall that $(\mathcal{K}, \mathbb{R})$ is a unitary module and there is a contraction $\mathfrak{J} : \mathcal{H} \rightarrow \mathcal{K}$ defined as $\mathfrak{J}h = Q^{\frac{1}{2}}h$ such that (2.5) holds. Moreover, now by the part of item (1) in Theorem 3, there exists an operator Y in the commutant of \mathbb{R} such that $Y|_{\overline{\text{Ran}Q}} = \tilde{X}$ and $\|Y\| = \|\tilde{X}\| \leq \|X\|$. Finally, we note that for every $h \in \mathcal{H}$,

$$\mathfrak{J}Xh = Q^{\frac{1}{2}}Xh = \tilde{X}Q^{\frac{1}{2}}h = YQ^{\frac{1}{2}}h = Y\mathfrak{J}h.$$

This completes the proof.

The following analog of the intertwining lifting theorem is easily obtained as a corollary to Theorem 4 by a 2×2 operator matrix trick.

Corollary 4.1. *Let (\mathcal{H}, S) and (\mathcal{H}', S') be two contractive modules. Let $(\mathfrak{J}, \mathcal{K}, R)$ and $(\mathfrak{J}', \mathcal{K}', R')$ be as obtained by part (3) of Theorem 1 corresponding to (\mathcal{H}, S) and (\mathcal{H}', S') , respectively. Then, corresponding to any operator $X : \mathcal{H} \rightarrow \mathcal{H}'$ intertwining S and S' , there exists another operator $Y : \mathcal{K} \rightarrow \mathcal{K}'$ such that Y intertwines R and R' , $Y\mathfrak{J} = \mathfrak{J}'X$ and $\|Y\| \leq \|X\|$.*

Conflict of Interest

The authors confirm that there is no conflict of interest. No datasets were generated or analyzed during this study.

Acknowledgments

The research work of Tirthankar Bhattacharyya is supported by the J C Bose National Fellowship JCB/2021/000041, and those of B. Krishna Das and Haripada Sau are supported by the DST-INSPIRE Faculty Fellowships DST/INSPIRE/04/2015/001094 and DST/INSPIRE/04/2018/002458, respectively.

References

- [1] W. Arveson, Interpolation problems in nest algebras, *J. Funct. Anal.* **20** (1975) 208–233.
- [2] T. Bhattacharyya, B. K. Das and H. Sau, Toeplitz operators on the symmetrized bidisc, *Int. Math. Res. Not.* **2021**(11) (2021) 8492–8520.
- [3] T. Bhattacharyya, S. Pal and S. Shyam Roy, Dilations of Γ -contractions by solving operator equations, *Adv. Math.* **230** (2012) 577–606.
- [4] S. Biswas and S. Shyam Roy, Functional models of Γ_n -contractions and characterization of Γ_n -isometries, *J. Funct. Anal.* **266** (2014) 6224–6255.
- [5] M. D. Choi and E. G. Effros, Injectivity and operator spaces, *J. Funct. Anal.* **24** (1977) 156–209.
- [6] B. K. Das and H. Sau, Algebraic properties of Toeplitz operators on the symmetrized polydisk, *Complex Anal. Oper. Theory* **15** (2021) 60.
- [7] R. G. Douglas, G. Misra and J. Sarkar, Contractive Hilbert modules and their dilations, *Isr. J. Math.* **187** (2012) 141–165.
- [8] R. G. Douglas and V. I. Paulsen, *Hilbert Modules Over Function Algebras*, Pitman Research Notes in Mathematics Series, Vol. 217 (Longman Scientific & Technical, Harlow, 1989) (co-published in the United States with John Wiley & Sons, Inc., New York, 1989).
- [9] J. Eschmeier and K. Everard, Toeplitz projections and essential commutants, *J. Funct. Anal.* **269** (2015) 1115–1135.
- [10] R. V. Kadison, Isometries of operator algebras, *Ann. Math.* **54** (1951) 325–338.
- [11] P. S. Muhly, Toeplitz operators and semigroups, *J. Math. Anal. Appl.* **38** (1972) 312–319.
- [12] A. Pal, On Γ_n -contractions and their conditional dilations, preprint, arXiv:1704.04508 [math.FA].

- [13] S. Pal, Dilation, functional model and a complete unitary invariant for C_0 Γ_n -contractions, preprint, arXiv:1708.06015 [math.FA].
- [14] S. Pal, Canonical decomposition of operators associated with the symmetrized polydisc, *Complex Anal. Oper. Theory* **12**(4) (2018) 931–943.
- [15] B. Prunaru, Toeplitz operators associated to commuting row contractions, *J. Funct. Anal.* **254**(6) (2008) 1626–1641.
- [16] J. Sarkar, Operator theory on symmetrized bidisc, *Indiana Univ. Math. J.* **64**(3) (2015) 847–873.